THO)UKE
IOSPITAL

LIBRARY
PERMANENT LOAN
GEORGIA MEDICAL SOCIETY

Duke University Medical Center Library
Historical Collection

Digitized by the Internet Archive
in 2016

https://archive.org/details/systemofanatomyp11fyfe

A

SYSTEM

OF

ANATOMY,

PHYSIOLOGY, &c.

Entered in Stationers Hall.



Alexander Monro Senr. M. D.

Professor of Anatomy, Fellow of the

Royal College of Physicians at Edinburgh and F.R.S.

A

SYSTEM

OF

ANATOMY

AND

PHYSIOLOGY,

WITH THE

COMPARATIVE ANATOMY OF ANIMALS.

COMPILED FROM

THE LATEST AND BEST AUTHORS,

AND ARRANGED,

AS NEARLY AS THE WORK WOULD ADMIT,

IN THE ORDER OF THE

LECTURES

DELIVERED IN THE

UNIVERSITY OF EDINBURGH.

IN THREE VOLUMES.

With a complete Index to each Volume.

A NEW EDITION,

WITH ADDITIONS, CORRECTIONS, AND NOTES;

ILLUSTRATED WITH

TWENTY NEW COPPERPLATES.

And an elegantly engraved head of

DR ALEXANDER MONRO, SENIOR.

VOL. I.

EDINBURGH:

PRINTED BY ALEX. SMELLIE, ANCHOR CLOSE;

FOR WILLIAM CREECH.

AND SOLD BY G. G. AND J. ROBINSON,

LONDON.

1801.

I
Edinburgh
V.1

TO

ALEXANDER MONRO, M. D.

FELLOW OF THE ROYAL SOCIETY,

AND OF THE COLLEGE OF PHYSICIANS,

AND PROFESSOR OF PHYSIC, ANATOMY, AND SURGERY,

IN THE UNIVERSITY OF EDINBURGH.

SIR,

THE great number and variety of books neceſſary for Students in Anatomy, many of which had become extremely ſcarce and valuable, rendered an Anatomical library very expenſive. In order to remove ſo great an inconvenience, a Gentleman connected with this Univerſity, whoſe great Anatomical knowledge and accuracy fully qualified him for the taſk, compiled from the beſt Authors the following Syſtem, which has already gone through three Editions, in the ſpace of a very few years.

MR CREECH, whoſe indefatigable zeal for promoting literature, and the intereſts of this Univerſity, is well known, bought the copy-right of this Work; and the urgent demand for a new Edition induced him to put each of the volumes under

21997

der different preſſes, leſt, at the commencement of your Lectures, the Students might be diſappointed in not having aſſiſtance from this uſeful Compilation. Being unable, from the multiplicity of his buſineſs, to ſuperintend the Publication, he requeſted me to reviſe the Work for the preſs. As I was acquainted with the ſubject, and the Authors whence the ſelection was made, I readily undertook the taſk.

In the firſt volume, which contains chiefly the Oſteology by the late Profeſſor, your illuſtrious father and the founder of the Medical School here, little more was requiſite than to correct ſome typographical errors, and to adapt a few phraſes to the idiom of the Engliſh language, in its preſent improved ſtate.

In the other volumes, where phyſiology was intermixed with anatomical deſcription, the caſe was widely different. This circumſtance, however, happened from no fault of the original Editor or Projector of the Work; for he had choſen the beſt books to ſelect from, and his knowledge and diſcrimination, in this reſpect, are ſufficiently conſpicuous; but theſe works being in foreign languages, and an Engliſh book being wanted, unfortunately erroneous tranſlations had been uſed. The Editor, who directed the choice of the books, could

could not be ſuppoſed, amid the multifarious and accumulated duties of his office, to attend to the faithfulneſs of the tranſlations; and the Book came before the Public therefore in a very imperfect ſtate.

THESE defects I have endeavoured to remedy in this Edition; and I am not conſcious, that any errors have eſcaped my notice. Should any, however, be diſcovered, I truſt that your candour will attribute them to their true cauſe, viz. the hurry of correcting three large octavo volumes for three preſſes, and preparing tranſlations for two of them in the ſpace of a few weeks.

THE thanks of the Public are juſtly due to you for patroniſing, and to the Editor for projecting, ſo uſeful a work. The faults of the former Editions were unavoidable from the circumſtances above mentioned; and if, by amending them, I have any way contributed to render the work more worthy of your patronage, or more uſeful to the attendants at your crowded theatre, I ſhall think myſelf highly honoured.

I have the honour to be,
With the greateſt reſpect,
SIR,
Your moſt obedient humble ſervant,

JOHN ROTHERAM.

ADVERTISEMENT

TO THIS NEW EDITION,

BY

THE PUBLISHER.

THE firſt Volume of this work contains the Oſteology of Dr Alexander Monro, ſenior, a performance which has been tranſlated and publiſhed in the moſt ſplendid form in foreign countries. The publiſher being in poſſeſſion of the beautifully engraved head of the Author, by Baſire from the original painting of Ramſay, ſo much celebrated by Mr Lavater in his ingenious Book on Phyſiognomy, thought it might not be unacceptable to the public, to be preſented with ſo elegant an engraving of the father of the Medical School of Edinburgh.

IT is therefore given as an embelliſhment to this new edition of the Syſtem of Anatomy and Phyſiology, which contains ſo much of his valuable writings.

THE words of Lavater, on ſeeing an impreſſion from this very plate, ſhall here be added;

and thoſe who knew the man beſt, will teſtify to the juſtneſs of the character drawn of Dr Monro.

Mr Lavater ſays,

" A good, gentle, and peaceable character, of a " ſanguine phlegmatic temperament. The good- " neſs is depicted in his eyes; the mouth breathes " only peace; and an amiable ſerenity is diffuſ- " ed over the whole countenance. This man is " incapable of giving offence to any one, and " who could ever ſuffer himſelf deſignedly to " offend him?—He loves tranquillity, order, " and ſimple elegance. He takes a clear view " of the ſubject he examines; he thinks accu- " rately; his ideas, and his reaſonings, are al- " ways equally well followed up; his mind re- " jects all that is falſe or obſcure. He gives " with a liberal hand, he forgives with a ge- " nerous heart, and takes delight in ſerving his " fellow creatures. You may ſafely depend on " what he ſays, on what he promiſes. His " ſenſibility never degenerates into weakneſs; " he eſteems worth, find it where he may. He " is not indifferent to the pleaſures of life; but " ſuffers not himſelf to be enervated by them. " This is not what is uſually denominated a

" great

" great man ; but he poſſeſſes a much more exalt-
" ed character ;—he is the honour of humanity,
" and of his rank in life.

" Reſpectable perſonage, I know you not ;
" I am entirely in the dark concerning you ;
" —but you ſhall not eſcape me in the great
" day which ſhall collect us all together ; and
" your form, diſengaged and purified from all
" earthly imperfection, ſhall appear to me, and
" ſtrike my raviſhed eye, in the midſt of my-
" riads."

CONTENTS

OF THE FIRST VOLUME.

Page

INTRODUCTION, or a Compendious View of the Parts of the Human Body - - - xv

PART I.

Of the Human Bones.

CHAP. I. *Of the Bones in General,* - 5

—— II. *Of the Skeleton,* - - 52

Sect. I. *Of the Head,* - - 53

§ 1. *The Cranium,* - - ib.

§ 2. *Bones of the Face,* - - 101

Sect. II. *Of the Trunk,* - - - 137

§ 1. *The Spine,* - - - ib.

§ 2. *The Pelvis,* - - 166

§ 3. *The Thorax,* - - 177

Sect. III. *Of the Superior Extremities,* - 192

§ 1. *The Shoulder,* - - - ib.

§ 2. *The Arm,* - - - 201

§ 3. *The Fore-arm,* - - 206

§ 4. *The Hand,* - - 212

Page

Sect. IV. *Of the Inferior Extremities,* - 226
§ 1. *The Thigh,* - - 227
§ 2. *The Leg,* - - - 232
§ 3. *The Foot,* - - 242
APPENDIX. *Of the Female Skeleton,* - 256

PART II.

Deſcription of the Human Muſcles.

CHAP. I. *Muſcles of the Integuments of the Cranium,* 264
—— II. *Of the Muſcles of the Ear,* - 265
—— III. *Of the Muſcles of the Eye-lids,* - 269
—— IV. *Muſcles of the Eye-ball,* - - 271
—— V. *Muſcles of the Noſe,* - - 273
—— VI. *Muſcles of the Mouth and Lips,* - 274
—— VII. *Muſcles of the Lower Jaw,* - 279
—— VIII. *The Muſcles which appear about the anterior Part of the Neck,* - 281
—— IX. *Muſcles ſituated between the Lower Jaw and Os Hyoides,* - 283
—— X. *Muſcles ſituated between the Os Hyoides and Trunk,* - - 285
—— XI. *Muſcles ſituated between the Lower Jaw and Os Hyoides laterally,* - - 287
—— XII. *Muſcles ſituated about the Entry to the Fauces,* - - 290
—— XIII. *Muſcles ſituated on the poſterior Part of the Pharynx,* - - 292
—— XIV. *Muſcles ſituated about the Glottis,* 294
—— XV. *Muſcles ſituated on the anterior Part of the Abdomen,* - - 296

Page

CHAP. XVI. *Muſcles about the Male Organs of Generation,* - - 301

—— XVII. *Muſcles of the Anus,* - 304

—— XVIII. *Muſcles of the Female Organs of Generation,* - - 305

—— XIX. *Muſcles ſituated within the Pelvis,* 307

—— XX. *Muſcles ſituated within the Cavity of the Abdomen,* - - 308

—— XXI. *Muſcles ſituated on the anterior Part of the Thorax,* - - 312

—— XXII. *Muſcles ſituated between the Ribs and within the Thorax,* - - 314

—— XXIII. *Muſcles ſituated on the anterior part of the Neck cloſe to the Vertebræ,* 316

—— XXIV. *Muſcles ſituated on the poſterior Part of the Trunk,* - 317

—— XXV. *Muſcles of the Superior Extremities,* 329

—— XXVI. *Muſcles ſituated on the Os Humeri,* 333

—— XXVII. *Muſcles ſituated on the Cubit or Fore-arm,* 335

—— XXVIII. *Muſcles ſituated on the Hand chiefly,* 342

—— XXIX. *Muſcles of the Inferior Extremities,* 349

—— XXX. *Muſcles ſituated on the Thigh,* 354

—— XXXI. *Muſcles ſituated on the Leg,* 360

—— XXXII. *Muſcles which are ſituated chiefly on the Foot,* - - 367

Of Muſcular Motion, - - 373

PART III.

Explanations of the Skeleton and Muſcles.

Page

Table I. *A front view of the Human Skeleton, with ſome of the Ligaments and Cartilages which connect the Bones of each other,* - 397

—— II. *The firſt Layer of Muſcles ſituated on the anterior Part of the whole Body, immediately under the common Teguments, and tendinous Faſciæ,* 401

—— III. *The ſecond Layer of Muſcles on the anterior Part of the whole body,* - 404

—— IV. *Repreſents the right eye ball, the ſix muſcles which move it, taken out of the orbit, with the Optic Nerve,* - 407

—— V. *A back view of the Human Skeleton, with ſome of the Ligaments, and Cartilages which connect the bones,* - 413

—— VI. *A back view of the Muſcles which are immediately ſituated below the common Integuments,* - - 414

—— VII. *The ſecond Layer of the Muſcles on the Back Part of the Body,* - 416

—— VIII. *The third Layer of Muſcles on the poſterior Part of the Body, with ſome of the Ligaments and naked bones,* - 420

—— IX. *A front view of the Bones of a Female Skeleton,* 425

—— X. *A lateral View of the Bones of a Female Skeleton,* - - 427

PART IV.

An Enumeration of the Cartilages, Ligaments, &c. of the Joints.

	Page
Those of the Head,	428
of the Vertebræ,	431
of the Sternum and Ribs,	436
of the Superior Extremities,	439
of the Pelvis and Inferior Extremities,	448

PART V.

A Description of the Common Integuments.

Cuticula	464
Corpus Mucosum,	466
Cutis Vera,	467
Glands of the Skin,	471
Uses of the Skin,	ib.
Membrana Adiposa and Fat,	472
The Nails,	476
The Hairs,	477
The supposed Integuments of the Antients,	479
Use of the Skin and the Nature of Touch,	480

INTRODUCTION.

A COMPENDIOUS VIEW *of the* PARTS *of the* HUMAN BODY.

Introduction.

THE human body is compoſed of firm and liquid parts, commonly called *ſolids* and *fluids*. Of the ſolid parts ſome are hard, others ſoft and flexible.

The ſolid parts are the chief ſubject of anatomy properly ſo called; by which term, borrowed from the Greek, we underſtand not only an artful decompoſition of the parts of the body, but alſo a methodical demonſtration and deſcription of the ſeveral parts when diſſected.

The hiſtory of the fluid parts comes into a ſyſtem of anatomy, only occaſionally; becauſe it properly belongs to what is called *phyſiology*, or the *animal economy*.

Anatomiſts ordinarily reduce all the ſolid parts under certain general claſſes, expreſſed by the common or generical names of *bone*, *cartilage*, *ligament*, *fibre*, *membrane*, *veſſel*, *artery*, *vein*, *nerve*, *muſcle*, *gland*, *fat*, *viſcus*, *organ*, &c. *.

Theſe terms are anatomical words, which expreſs ſeveral parts that have nearly the ſame ſtructure to outward appearance;

* The antients, who ſettled a general diviſion of the parts of the human body, from the mere outward appearance of the ſtructure, called ſome *ſimilar* or *ſimple*, and the reſt *organical* or *compound*.

pearance; and as they are often mentioned, we ſhall here give an explanation of them.

Explication of the general Terms of Anatomy.

By Bones, we mean in general the hardeſt, moſt ſolid, and moſt inflexible parts of the human body; the particular hiſtory of which is contained in the deſcription which ſhall be given of the Dry Bones.

A *cartilage* is a whitiſh or pearl coloured ſubſtance, ſofter than a bone, but harder than any other part, ſmooth, poliſhed, pliable, and elaſtic. The cartilages ſhall be explained with the Dry Bones and Joints.

A *ligament* is a white, fibrous, compact ſubſtance, more pliable than a cartilage, difficult to be broken or to be torn, and yielding but a very little when drawn out with force. The ligaments ſhall be explained at full length in the deſcription of the Joints.

The name of *fibre* is given to ſmall filaments, which appear to be the moſt ſimple parts of the body, and which, by their different diſpoſition and connections, compoſe all the other parts. The fibres differ in ſubſtance, being either membranous, fleſhy, tendinous, or bony; in direction, being either ſtraight, oblique, longitudinal, tranſverſe, circular, or ſpiral; and in ſize, being either large, ſmall, long, or ſhort.

By *membrane*, we underſtand a pliable texture of fibres interwoven or diſpoſed together in the ſame plane. They differ in thickneſs, according to the ſmallneſs of their fibres and number of their planes. Theſe particular planes are termed *laminæ*, and are diſtinguiſhed into external, internal, middle, &c.

The difference of membranes in general depends on that of the fibres, of which they are compoſed. Small portions

tions of membranes, especially when they are very thin, are called *pelliculæ*; and some membranous laminæ are united together by the intervention of a particular substance, composed of this sort of pellicles, and called *the cellular* or *spongy substance*.

Vessels are tubes, ducts, or canals, more or less flexible, composed of different membranes, the strata of which are generally termed *tunicæ* or *coats*. Some of them are divided into branches, and these again into rami and ramifications, which gradually diminish, but still remain hollow.

The general design of the vessels is to contain fluids; from the diversity of which they are distinguished into blood-vessels, vasa lactea, lymphatica, &c. The last and smallest extremities of all sorts of vessels are generally termed *capillaries*.

The *blood vessels* are of two kinds; one of which receives the blood from the heart, and distributes it to all the parts of the body, and are named *arteries*; the other brings the blood from all the parts back to the heart, and are called *veins*; and some of these have the name of *sinuses*.

The arteries are thicker than the veins, and may be distinguished by this mark in dissected dead bodies; and in living bodies they are known by a certain beating called the *pulse*. The trunks of the veins lie nearer the surface of the body than the arteries, and are furnished with valves, that is, with small membranous sacculi, fixed at different distances to the sides of their cavities. The openings of these valves are broad, and turned toward the heart; but their bottoms are turned the contrary way. In some places these valves are single, in others double, triple, &c.

By *nerves*, anatomists mean the white ropes which proceed from the cerebrum, cerebellum, and spinal marrow,

and are ſpread over all the parts of the body by filaments and ramifications.

Each nervous rope may be conſidered as a membranous veſſel, the cavity of which is filled by a great number of membranous longitudinal ſepta, and by medullary filaments which lie between the ſepta.

Muſcles are bundles of fibres, called by anatomiſts *fibræ motrices*, of a reddiſh colour, and of different lengths.

The middle portion of the moving fibres is the principal, and differs from the extremities in being red, thick, ſoft, and capable of contraction.

This middle portion of each moving fibre is ſaid to be fleſhy, and forms what is properly called *fleſh*. The extremities are called *tendinous*, and the ſubſtance formed by them *tendons*.

Glands are cluſters or moleculæ, diſtinguiſhable from all the other parts of the body by their form, conſiſtence, texture, and connection.

They are, in general, compoſed of arteries, veins, nerves, and other particular veſſels, and of a ſubſtance which unites all theſe together in their different folds, contortions, and intertextures, all inveſted by a membranous covering.

The office of glands is to ſeparate from the maſs of blood, by means of certain ſecretory veſſels, fluids, which they diſcharge, either immediately, or by other veſſels termed *excretory*; and theſe fluids are either accumulated in particular reſervoirs, collected in the common cavities, or forced out of the body.

Fat and *marrow* are equivocal terms. By the firſt, we generally underſtand an oily, ſoft, white, or yellowiſh ſubſtance, of different conſiſtencies, collected between the ſkin and the muſcles, in the interſtices of the muſcles about the viſcera, &c. and compoſed partly of a cellulous or ſpongy ſubſtance,

ſubſtance, purely membranous, and partly of an oily matter of different thickneſs. This oily matter is called *fat*, eſpecially when ſeparated from the cellulous ſubſtance, and likewiſe *corpus adipoſum* by anatomiſts.

Marrow differs from fat only in the fineneſs of the membranous texture, in the ſubtility of the oily matter, and its ſituation within the bones.

By *viſcera*, we commonly underſtand parts contained in a great cavity, without being connected to it through their whole extent or circumference. Such are the ſtomach, inteſtines, &c. in the abdomen; and the lungs in the thorax.

Organ or *inſtrument* is a term given to every part capable of any function, whether it be ſimple or complex; and in this ſenſe we talk of the organ of ſight, of reſpiration, &c.

General Diviſion of the Human Body.

The HUMAN BODY is commonly divided into the *head*, *trunk*, and *extremities*. The trunk is again ſubdivided into the *neck*, *thorax*, and *abdomen*; and the extremities into *ſuperior*, called the *arms*, and *inferior*, called the *legs*.

The ancients divided the body into three great cavities, which they termed *venters*, and into four extremities. They called the head the *upper venter*, the thorax the *middle venter*, and the abdomen the *lower venter*. The neck was by ſome joined to the head; by others, to the thorax.

The moſt natural and plaineſt diviſion of the body, is into the head, neck, thorax, abdomen, arms, and legs; each of which portions may afterwards be ſubdivided.

Each portion is to be examined not only with regard to its ſurface or external confirmation, but alſo with regard

to

to its internal ſtructure or compoſition, and to the viſcera or organs which it contains or ſupports.

This is what gave occaſion to the antients to divide the body into parts containing and parts contained; and to ſubdivide the containing parts into common and proper. The common containing parts have been named *integuments*, by which they meant chiefly the ſkin and fat.

External Parts of the Head.

The HEAD, viewed on the outſide, is divided into the *hairy ſcalp* and *face*.

The inner parts of the head are ſurrounded by the bones of the cranium.

The *hairy ſcalp* covers the upper parts of the os frontis, the oſſa parietalia, the os occipitis, and the upper and lower portions of the oſſa temporum.

The uppermoſt part of the hairy ſcalp is termed the *vertex* or *fontanella*; the back part, *occiput*; the lateral parts, the *temples*. The vertex is diſtinguiſhed from the occiput by contorted hair, and from the temples by the ears.

The *arteries* on each ſide of the hairy ſcalp are: Arteria carotis externa, in general; arteria temporalis; arteria occipitalis; arteria angularis, by communication; arteria frontalis, a branch of the internal carotid.

The *veins* on each ſide of the hairy ſcalp are: Vena jugularis externa, in general; vena jugularis externa poſterior; vena temporalis; vena occipitalis; vena jugularis externa anterior, by communication; vena jugularis interna, by communication.

The *nerves* on each ſide of the hairy ſcalp are: Nervi ſuboccipitales, commonly called *the tenth pair from the medulla oblongata*; par primum cervicale; par ſecundum cervicale,

vicale, by communication; ramus frontalis of the orbitary or ophthalmic nerve; nervus ſympatheticus minor, called the *portio dura* of the auditory nerve.

The *face* comprehends all that portion of the ſurface of the head which lies between the hairy ſcalp and the neck, viz. the forehead, eyebrows, palpebræ, eyes, noſe, mouth, chin, cheeks, and ears.

The external parts of the *eye* are: The anterior portion of the globe of the eye, the membrana conjunctiva, the cornea lucida, caruncula lachrymalis, angles of the palpebræ, and the cilia, or hairs of each palpebra. The internal parts are; the globe of the eye, the iris and pupil, the tunica ſclerotica or cornea opaca, the choroides, arachnoides, cryſtalline, vitreous humour, aqueous humour, the anterior and poſterior chambers, the muſcles, and the optic nerve.

The external parts of the *ear* are: The great concha, the convex ſide of this concha, or hinder part of the ear, the great border, the fold or helix, the concavity, the broad eminence or antihelix, the ſmall anterior eminence or tragus, the ſmall poſterior eminence or antitragus, the lobe or lower extremity of the ear, and the meatus.

The external parts of the *noſe* are: The upper extremity or root of the noſe, the arch or back, the ſides of that arch, the tip of the noſe, the alæ, the nares, and the ſeptum narium. The internal parts are the cavity and bottom of the nares, the convolutions, the maxillary, ſphenoidal, frontal, and ethmoidal ſinuſes.

The external parts of the *mouth* are: The lips, one upper, the other lower, the angles or commiſſures of the lips, the border or edge of each lip, the foſſula which runs from the ſeptum narium to the edge of the upper lip, and the

the tranſverſe fold which ſeparates the upper lip from the chin.

The internal parts of the *mouth* are: The palate, the ſeptum palati, the uvula, the amygdalæ, gums, frœna of the lips, the tongue, its apex, roots, ſides, and frœnum. The other internal parts of the mouth, eye, noſe, and ear, ſuch as the glands, membranes, muſcles, &c. muſt be referred to the particular deſcriptions of theſe parts.

The *cheeks* are the lateral parts of the face, reaching downward from the eyes and temples, between the noſe and ears. The upper prominent part of the cheek is commonly termed *mala*.

The *chin* is the anterior protuberance, by which the lower part of the face is terminated, from whence it runs all the way to the neck. This under part of the chin, is termed the *baſis*; and it is diſtinguiſhed from the throat by a tranſverſe fold, which reaches from ear to ear. In the middle of the chin, there is ſometimes a foſſula or depreſſion.

The exterior *arteries* which belong to each ſide of the *face* are: Arteria carotis externa, in general; arteria facialis; arteria temporalis; arteria carotis interna, by communication.

The exterior *veins* diſtributed to each ſide of the face are: Vena jugularis externa; vena jugularis interna, by communication; vena facialis; vena temporalis.

The exterior *nerves* ſpread on each ſide of the face are: Nervus olfactorius; nervus orbitarius ſive ophthalmicus; nervus maxillaris ſuperior; nervus maxillaris inferior; portio dura of the auditory nerve; the ſecond pair of the nervi cervicales.

The *arteries* of the *forehead* are: Arteria temporalis, arteria angularis, which are branches of the external carotid;

tid; arteria frontalis, which is a branch of the internal carotid.

The *veins* of the forehead are: Vena frontalis; vena temporalis; vena angularis; ſinus orbitarius.

The *nerves* of the forehead are: Nervus orbitarius ſive ophthalmicus; nervus maxillaris ſuperior; nervus maxillaris inferior; portio dura of the nervus auditorius.

The *arteries* which go to the *eye* are: Arteria occularis, which is a branch of the internal carotid: arteria facialis ſive angularis; arteria maxillaris interna, which are branches of the external carotid.

The *veins* which belong to the eye are: Vena temporalis, which is a branch of the poſterior external jugular; vena frontalis, and vena angularis, which are branches of the vena faſcialis; ſinus orbitarius; vena jugularis interna, by communication.

The *nerves* belonging to the eye are: Nervis opticus; nervis motor communis, or the third pair; nervus trochlearis, or the fourth pair; nervus orbitarius ſive ophthalmicus; and nervus maxillaris ſuperior, branches of the fifth pair; nervus motor externus, or the ſixth pair; portio dura of the auditory nerve; nervus ſympatheticus maximus, by communication.

The *arteries* diſtributed to the *noſe* are: Arteriæ maxillaris externæ; arteriæ maxillaris internæ; arteria occularis; arteria labiorum orbicularis, by communication.

The *veins* belonging to the noſe have their names correſponding with thoſe of the arteries.

The *nerves* which go to each ſide of the noſe are: Nervus olfactorus: nervus orbitarius ſive ophthalmicus, both immediately and by communication; nervus maxillaris ſuperior; portio dura of the auditory nerve.

The *arteries* which go to the *ear* are: Arteria temporalis,

lis, a branch of the external carotid; arteria auricularis, a branch of the temporalis; arteria occipitalis, by communication; arteria vertebralis, by means of the arteria basilaris, which is a continuation of it; arteria carotis interna, by communication with the arteria basilaris.

The *veins* belonging to the ear are: Vena temporalis; vena occipitalis; vena cervicalis; vena jugularis interna, by several communications; sinus petrosus durae matris.

The *nerves* distributed to the ear are: Nervus maxillaris inferior, the third branch of the fifth pair; nervus auditorius, the seventh pair; nervus suboccipitalis, the tenth pair by communication; the second cervical pair; nervus sympatheticus maximus.

The *arteries* which go to the *mouth*, *tongue*, *&c.* are: The artery of the chin; arteria coronaria sive orbicularis labiorum, both being branches of the external carotid; arteria maxillaris interna; arteria palatina, a branch of the maxillaris externa; arteria sublingualis.

The *veins* belonging to each side of the mouth, tongue, *&c.* are: Vena maxillaris externa; vena maxillaris interna; venæ raninæ. All these are branches of the external jugular. Vena jugularis interna, by several communications; vena gutturalis superior, a branch of the internal jugular.

The *nerves* distributed to the mouth, tongue, and salivary glands, &c. are: Nervus maxillaris superior; nervus maxillaris inferior, both branches of the fifth pair; portio dura of the auditory nerve; the eighth pair; the ninth pair; the second pair of cervical nerves; nervus sympatheticus maximus, by communication.

The *cheeks* on each side are furnished with *arteries* and *veins* from the nearest ramifications of the temporal and maxillary arteries and veins; and with *nerves* from the portio

portio dura of the auditory nerve, and from the ſuperior and inferior maxillary nerves.

Parts within the Cranium.

The CRANIUM comprehends all that portion of the head which reaches from the upper part of the orbit to the upper and back part of the neck; at the ſides it extends as low as the paſſage into the ears.

It is lined internally by the dura mater, and divided by a proceſs of that membrane into a large upper cavity and a ſmall under one. The upper cavity contains the cerebrum, the under one contains the cerebellum and medulla oblongata.

The arteries of the brain are: Arteriae carotides internae; arteriae vertebrales.—The arteries of the dura mater are, the arteriae durae matris, mediae maximae, &c. The veins of the brain run to the ſinuſes, and theſe end in the venae jugulares internae.

Parts of the Neck in general.

The NECK, in general, is divided into the anterior part or throat, and poſterior part or napé. The throat begins by an eminence, and terminates by a foſſula. The nape begins by a foſſula, which, as it deſcends, is gradually loſt. The neck contains the larynx, a part of the trachea arteria, the pharynx, a part of the œſophagus, the muſculi cutanei, ſterno-maſtoidaei, ſterno-hyoidaei, thyro-hyoidaei, omo-hyoidaei, ſplenius complexus, the muſculi vertebrales, which lie upon the firſt ſeven vertebrae, and a portion of the medulla ſpinalis.

The *arteries* which go to the neck are: Arteriae carotides externae, and internae; arteriae vertebrales; arteriae cervicales.

The *veins* belonging to the neck are: Venae jugulares in

general; venae jugulares externae, and internae; venae cervicales; venae vertebrales.

The *nerves* diſtributed to each ſide of the neck are: Portio dura of the auditory nerves; the eighth pair; nervus acceſſorius octavi paris; the ninth pair; nervus ſuboccipitalis, or the tenth pair; the ſeventh cervical pair; nervus ſympatheticus maximus.

Parts of the Thorax.

By the THORAX we commonly underſtand all that part of the body which anſwers to the extent of the ſternum, ribs, and vertebrae of the back, both outwardly and inwardly.

The thorax is divided into the anterior part, called commonly the *breaſt;* the poſterior part, called the *back;* and the lateral parts, called the *right* and *left ſides.*

The external parts of the thorax, beſides the ſkin and membrana adipoſa, are principally the mammae, and the muſcles which cover the ribs and fill the ſpaces between them. In the mammae we ſee the papillae or nipples, and a ſmall coloured circle which ſurrounds them. The muſcles are, the pectorales majores and minores, ſubclavii, ſerrati majores, ſerrati ſuperiores poſtici, latiſſimi dorſi, and vertebrales: and to theſe we may add the cuculares rhomboides and muſcles which cover the ſcapula.

The internal parts of the thorax are contained in the large cavity of that portion of the trunk which the antients called the *middle venter;* but the moderns name it ſimply *the cavity of the breaſt.* The cavity is lined by a membrane named *pleura*, and divided into two lateral cavities by a membranous ſeptum named *mediaſtinum*, which is a production or duplicature of the pleura.

Theſe parts are the heart, pericardium, trunk of the aorta, great arch of the aorta, beginning of the carotid and

and ſubclavian arteries, the ſuperior portion of the deſcending aorta, the intercoſtal arteries, the vena cava ſuperior, vena azygos, termination of the ſubclavian veins, a portion of the aſpera arteria, and of the œſophagus; the ductus lacteus or thoracicus, the lungs, pulmonary artery, pulmonary veins, &c.

The *arteries* and *veins* which particularly belong to the thorax are; Arteriae and venae thoracicae, ſuperiores and inferiores; arteriae and venae mammariae, internae and externae; arteriae and venae intercoſtales, ſuperiores and inferiores; arteriae and venae ſpinales, with the venal ſinuſes of the canal of the ſpine.

The *nerves* diſtributed to the thorax are: Nervi ſympathetici medii, or the eighth pair; nervi ſympathetici univerſales, commonly called *intercoſtales;* the twelve dorſal pairs; nervi diaphragmatici; nervi thoracici externi.

The cavity of the thorax is terminated downward by the diaphragm, which parts it from the abdomen.

Parts of the Abdomen.

The ABDOMEN begins immediately under the thorax, and terminates at the bottom of the pelvis of the oſſa innominata. Its circumference or outer ſurface is divided into regions; of which there are three anterior, *viz.* the epigaſtric or ſuperior region, the umbilical or middle region, and the hypogaſtric, or lower region. There is but one poſterior region, named *regio lumbaris.*

The *epigaſtric region* begins immediately under the appendix enſiformis at a ſmall ſuperficial depreſſion called *the pit of the ſtomach,* and in adult ſubjects ends above the navel at a tranſverſe line ſuppoſed to be drawn between the laſt falſe ribs on each ſide.

This region is ſubdivided into three parts, *epigaſtrium,* already named; and two lateral regions, termed *hypochondria.*

dria. The epigaſtrium takes in all that ſpace which lies between the falſe ribs of both ſides, and the hypochondria are the places covered by the falſe ribs.

The *umbilical region* begins in adults, above the navel, at the tranſverſe line already mentioned; and ends below the navel at another tranſverſe line, ſuppoſed to be drawn parallel to the former, between the two criſtae of the oſſa ilium.

This region is likewiſe divided into three parts; one middle, which is properly the regio umbilicalis; and two lateral, called *ilia* or *the flanks;* and they comprehend the ſpace between the falſe ribs and upper part of the os ilium on each ſide.

The *hypogaſtric region* is extended downward from the inferior limit of the umbilical region, and is divided into three parts; one middle, called *pubis*; and two lateral, called *inguina* or *the groins.*

The *lumbar region* is the poſterior part of the abdomen, and comprehends all that ſpace which reaches from the loweſt ribs on each ſide, and the laſt vertebra of the back, to the os ſacrum and neighbouring parts of the oſſa ilium. The lateral parts of this region are termed the *loins*, but the middle part has no proper name in men.

Laſtly, the bottom of the abdomen, which anſwers to the pelvis of the ſkeleton, is terminated anteriorly by the pudenda or parts of generation; and poſteriorly by the clunes or buttocks, and anus. The buttocks are ſeparated by a foſſa, which leads to the anus, and each buttock is terminated downward by a large fold which diſtinguiſhes it from the reſt of the thigh.

This lumbar region takes in likewiſe the muſculus quadratus lumborum on each ſide, the lower portions of the ſacro-lumbales, of the longiſſimi and latiſſimi dorſi, the multiſidus ſpinae, &c.

The

The ſpace between the anus and the parts of generation is called *perinæum;* and is divided into two equal lateral parts by a very diſtinct line, which is longer in males than in females, as we ſhall ſee in another place.

The cavity of the abdomen, formed by the parts already mentioned, and covered by the ſkin and membrana adipoſa, is lined on the inſide by a particular membrane, called *peritonæum.* It is ſeparated from the cavity of the thorax by the diaphragm, and terminated below by the muſculi levatores ani.

This cavity contains the ſtomach and the inteſtines; which are commonly divided into three ſmall portions, named *duodenum*, *jejunum*, and *ileum;* and three large, called *cæcum*, *colon*, and *rectum.* It contains likewiſe the meſentery, meſocolon, omentum, liver, gall-bladder, ſpleen, pancreas, glands of the meſentery, vaſa lactea, receptaculum chyli, kidneys, renal glands, ureters, bladder, and the internal parts of generation in both ſexes.

The principal *arteries* of the abdomen are; Arteriae epigaſtricae ſuperiores, which are the loweſt portions of the mammariae internae; aorta inferior; arteria caeliaca; arteria meſenterica ſuperior; arteriae renales, called formerly *emulgentes;* arteriae ſpermaticae; arteria meſenterica inferior; arteriae lumbares; arteriae iliacae; arteriae hypogaſtricae; arteriae epigaſtricae inferiores; arteriae haemorrhoidales; arteriae pudicae.

The principal *veins* of the abdomen are: The inferior portions of the venae mammariae internae; venae renales; venae lumbares; venae ſpermaticae; venae iliacae; venae hypogaſtricae; vena meſaraica minor, ſive haemorrhoidales internae; vena meſaraica major; vena ſplenica; vena portae ventralis; vena portae hepatica.

The principal *nerves* of the abdomen are: Nervi ſtomachici, formed by the extremity of the eighth pair; nervi ſympathetici

ſympathetici maximi, the inferior portion; the two ſemilunar or plexiform ganglions; plexus ſtomachicus; plexus hepaticus; plexus ſplenicus; plexus renalis; plexus meſentericus ſuperior; plexus meſentericus inferior; nervi lumbares; nervi ſacri; nervi crurales, their origin; nervi ſciatici, their origin.

Parts of the Upper Extremities.

The whole ARM is divided, as in the ſkeleton, into the ſhoulder, the arm properly ſo called, the fore-arm, and the hand. But to theſe we muſt here add the ſhoulder, the axilla or arm-pit, the elbow, the fold of the arm, and the hollow of the hand.

What is called the ſhoulder, is formed by the fleſhy belly of the muſculus deltoides; and the axilla, by the correſponding edges of the pectoralis major and latiſſimus dorſi. The elbow anſwers to the olecranum; the fold of the arm is on the fore-ſide of the articulation of the os humeri, with the bones of the fore-arm, and the hollow of the hand is in the middle of the palm.

The arm, properly ſo called, is principally covered, from the ſhoulder downward, by the biceps, brachialis, and the triceps. The fore-arm is furniſhed with thoſe muſcles which move the radius on the ulna, and the carpus on the fore-arm. The hand has few very conſiderable fleſhy parts, except the muſcles of the thumb and little finger, the lumbricales and interoſſei.

The *arteries* of the upper extremity are: Arteria axillaris; arteria humeralis; arteriae ſcapulares; arteria articularis; arteria brachialis; arteriae collaterales; arteria cubitalis; arteria radialis; arteria interoſſeae anterior; arteriae interoſſeae poſteriores. The ſuperficial and deep arches in the palm of the hand.

The *veins* of the upper extremity are: Areolae venoſae dorſi

dorſi manus; vena ſalvatella, ſive auricularis; vena cephalica cubiti, ſive radialis; venae cubiti ſatellites; vena baſilica cubiti, ſive ulnaris; vena mediana, or major; vena mediana cephalica; vena mediana baſilica; vena profunda cubiti; vena profunda ſuperior; venae ſatellites arteriae brachialis; vena brachii cephalica; vena brachii baſilica; venae muſculares; venae ſcapulares; vena axillaris.

The *nerves* of the upper extremity are; Nervi brachiales in general, formed by the laſt four cervical and firſt dorſal pairs; nervus ſcapularis; nervus articularis; nervus cutaneus; nervus muſculo-cutaneus; nervus muſcularis; nervus ulnaris; nervus radialis.

Parts of the Lower Extremities.

The LOWER EXTREMITIES of the whole body are divided, as thoſe of the ſkeleton, into the thigh, leg, and foot.

The *thigh* begins anteriorly on one ſide of the fold of the groin; and poſteriorly, a little above the lower half of the buttock. It terminates anteriorly at the patella on the knee, and poſteriorly at the poples or ham. It is formed chiefly by the muſcles which ſurround the os femoris, which are themſelves inveſted by the faſcia lata, *viz.* the glutaeus maximus, two vaſti, crureus, biceps, triceps, ſemi-membranoſus, ſemi-tendinoſus, gracilis internus, gracilis anterior or rectus, and ſartorius.

The *leg* has but very few muſcles on the fore-part, but a great many large ones behind: where the gaſtrocnemii and ſoleus, form a kind of belly, called the *calf of the leg*. The leg begins anteriorly at the knee below the patella, and poſteriorly at the poples; and it terminates below at the ancles.

Beſides the parts of the *foot* mentioned in the deſcription of the ſkeleton, that convex part near its articulation with the

the leg is termed the *neck of the foot:* and the under part, which is the basis of the whole lower extremity, the *sole of the foot.* The fleshy parts are not more considerable on the foot than on the hand.

The *arteries* of the lower extremity are: Arteria obturatrix, a branch of the hypogastrica; arteria glutaea, a branch of the hypogastrica; arteria sciatica, by communication; arteria pudica, by communication; arteria circumflexa externa; arteria circumflexa interna; arteria vasta; arteria cruralis; arteria poplitea; arteria tibialis anterior; arteria tibialis posterior; arteria peronaea; arteria plantaris.

The *veins* of the lower extremity are: Vena plantaris; vena parva saphena; vena magna saphena; venae tibiales; venae fibulares; vena poplitea; vena sciatica; vena glutea; venae femorales; vena obturatrix, &c.

The *nerves* of the lower extremity are: Nervus cruralis, formed by a complication of the five lumbares, especially of the first four; nervus sciaticus, formed by the union of the last two lumbares, and first three sacri; nervus sympathetic. cus maximus by communication with the nervi lumbares and sacri; nervus popliteus; nervus sciaticus internus, sive popliteus internus; nervus sciaticus externus, sive popliteus externus; nervus tibialis; nervus fibularis; nervi plantares.

A

SYSTEM OF ANATOMY,

WITH THE

PHYSIOLOGY.

PART I.

CONTAINING THE

ANATOMY

OF THE

HUMAN BONES.

BY THE LATE

ALEXANDER MONRO, M. D. AND F. R. S.

Fellow of the Royal College of Physicians, and Professor of Medicine and Anatomy in the University of Edinburgh.

TO THE

STUDENTS OF ANATOMY

IN THE

UNIVERSITY OF EDINBURGH.

GENTLEMEN,

WHEN this *Osteology* was first printed in 1726, I did not know that Albinus, Winslow, and Palfyn, were to publish descriptions of the bones; otherwise my papers probably would have remained yet undelivered to the printers. I flatter myself, however, that this *essay* has been of use to the gentlemen who did me the honour to attend my lectures, by assisting them to understand my sense and representation of things in this fundamental part of anatomy; and that it has possibly been of more advantage to them than a more complete work from an abler hand, unless my demonstrations had been in the order and method of such an author.

This view of your improvement, Gentlemen, is a prevailing argument with me to cause this essay to be reprinted; and you cannot reasonably blame me, if I likewise acknowledge another motive for it, which more particularly relates to myself. In a new edition an author has an opportunity of making his works more correct, complete, and consequently more acceptable to the public, who may perhaps be indulgent enough to think this little treatise not altogether useless; since more reasoning on the structure and morbid *phænomena* of bones is to be found in it, than in the other writers, who have confined themselves almost entirely to the descriptive or proper anatomical part of the *osteology*.

I have here kept in the plan of the former editions, by first considering, in the order that seemed to me most natural and methodical, every thing which I thought necessary to be known concerning bones in general; and, in the second place, I have described the several bones composing the skeleton.

The bones of adults are what I principally endeavour to describe; but I have added as much of the *osteogenea* as I think serviceable in the practice of physic and surgery.

That

That little might be omitted of what was formerly done on this ſubject, I have taken all the aſſiſtance I could from books; but have never aſſerted any anatomical fact on their authority, without conſulting nature, from which all the deſcriptions are made; and therefore the quotations from ſuch books ſerve only to do juſtice to the authors, who have remarked any thing in the ſtructure of the parts that was commonly omitted, and to initiate you in the hiſtory of anatomy, which I once propoſed to make complete, ſo far as related to this ſubject. But not being able to procure ſeveral books, and being ſenſible how many more may have never come to my knowledge, I laid aſide this deſign, of purpoſe omitted many I could have inſerted, and in ſome places I have changed an older author for a later one who has more fully or clearly deſcribed what I treated of. Beſide anatomiſts, I have alſo named ſeveral other authors to confirm my reaſoning by practical caſes; of which, it is not to be ſuppoſed, my own experience could furniſh a ſufficient variety.

You will readily obſerve, that I quote no paſſages with a view to criticiſe or condemn them. This precaution of giving no offence, is very neceſſary in thoſe who are ſufficiently conſcious of their being liable to lay themſelves open to juſt cenſure; and it prevents occaſions of uſeleſs wrangling, in which generally both parties are loſers, and the public has little advantage.

In this treatiſe I always make uſe of the moſt common name of each part, and have put the ſynonymous name to be met with in books at the foot of the page, that the reading might be ſmoother, and you might conſult them at your leiſure to aſſiſt you in underſtanding different authors.

The deſcriptions and reaſonings are blended, without which I always find young anatomiſts are ſoon diſguſted with authors. Their imaginations cannot follow a long chain of deſcriptions, eſpecially when they are not taught at the ſame time the uſes which the deſcribed parts ſerve. Their minds muſt have ſome relaxation, by a mixture of reaſoning, which never miſſes to ſtrike the fancy agreeably, and raiſes a ſtrong deſire to underſtand the principles on which it depends.

The phenomena of diſeaſes are all deduced in this eſſay from

the ſtructure of the parts, by way of corollaries and queſtions; which ſuch an anatomical work confined me to. And this method has otherwiſe a good effect; for, when a perſon meets with an uſeful propoſition, and is obliged to employ a little thought to find out its ſolution, the impreſſion it makes is deeper, and he acquires a fondneſs for it, as being in part his own diſcovery. My pupils have frequently aſſured me, that they could, with very ſmall reflection, trace out the whole reaſonings from which my concluſions were drawn. I hope that their ſucceſſors will alſo think this an agreeable manner of being inſtructed.

Thoſe gentlemen who deſired I would add the lectures which I pronounce in my college as a commentary upon the text, where the diſeaſes are mentioned, will, I perſuade myſelf, excuſe me for not complying with their deſire, when they conſider the deſign of this is to be a ſchool-book, and how great the difference is between inſtructing youth in private, and pretending to inform the public. *Art.* xxv. *vol.* v. of Medical Eſſays and Obſervations, publiſhed in this place, is one of theſe lectures which I gave as a commentary on the paragraph *(p.* 14.*)* concerning the different kinds of *caries*.

In this edition, I have corrected the miſtakes and obſcure paſſages which I diſcovered in the former, and in ſome places I have made the deſcriptions more full and exact, aiming all I could to ſhun unneceſſary minuteneſs on the one hand, and a blameable inaccuracy on the other. Whether I have hit that juſt medium, is what you and the public muſt now judge.

You have advantageous opportunities in this place of ſtudying all parts of medicine, under the profeſſors of its different branches in the Univerſity, and of ſeeing the practice of pharmacy, ſurgery, and phyſic, with our ſurgeon apothecaries, and in the Royal Infirmary, where the diſeaſed poor are carefully treated. Theſe your intereſt, and, I hope, your inclinations, will lead you, Gentlemen, ſo to improve, as that they become the happy means of your making a conſiderable figure in your ſeveral ſtations. Whatever aſſiſtance is in my power towards ſuch a deſirable event, ſhall be given with the greateſt pleaſure by,

Your humble ſervant,

ALEX. MONRO.

A

SYSTEM

OF

ANATOMY,

WITH THE

PHYSIOLOGY.

PART I.

OF THE

HUMAN BONES.

CHAP. I.

Of the BONES in general.

THE PERIOSTEUM.

BONES are covered by a membrane, named on that account PERIOSTEUM (*a*), which is ſo neceſſary to them, that we muſt examine its texture and uſes before we can underſtand their ſtructure.

The *perioſteum*, as well as moſt other membranes, can be divided into *layers* of fibres. The *exterior* layers, compoſed of the fibres of the muſcles connected to the bones, vary in their number, ſize, and direction, and conſequently occaſion

(*a*) Membrana circumoſſalis, omentum oſſibus impoſitum.

ſion a very great difference in the thickneſs and ſtrength of the perioſteum of different bones, and even of the different parts of the ſame bone. The *internal* layer is every where nearly of a ſimilar ſtructure, and has its fibres in the ſame direction with thoſe of the bone to which they are contiguous. Ought not then the name *perioſteum* to be applied, ſtrictly ſpeaking, only to this internal layer, to which the others are joined in an uncertain manner and number?

Some authors (*b*) endeavour to prove the internal layers of fibres of the perioſteum to be derived from the *dura mater:* For, ſay they, ſince the membrane covering the ſkull is plainly a production or continuation of the dura mater, which paſſes out between the ſutures; and ſince there are muſcles on the head, as well as in other parts, which might furniſh a perioſteum, it is needleſs to aſſign different origins to membranes which have the ſame texture and uſes. They add farther, in proof of this doctrine, that the perioſteum extends itſelf along the ligaments of the articulations from one bone to another; and therefore is continued from its origin over all the bones of the body.—While anatomiſts were fond of the hypotheſis of all membranes being derived from one or other of the two that cover the brain, a diſpute of this kind might be thought of conſequence: but now that the hypotheſis is neglected as uſeleſs, it is needleſs to examine the arguments for or againſt it.

Except where muſcles, cartilages, or ligaments, are inſerted into the perioſteum, its external ſurface is connected to the ſurrounding parts by thin cellular membranes, which can eaſily be ſtretched conſiderably, but ſhorten themſelves whenever the ſtretching force is removed. When theſe membranes are cut off or broken, they collapſe into ſuch a ſmall ſpace, that the ſurface of the perioſteum ſeems ſmooth and equal.

When

(*b*) Havers, Oſteolog. Nov. diſc. 1. p. 16.

When we attempt to tear off the periosteum from bones, we see a great number of white threads produced from the membrane into them; and, after a successful injection of the arteries with a red liquor, numerous vessels are not only seen on the periosteum (*c*), but most of the fibres sent from the membranes to the bone, shew themselves to be vessels entering it, with the injected liquor in them; and when they are broken, by tearing off the periosteum, the surface of the bone is almost covered with red points.

The veins corresponding to these arteries are sometimes to be seen in subjects that die with their vessels full of blood; though such numerous ramifications of them, as of the arteries, can seldom be demonstrated, because few of them naturally contain coloured liquors, and such lipuors can difficultly be injected into them. This, however, is sometimes done (*d*).

The great sensibility of the periosteum in the deep seated species of paronychia, in exostoses, nodi, tophi, and gummata, from a lues venerea, or whenever this membrane is in an inflamed state, is a sufficient proof that it is well provided with nerves, though they are perhaps too small to be traced upon it; and therefore it is difficult to determine, whether they are sent along with the arteries in the common way, or are derived from the tendinous fibres of the muscles expanded on the periosteum (*e*).

Vessels also pass through the periosteum to the marrow; of which more hereafter. And frequently muscles, ligaments, or cartilages, pierce through the periosteum, to be inserted into the bones.

The

(*c*) Ruysch. Epist. 5. tab. 5. fig 1. 2. Epist. 8. tab. 9. fig. 1. 9.

(*d*) Sue Traité d'Osteologie, traduit de l'Anglois de Mr Monro; note in p. 9.

(*e*) See the dispute about the sensibility of this and other membranes, in Zimmerman. Dissert. de irritabilit.—Act. Gotting. vol. 2.——Haller sur la nature sensible et irritabile.—Whytt's Physiolog. essay 2.—Remar. Dissert. de fungo articular. § 26. 34.

The chief uſes of the perioſteum are: 1. To allow the muſcles, when they contract or are ſtretched, to move and ſlide eaſily upon the bones; the ſmooth ſurface of this membrane preventing any ill effects of their friction upon each other. 2. To keep in due order and to ſupport the veſſels in their paſſage to the bones. 3. By being firmly braced on the bones, to aſſiſt in ſetting limits to their increaſe, and to check their overgrowth. 4. To ſtrengthen the conjunction of the bones with their epiphyſes, ligaments, and cartilages, which are eaſily ſeparated in young creatures, when this membrane is taken away. 5. To afford convenient origin and inſertion to ſeveral muſcles which are fixed to this membrane. And, laſtly, To warn us when any injury is offered to the parts it covers; which being inſenſible, might otherwiſe be deſtroyed without our knowledge, or endeavouring to procure a remedy.

When the cellular ſubſtance connecting the perioſteum to the ſurrounding parts is deſtroyed, theſe parts are fixed to that membrane, and loſe the ſliding motion they had upon it; as we ſee daily in iſſues, or any other tedious ſuppurations near a bone.——When the veſſels which go from the perioſteum to the bones are broken or eroded, a collection of liquor is made between them, which produces a ſordid ulcer or rotten bone. This often is the caſe after fractures of bones and inflammations of the perioſteum, or after ſmall pox, meaſles, ſpotted fevers, end eryſipelas.——Do not the diſorders of the perioſteum, coming rather along with or ſoon after the cutaneous than other diſeaſes, indicate ſome ſimilarity of ſtructure in the perioſteum and ſkin?

OF THE COMPOSITION OF BONES.

The bones are the moſt hard and ſolid parts of the body: and like all other parts where large veſſels do not enter,

ter, are generally of a white colour; only in a living creature they are bluiſh, which is owing to the blood in the ſmall veſſels under their ſurface. The leſs therefore and fewer the veſſels are, and the thicker and firmer the bony ſurface covering the veſſel is, the bones are whiter. Hence the bones of adults are whiter than thoſe of children; and, in both young and old, the white colour of different bones, or of the ſeveral parts of the ſame bone, is always in proportion to their veſſels and ſolidities; which circumſtances ought to be regarded by ſurgeons, when they are to judge of the condition of bones laid bare.

Bones are compoſed of a great many plates (*f*), each of which is made up of fibres or ſtrings united by ſmaller fibrils (*g*); which being irregularly diſpoſed, and interwoven with the other larger fibres, make a reticular work.—— This texture is plainly ſeen in the bones of fœtuſes, which have not their parts cloſely compacted; and in the bones of adults which have been burnt, long expoſed to the weather, or whoſe compoſition has been made looſe by diſeaſes.—— The chinks which are generally made according to the directions of the larger fibres of bones that have undergone the action of fire or of the weather, ſhew the greater ſtrength of theſe than of the fibres which connect them.——Numerous accurate obſervations of the different times in which exfoliations are made from the ſides or ends of ſimilar bones, might bid fair to determine what is the proportional force of coheſion in the two ſorts of fibres.

The plates are ſaid (*h*) to be firmly joined to each other by a great number of *claviculi*, or ſmall bony proceſſes, which, riſing from the inner plates, pierce through ſome, and are fixed into the more external ones. Of theſe nails, four

(*f*) Squamae, bracteae, laminae.

(*g*) Malpigh. Anat. plant. et oper. poſthum.

(*h*) Gagliard. Anatom. oſſium nov. invent. illuſtrat. cap. 1. obſ. 2.

four kinds, viz. the *perpendicular*, *oblique*, *headed*, and *crooked*, have, been described: But in bones fitly prepared, I could only see numerous irregular processes rising out from the plates (*i*).

Though the exterior part of bones is composed of firm compact plates, yet they are all more or less cavernous internally. In some (*e. g.* middle thin part of the *scapula* and *os ilium*) the solid sides are brought so near, that little cavity can be seen; and in others (middle of *os humeri*, *femoris*, &c.) the cavities are so large, that such bones are generally esteemed to be hollow or fistular. But the internal spongy texture is evident in young animals; and some of it may be seen to remain in those of the greatest age, when bones are cautiously opened, after they have been kept so long as to be free of the oil they contain, or after being burnt.

This spongy cavernous internal part of bones is generally called their CANCELLI, or LATTICE-WORK, and is formed in the following manner. The plates are firmly joined about the middle of the bone; but as they are extended towards its ends, the more internal parts separate from the exterior, and stretch out their fibres towards the axis of the bone, where they are interwoven with the fibres of other plates that have been sent off in the same way. Seeing the plates are thus constantly going off, the solid sides of the bones must become thinner, and the lattice-work must be thicker and stronger towards their ends. This is evident in many of them, where the solid sides of their middle are very thick, and the cancelli are scarce observable; whereas, at the ends, where their diameter is greatest, the solid walls or sides are not thicker than paper, and the cancelli are numerous, and large enough to fill up the whole space left between the sides.

The

(*i*) Malpigh. Oper. posthum.

The twiſting and winding which theſe cancelli make, and the interſtices which they leave, differ conſiderably in figure, number, and ſize; and therefore form little cells, which are as different, but communicate with each other. Some writers (*k*) minutely remark theſe different appearances of the cancelli, after they begin to ſeparate from the plates; and from thence diſtinguiſh them into *wrinkled*, *perforated*, and *net-like*.

The cancelli ſuſtain the membranous bags of the marrow which are ſtretched upon them, and thereby hinder theſe membranous parts from being torn, or removed out of their proper places, in the violent motions and different poſtures which the bones are employed in. This ſupport which the cancelli afford to the marrow, alſo ſaves its membranes and veſſels, in the lower parts of the bones, from being compreſſed by the weight of the marrow above.

The depreſſions between the fibres of the external plates of bones appear like ſo many furrows on their ſurface, into each of which the perioſteum enters; by which the ſurface of contact, conſequently the coheſion, between it and the bone, is conſiderably increaſed, and a greater number of veſſels is ſent from it into the bone than if it was a plain ſurface.

Both on the ridges and furrows, numerous little pits or orifices of canals are to be ſeen, by which the veſſels paſs to and from the bones.

After a ſucceſsful injection, the arteries can be traced in their courſe from the pits to the plates and fibres; and, in ſawing, cutting, or raſping the bones of living creatures, theſe veſſels diſcover themſelves by the ſmall drops of blood which then ooze out from the moſt ſolid part of the bones. But the cleareſt demonſtration of the intimate diſtribution of theſe ſmall arteries, is, to obſerve the effect of ſuch a tinging

(*k*) Gagliard. Anat. oſſium, cap. 1. obſ. 4. 5. 6. 7.

tinging ſubſtance as can retain its colour, when ſwallowed, digeſted, and mixed with the blood of any living animal, and at the ſame time has particles ſmall enough to be conveyed into the veſſels of the bones; ſuch is rubia tinctorum, madder-root (*l*): For we ſee the gradual advances which this tincture makes from the perioſteum into the more internal parts of the bones, and how univerſally the diſtribution of the liquors is made, the whole bony ſubſtance being tinged by it. Whether the time in which this tinged liquor paſſes from the outer to the internal plates, till all the plates are made of its colour, and the time which the diſappearing of the dye, after giving the creature no more of this ſort of food, makes us think it takes to return, are the ſame in which the natural liquors circulate, is uncertain; becauſe this tinging ſubſtance may move more ſlowly, or may paſs more quickly than the natural liquors do.——The arteries are larger near each end than at the middle of the large bones that are much moved; becauſe they not only ſerve the bony plates near the ends, but paſs through them to the marrow.——As animals advance in age, the arteries of the bones become leſs capacious; as is evident, 1. From the bones of adults having leſs blood in them than thoſe of children have. 2. From many of them becoming incapable in old age of admitting the coloured powders uſed in injections, which eaſily paſs in youth. And, 3. From the bones of old creatures being more difficultly tinged with madder than thoſe of young ones. If authors have not miſtaken, the arteries of bones have ſometimes become very large (*m*).

We may conclude, from arteries being accompanied with veins, ſo far as we can trace them in every other part of the

(*l*) Philoſoph. Tranſact. num. 442. art. 8. num. 443. art. 2. num. 457. art. 4.——Mem. de l'acad. des ſciences, 1739, 1742.

(*m*) Diemerbroek Anat. lib. 9. cap. 1.——Mery Hiſt. de l'acad. des ſciences, 1704.

the body, that there are alſo veins in the bones; and the diſappearing of the tincture of madder, after the bones of living animals are coloured with it, could not be without ſuch veins to carry it away; nay, the veins of bones can ſometimes be injected, and then ſeen (*n*).

The bones of a living animal are ſo inſenſible, that they can be cut, raſped, or burnt, without putting the creature to pain, and the nerves diſtributed in their ſubſtance cannot be ſhewn by diſſection; from which it might be inferred that they have no nerves diſtributed to them: but the general tenor of nature, which beſtows nerves to all other parts, ſhould prevent our drawing ſuch a concluſion. And if ſenſibility is a ſure proof of nerves entering into the compoſition of any part, as it is generally allowed to be, we have ſufficient evidence of nerves here in the bones: for the granulated red fleſh which ſprouts out from them, after an amputation of a limb, or performing the operation of the trepan, or after an exfoliation, is exquiſitely ſenſible; and in ſome ulcers of bones, where the perioſteum was all ſeparated, the patient ſuffered racking pain, if the bone was touched with a rough inſtrument; nor was he free from pain after the bone was perforated (*o*).——The reaſon why the nerves of rigid hard bones become inſenſible, is, that all nerves muſt have a conſiderable degree of flexibility at the part where an object is applied, otherwiſe they cannot be affected by its impreſſions. We ſee this illuſtrated in a very common analogous caſe, the growth of a new nail; when the former one has ſuppurated off, the thin membrane which firſt appears, is exquiſitely ſenſible; but gradually becomes dull in its ſenſation, till it can be cut or ſcraped, without cauſing pain, after it is formed into a hard nail.

From what has been ſaid of the veſſels of bones, it is evident,

(*n*) Sue trad. d'oſteolog. p. 9.

(*o*) Nicol. Maſſa, lib. introd. anat. cap. 30.

dent, that there is a conſtant circulation of fluids in every part of them; and that there is a perpetual waſte and renewal of the particles which compoſe the ſolid fibres of bones, as well as of other parts of the body; the addition from the fluids exceeding the waſte during the growth of the bones; the renewal and waſte keeping nearly equal in adult middle age; and the waſte exceeding the ſupply from the liquors in old age; as is demonſtrable from their weight: for each bone increaſes in weight as a perſon approaches to maturity; continues of nearly the ſame weight till old age begins, and then becomes lighter. The ſpecific gravity of the ſolid ſides, on the contrary, increaſes by age; for then they become more hard, compact, and denſe. In conſequence of this the bones of old people, are thinner and firmer in their ſides, and have larger cavities, than thoſe of young perſons.

The vaſcular texture of bones muſt make them ſubject to obſtructions, ecchymoſes, ulcers, gangrenes, and moſt other diſeaſes with which the ſofter parts are affected; and therefore there may be a greater variety of caries than is commonly deſcribed (*p*).

Hence we can account for the following appearances.

Haemorrhagies from fungous fleſh riſing out from the moſt ſolid part of a cut bone (*q*).

The regular alternate elevation and ſubſiding, or apparent pulſation, frequently to be ſeen in ſome of the cells of a carious bone.

Cells reſembling *cancelli*, ſometimes ſeen in the part of a bone, which, in a natural ſtate, is the moſt ſolid and firm (*r*).

A bone, as a tube, including another bone within it (*s*).

On

(*p*) Edin. Medical eſſays and obſ. vol. 5. art. 25,

(*q*) Medical eſſays, vol. 4. art. 21.

(*r*) Ruyſch. Theſ. 8. num. 8. Theſ. 10. num. 176.

(*s*) Idem, ibid.

On the internal ſurface of the ſolid parts of the bones there are orifices of canals, which paſs outwards through the plates to open into other canals that are in a longitudinal direction; from which other *tranſverſe* paſſages go out to terminate in other *longitudinal* canals; and this ſtructure is continued through the whole ſubſtance of bones; both theſe kinds of canals becoming gradually ſmaller as they approach the outer ſurface (*t*).——Theſe canals are beſt ſeen in a bone burnt till it is white. When it is broken tranſverſely, the orifices of the *longitudinal* canals are in view; and when we ſeparate the plates, the *tranſverſe* ones are to be obſerved. Here, however, we are in danger of believing both theſe ſorts of canals more numerous than they really are; becauſe the holes made by the proceſſes connecting the plates of bones have the appearance of the *tranſverſe* (*u*), and the paſſages for the blood-veſſels reſemble the *longitudinal* canals. I do not know how we are to keep free of error about the *tranſverſe* canals; but think we may diſtinguiſh between the two kinds of *longitudinal* ones: for the paſſages of the veſſels are largeſt near the external ſurface of the bone, and every tranſverſe ſection of them is circular; whereas the longitudinal canals are largeſt near the cancelli, and their tranſverſe ſections appear to me of a flat oval figure, which may be owing to the different momentum of the fluids conveyed in them.——The ſituation of the larger longitudinal canals, and of the paſſages of the larger veſſels, makes a bone appear more denſe and compact in the middle of its ſolid ſides, than towards its outer and inner ſurfaces, where it is ſpongy.

We ſee marrow contained in the larger tranſverſe and longitudinal canals juſt now deſcribed, and from thence judge that it paſſes alſo into the ſmaller ones. The drops of oil which we diſcover with a microſcope every where on the

(*t*) Havers Oſteolog. Nov. p. 43. (*u*) Morgagn. Adverſ. 3. animad. 25.

the ſurface of a recent bone fractured tranſverſely, and the oozing of oil through the moſt ſolid bones of a ſkeleton, which renders them greaſy and yellow, are a confirmation of the uſe of theſe canals. Of what advantage this diſtribution of the marrow through the ſubſtance of bones is, will be mentioned when the nature and uſe of this animal oil is inquired into.

Moſt bones have one or more large oblique canals formed through their ſides for the paſſage of the medullary veſſels, which are to be deſcribed afterwards.

Bones expoſed to a ſtrong fire in chemical veſſels, are reſolved, in the ſame manner as the other parts of animals, into *phlegm*, *ſpirit*, *volatile ſalt*, *fetid oil*, and a black *caput mortuum*. But the proportion of theſe principles varies according to the age, ſolidities, and other circumſtances of bones. Young bones yield the largeſt proportion of phlegm; ſpongy bones afford much oil; and ſolid ones give moſt ſalt and black reſiduum.——Though this reſiduum can ſcarce be changed by the force of fire while it is in cloſe veſſels; yet, when it is burnt in an open fire, the tenacious oil, to which it owes its black colour, is forced away, and a white earth is left that has little or no fixed ſalt in it (*v*). This earth ſeems to be the proper conſtituent ſolid part of bones, and the other principles give it firmneſs and tenacity: for the quantity of the earth is ſo great, that, after all the other principles are ſeparated from a bone, its former ſhape and ſize remain (*w*); but it is very brittle till it is moiſtened with water or oil, when it recovers ſome tenacity.—The increaſe of the proportion of earth in old peoples bones, is one reaſon of their being more brittle than thoſe of young people.

Leſt

(*v*) Later chemiſts have diſcovered this earth to be calcareous earth ſaturated with phoſphoric acid.

(*w*) Havers Oſteolog. Nov. diſc. I. p. 32.

Leſt any imagine the ſalts and oils of bones, while in a natural ſtate, to be of the ſame acrid kind with thoſe obtained from them by the chemical analyſis, it is to be obſerved, that theſe principles may be extracted from bones in the form of a very mild jelly, by boiling them in water.

The bones ſuſtain and defend the other parts of the body.

Bones are lined within, as well as covered externally, with a membrane; which is therefore commonly called PERIOSTEUM INTERNUM.

INTERNAL PERIOSTEUM.

THE *internal perioſteum* is an extremely fine membrane; nay, frequently, it has a looſe reticular texture; and therefore it is compared by ſome to the arachnoid coat of the ſpinal marrow: ſo that we cannot expect to divide it into layers as we can divide the external perioſteum. We can, however, obſerve its proceſſes entering into the tranſverſe pores of the bones, where probably they are continued to form the intermediate canals for the marrow diſtributed through the ſubſtance of the bones; and along with them veſſels are ſent, as from the external perioſteum, into the bone *(y)*. Theſe proceſſes being of a very delicate texture, the adheſion of this membrane to the bone is ſo ſmall, that it ſeparates commonly more eaſily from the bone than from the marrow which it contains: wherefore one might call it the common membrane of the marrow, rather than by the name it now has. But whether the one or the other deſignation ought to be given it, is not worthy a diſpute.

From the internal ſurface of the *internal perioſteum*, a great number of thin membranes are produced; which, paſſing acroſs the cavity, unite with others of the ſame kind, and form ſo many diſtinct bags, which communicate

(y) Winſlow Expoſition anat. des os frais, ſect. 82. 83.

with each other; and these again are subdivided into communicating vesicular cells, in which the marrow is contained. Hence it is, that the marrow, when hardened, and viewed with a microscope, appears like a cluster of small pearls; and that the hardened marrow of bones, buried long under ground, or laid some time in water, and then dried, is granulous (z).——This texture is much the same with what obtains in other cellular parts of the body, where fat is collected; only that the cells containing the marrow are smaller than those of the *tunica adiposa* or *cellulosa* elsewhere; which probably is owing to their being inclosed in the bones, where they are not so much stretched or extended as in other parts.

OF THE MARROW.

THE *Marrow* is the oily part of the blood, separated by small arteries, and deposited in these cells. Its colour and consistence may therefore vary according to the state of the vessels, and their distribution on the membranes of the cells.

The marrow, as well as the other fat of the body, chemically analized, yields, besides oil and water, a considerable proportion of an acid liquor, but no alkali (a). This may be the reason of its being less putrescent than the blood or most other parts of animals (b); which is a necessary quality in a substance that is constantly exposed to a considerable degree of heat, and is more in a stagnating condition than the other liquors.

Besides the arteries, which I mentioned already, (p. 12.) to be sent from the bones to the marrow, there is at least one artery for each bone; several bones have more, whose principal

(z) Ruysch. Thesaur. 9. num. 2. et Adversf. dec. III. obs. 9.

(a) Grutzmaker Dissert. de ossium medulla.—Haller element. physiolog. lib. 4. sect. 4.

(b) Pringle Appendix to camp diseases, exper. 47.

principal uſe is to convey and ſecern this oily matter. After theſe arteries have pierced the ſolid ſide of a bone, they are divided into ſeveral branches; which are ſoon diſtributed every where on the internal perioſteum, and afterwards ſpread their branches inwards on the medullary cells, and outwards through the tables of the bone.

The blood which remains after the ſecretion of the marrow is returned by proper veins, which are collected from the membranes into one or more large trunks, to paſs out at the ſame holes or paſſages at which the artery or arteries enter.

The general rule of the ſmall veſſels decreaſing in their capacities as animals advance in age, to which many phenomena in the animal oeconomy are owing, obtains here: for though the trunks of the medullary veſſels enlarge as animals turn older, yet the ſmall branches become ſmaller; as is evident from injections, which cannot be made to paſs near ſo far in theſe veſſels of adults as of children. Hence the marrow is bloody in children, oily and balmy in middle age, and thin and watery in old people.

By experiments made on the marrow when bones of living animals are opened or cut through (*c*), and from the racking pain with which ſuppurations within bones are frequently attended, we have ſufficient proof that the membranes here are ſenſible, and conſequently have nerves diſtributed to them. Hippocrates (*d*) might therefore ſay juſtly, that a wound penetrating into the cavity of a bone may produce a delirium.

The veſſels of the marrow, wrapt up in one common coat from the perioſteum, paſs through the bones by proper canals; the moſt conſiderable of which are about the middle of each bone, and are very oblique. Sometimes theſe veſſels

(*c*) Du Verney, Memoires de l'acad. des ſciences, 1700.
(*d*) Aphoriſm. § 7. aph. 24.

veſſels continue at a little diſtance in their paſſage, when the canal is divided by a ſmall bony partition or two.

From the ſtructure of the contents of the bones, we may judge how theſe parts, as well as others, may be ſubject to œdema, phlegmon, eryſipelas, ſcirrhus, &c. and may thence be led to a cure of each, before the common conſequence, putrefaction, takes place, which frequently occaſions the loſs of the limb, if not of the patient.

The marrow is of very conſiderable uſe to the bones: for by entering their tranſverſe canals, and paſſing from them into the longitudinal ones, it is communicated to all the plates to ſoften and connect their fibres, whereby they are preſerved from becoming too brittle; as we ſee they do in burnt bones, or thoſe long expoſed to the air, in people labouring under old age, pox, or ſcurvy. In all which caſes, the oil is either in too little quantity, or has its natural good qualities changed for worſe ones.

Beſides this advantage which the ſubſtance of bones has from the marrow, their articulations are ſaid (*e*) to receive no leſs benefit from it: for it is thought that the marrow paſſes into the articular cavities through the holes which are in the bones near the large joints. And as a proof of this it is alleged, that butchers, upon ſeeing the greater or leſſer quantity of marrow in the bones of cows, can tell whether they have travelled far or little before they were ſlaughtered.

When the marrow, after having ſerved the uſes above mentioned, is reaſſumed into the maſs of blood (as it is continually in common with all other ſecreted liquors that have not paſſages formed for conveying them out of the body), it corrects the too great acrimony communicated to the ſaline particles of our fluids by their circulation and heat, in the ſame manner as lixivial ſalts are blunted by oil in making

(*e*) Joan de Muralto Vade-mecum Anat. exercit. 5. § 3. Havers Oſteolog. Nov. diſ. 3. p. 170.

making ſoap. Hence, in acute diſeaſes, the marrow, as well as the other fat of the body, is quickly waſted, but muſt be immediately ſupplied by liquors from the veſſels; ſeeing the cells within the bones, which have no aſſiſtance to their contraction from the preſſure of the atmoſphere, cannot collapſe, as the tela cellularis under the ſkin does when the liqour in its cells is abſorbed; the bones therefore are always full.

Since it is the nature of all oil to become thin and rancid when expoſed long to heat, and bones have much oil in their firm hard ſubſtance, we may know why an ungrateful ſmell and dark-coloured thin ichor proceed more from corrupted bones than from the other parts of the body; and we can underſtand the reaſon of the changes of colour which bones undergo, according to their different degrees of mortification.—Hence likewiſe we may learn the cauſe of a ſpina ventoſa, and of the difficulty of curing all caries of bones proceeding from an obſtruction and conſequent putrefaction of the marrow; and of the quick pulſe, thirſt, and hectic paroxyſms, ſo often attending theſe diſeaſes. Theſe phenomena alſo teach us the reaſon of the fatal prognoſis taken from black fetid urine in fevers.

Though bones ſo far agree in their ſtructure and annexed parts, yet we may obſerve a conſiderable difference among them in their magnitude, figure, ſituation, ſubſtance, connection, uſes, &c. From which authors have taken occaſion to diſtinguiſh them into as many claſſes as they could enumerate of theſe different circumſtances. But theſe being obvious to every perſon that looks on bones, I ſhall only mention one of them; which comprehends very near the whole bones of the body, and at the ſame time leads us to examine the moſt conſiderable variety that is to be found in the diſpoſition of their conſtituent parts, and in their uſes. It is this, that ſome bones are *broad* and *flat*, while others are *long* and *round*.

THE

THE DIFFERENT CLASSES, &c. OF BONES.

THE *broad* bones have thin ſides, by the plates being ſoon and equally ſent off to form the lattice-work; which therefore is thicker, and nearly of an equal form all through. By this ſtructure they are well adapted to their uſes, of affording a large enough ſurface for the muſcles to riſe from and move upon, and of defending ſufficiently the parts which they incloſe.

The *round* bones have thick ſtrong walls in the middle, and become very thin towards their ends; which is owing to very few plates ſeparating at their middle; where, on that account, the cancelli are ſo fine and ſmall, that they are not taken notice of: but ſuch bones are ſaid to have a large reſervoir of oil in this place. Towards their ends the lattice-work becomes very thick, and rather more complete than in the other ſort of bones.—Theſe round bones having ſtrong forces naturally applied to them, and being otherwiſe expoſed to violent injuries, have need of a cylindrical figure to reſiſt external preſſure, and of a conſiderable quantity of oil to preſerve them from becoming too brittle. Beſides which, they are advantageouſly provided with thick ſides towards their middle, where the greateſt forces are applied to injure them; while their hollowneſs increaſes their diameter, and conſequently their ſtrength to reſiſt forces applied to break them tranſverſely (*f*). Thus, for inſtance, in eſtimating the proportional reſiſtance of two cylindrical bones of unequal diameters, but conſiſting of an equal number of ſimilar fibres uniformly diſpoſed round each, it is plain,

1. That the abſolute force of theſe two bones is equal, becauſe they conſiſt of equal numbers of ſimilar fibres.

2. That the abſolute forces of all the fibres in each bone have

(*f*) Galilei Mechanic. dialog. 1.

have the ſame effect in reſiſting any power applied to break them, as if the ſum of all their forces was united in the reſpective centres of the tranſverſe ſections where the fractures are to be made. For, by hypotheſis, the fibres being uniformly diſpoſed in each, there is not any fibre in either bone that has not a correſponding fibre; the ſum of both whoſe diſtances from the axis of revolution (about which all the parts of the bone muſt revolve in breaking) is equal to two ſemi-diameters of the bone: conſequently each fibre, and all the fibres, may be regarded as reſiſting at the diſtance of one ſemi-diameter or radius from this axis, that is, in the centre.

3. Since the united force of all the fibres is to be regarded as reſiſting at a diſtance from the centre of motion equal to the ſemi-diameter, it follows that the total reſiſtance of all theſe fibres, or the ſtrength of the bone is proportional to its ſemi-diameter, and conſequently to its diameter.

I have here taken for an example one of the moſt ſimple caſes for calculating the proportional forces of bones. But, were it not too foreign to the preſent deſign, it might be univerſally demonſtrated, that of whatever figure bones are, and in whatever manner their fibres are diſpoſed, their ſtrength muſt always be in a ratio, compounded of the area of their tranſverſe ſections, or of their quantity of bony mater, and of the diſtance of the centre of gravity of theſe ſections from the centre of motion or fulcrum, on which the bone is ſuppoſed to be broken (*g*).

Since, therefore, the ſtrength of bones depends on their number of fibres, or quantity of matter, and the largeneſs of their diameters, one may conclude, that the part of a bone formerly fractured, and reunited by a *callus*, muſt be ſtronger than it was before the fracture happened; becauſe both

(*g*) See the demonſtration of this theorem by Dr Porterfield in the Edinburgh medical eſſays, vol. 1. art. 10.

both theſe advantages are obtained by a callus: which is a wiſe proviſion, ſince bones are never ſet in ſuch a good direction as they were naturally of; and then wherever a callus is formed, there is ſuch an obſtruction of the veſſels, that if the bone was again broken in the ſame place, the oſſific matter could not ſo eaſily be conveyed to reunite it. This callus may indeed, for want of compreſſion, be allowed to form into a ſpongy cellular ſubſtance (*b*); but even in this caſe the ſtrength of the bone is here increaſed by one or both of the cauſes above-mentioned.

Many bones have protuberances or *proceſſes* (*i*) riſing out from them. If a proceſs ſtands out in a roundiſh ball, it is called *caput* or *head*.—If the head is flatted, it obtains the appellation of *condyle*.—A rough unequal protuberance is called *tuberoſity*.—When a proceſs riſes narrow, and then becomes large, the narrow or ſmall part is named *cervix* or *neck*.—Long ridges of bones are called *ſpines*.—Such proceſſes as terminate in a ſharp point have the general name of *coronæ* (*k*) or *coronoid* beſtowed on them; though moſt of them receive particular names from the reſemblance they have, or are imagined to have, to other ſubſtances, *e. g. maſtoid*, *ſtyloid*, *anchoroid*, *coracoid*, *ſpinal*, &c.—Such proceſſes as form brims of cavities, are called *ſupercilia* (*l*).

Proceſſes ſerve for the advantageous origin and inſertion of muſcles, and render the articulations firm and ſtable.

Before leaving this ſubject, we muſt remark, that much the greater number of what are called proceſſes in adult bones, diſcover themſelves in children to be *epiphyſes*, or diſtinct bones, which are afterwards united to the other parts:

(*b*) Ruyſch. Theſaur. 8. n. 49. Muſ. anat. thec. B. repoſit. 2. n. 2.

(*i*) Αποφυσεις, εκφυσεις, εξοχη, προβολη, προβλημα, Exceſſus, explanatio, tuberculum, gibbus, eminentia, productio, extuberantia, protectura, enaſcentia.

(*k*) Roſtra, glandes.

(*l*) Ιτυες, οφρυες, αμφωνις, χειλη, Labra.

parts: ſuch are the *ſtyloid* proceſſes of the temporal bones, proceſſes of the *vertebræ*, *trochanteri* of the thigh, &c. However, as I deſign to inſiſt chiefly on the deſcription of the adult ſkeleton, in which the union of theſe parts is ſo intimate, that ſcarce any veſtige remains of their former ſeparation, I ſhall retain the common appellation of *apophyſe*, or *proceſs*, to all ſuch protuberances; but ſhall remark the principal ones that have no juſt title to this name, when they occur in the deſcription of particular bones.

On the ſurface of a great many of the bones there are cavities or depreſſions. If theſe are deep, with large brims, authors name them *cotylæ* (*m*).—If they are ſuperficial, they obtain the deſignation of *glenæ* or *glenoid*. Theſe general claſſes are again divided into ſeveral ſpecies:—Of which, *pits* are ſmall roundiſh channels ſunk perpendicularly into the bone;—*furrows*, long narrow canals formed in the ſurface;—*nitches*, or *notches*, ſmall breaches in the bone;—*ſinuoſities*, broad, but ſuperficial depreſſions without brims;—*foſſæ*, large deep cavities, which are not equally ſurrounded by high brims;—*ſinuſes*, large cavities within the ſubſtance of the bones, with ſmall apertures;—*foramina*, or holes, canals that pierce quite through the ſubſtance of the bones.—When this laſt ſort of cavity is extended any long way within a bone, the middle part retains the name of *canal*, and its ends are called *holes*.

The cavities allow the heads of bones to play in them; they lodge and defend other parts; they afford ſafe paſſage to veſſels, muſcles, &c. To mention more would engage us too much in the hiſtory of particular bones, which more properly belongs to the demonſtration of the *ſkeleton*, where we ſhall have occaſion to obſerve theſe ſeveral ſpecies of cavities.

To far the greater number of bones, whoſe ends are not

(*m*) Κοτυλδις, ὀξυβαφοι, Acetabula, pyxides, buccellæ.

joined to other bones by an immoveable articulation, there are ſmaller ones annexed, which afterwards become ſcarce diſtinguiſhable from the ſubſtance of the bone itſelf. Theſe are called *epiphyſes* or *appendices* (*n*). Some bones have one, others have two, three, or four of thoſe appendices annexed by the means of cartilages, which are of a conſiderable thickneſs in children, but by age become thinner, the oſſification proceeding from the end of the bone on one ſide, and from the epiphyſes on the other, till at laſt, in adults, the place of their conjunction can ſcarcely be ſeen on the external ſurface; and it is only ſometimes that we can then ſee any mark of diſtinction in the cancelli (*o*).

Several proceſſes (*e. g. trochanters* of the thigh, *ſpine* of the ſcapula, &c.) have *epiphyſes*; and proceſſes frequently riſe out from epiphyſes; for example, at the lower end of the femur, ulna, tibia, &c. (*p*).

The epiphyſes are united chiefly to ſuch bones as are deſtined for frequent and violent motion; and for this purpoſe they are wiſely framed of a larger diameter than the bone they belong to: For, by this means, the ſurface of contact between the two bones of any articulation being increaſed, their conjunction becomes firmer, and the muſcles inſerted into them act with greater force by reaſon of their axes being further removed from the centre of motion. Theſe advantages might indeed have been obtained by the expanſion of the end of the bone itſelf to a thickneſs equal to that of the epiphyſes; but then the conſtant ſeparation of new plates to form ſo wide a cellular ſtructure, muſt have left the ſolid ſides of the bones ſo thin as to yield eaſily, either to the action of the muſcles fixed to them and paſſing over them, to the

(*n*) Applantatio, additamentum, adnaſcentia, adnexum, perone.

(*o*) Winſlow, Expoſition anatomique de corps humain, traité des os ſecs, § 116.

(*p*) Veſal. De human. corp. fabrica, lib. 1. cap. 3.

the weight ſeveral of them are obliged to ſupport, or to the application of any other external force.

Several anatomiſts (*q*) thought that the *epiphyſes* ſerve other purpoſes; ſuch as ſecuring the ligaments of the articulations which riſe out from between the bones and them; for, as ſoon as theſe parts are intimately joined, the ligaments inſinuated betwixt them muſt have a much ſtronger connection than they could have to the ſmooth ſurface of the bones. Such an interception of the ligament between the body of the bone and its epiphyſe is not to be ſeen; but, the adheſion of the perioſteum and ligaments to bones being always ſtronger in proportion to the ſimilarity of their conſiſtence, and the bones remaining longer ſoft, or of a ſimilar conſiſtence with ligaments, at theſe places than any where elſe, the opinion of theſe writers, concerning the ſtronger connection of the ligaments where the bones and epiphyſes join, is not without ſome foundation.

Poſſibly too, by the fibres of epiphyſes not extending themſelves ſo longitudinally as thoſe of the bones, there may be leſs chance of the former running into each other than of the latter.

The ſoftneſs of the ends of bones may be of ſome advantage in the womb and in parturition; after which the offification begins at different points to form epiphyſes, before the offification can extend from the middle to the ends of the bones (*r*).

OF OSSIFICATION.

HOWEVER ſolid and compact adult bones are, yet they were once cartilages, membranes, nay, a mere jelly. This needs no further proof, than repeated obſervations of embryos

(*q*) Columb. De re anatomica, lib. 1. cap. 2.——Fallop. Expoſ. de oſſibus, cap. 14.

(*r*) Haller de ſtudio medic. p. 267.

bryos when diſſected: And how much more tender muſt the bones be before that time, when neither knife nor eye is capable to diſcover the leaſt rudiments of them? By degrees they become more ſolid, then aſſume the nature of griſtles, and at laſt oſſify; the coheſion of their plates and fibres always increaſing in proportion to their increaſed ſolidities; as is evident from the time neceſſary to unravel the texture of bones of people of different ages, or of denſe and of ſpongy bones, or of the different parts of the ſame bone, and from the more tedious exfoliations of the bones of adults than of children.

After any part of a bone is fully oſſified, its fibres are extended little more in length at that part, though they increaſe there in thickneſs, and though their ſofter parts continue to become longer (*s*).

As the ſolidity of bones increaſes, their perioſteum more eaſily ſeparates from them. When bones are membranous, the perioſteum and they cannot be diſtinguiſhed; they appear to be the ſame ſubſtance. When they are cartilages, their membrane adheres ſo firmly to them, that it is difficult to ſeparate it from them. Where the bony fibres are rigid, the perioſteum is eaſily taken off.—Is the ſimiliarity of ſtructure and conſequent greater attraction of the membrane and ſubſtance it incloſes, while they are both flexible, the cauſe of their greater adheſion? or is it owing to the veſſels that go from the one to the other being then larger? or do both theſe cauſes combine to produce this effect? or is the membrane or cartilage, which becomes bone afterwards, to be conſidered as the ſame ſubſtance with the perioſteum (*t*)? and muſt all theſe plates of bones be therefore ſaid to be layers of the perioſteum hardened (*u*)?

The

(*s*) Hale's Vegetable ſtatics, p. 293.——Du Hamel. Memoires de l'acad. des ſciences, 1742.

(*t*) Memoires de l'acad. des ſciences, 1744.

(*u*) Ibid. 1743.

The offification of bones depends principally on their veſſels being ſo diſpoſed, and of ſuch diameters, as to ſeparate a liquor, which may eaſily turn into a bony ſubſtance, when it is deprived of its thinner parts; as ſeems plain from the obſervation of the callous matter ſeparated after fractures and ulcers, where part of the bone is taken out: For, in theſe caſes, the veſſels extending themſelves, and the liquors added to them, are gradually formed into granulated fleſh; which fills up all the ſpace where the bone is taken from, then hardens till it becomes as firm as any other part of the bone. This happens frequently, even when the ends of the diſeaſed bone are at a conſiderable diſtance from each other (*v*).

The induration of bones is alſo greatly aſſiſted by their being expoſed, more than any other parts, to the ſtrong preſſure of the great weights they ſupport, to the violent contraction of the muſcles fixed to them, and to the force of the parts they contain, which endeavour to make way for their own further growth. By all this preſſing force, the ſolid fibres and veſſels of bones are thruſt cloſer, and ſuch particles of the fluids conveyed in theſe veſſels as are fit to be united to the fibres are ſooner and more firmly incorporated with them, while the remaining fluids are forcibly driven out by the veins, to be mixed with the maſs of blood. In conſequence of this, the veſſels gradually diminiſh as the bones harden. From which again we can underſtand one reaſon why the bones of young animals ſooner reunite after a fracture than thoſe of old, and why cattle that are put too ſoon to hard labour ſeldom are of ſuch large ſize as others of the ſame brood who are longer kept from labour.

That

(*v*) Hildan. de vuln. graviſ.—Med. eſſays, vol. 1. art. 23.———Job a Meckren, obſ. 69.—Mem. de l'acad. des ſciences, 1742.—See a collection of ſuch caſes in Bochmer de oſſium callo.

That the offifying of bones greatly depends on preſſure, ſeems to be evinced from the frequent examples we meet with of other parts turning bony, when long expoſed to the preſſing force of the ſurrounding parts, or when they are ſubjected to the like circumſtances by their own frequent and violent contraction. Witneſs the bones found frequently near the baſe of the heart in ſome old men (*y*), and in ſeveral other creatures. Nay, the muſcular ſubſtance of the heart has been offified in ſuch (*z*), and the arteries of old men often become bony.——The cartilages of the larynx are generally offified in adults.——In beaſts of burden, the cartilages between the vertebrae of the back very often change into complete bones; and, being intimately united with the vertebrae, the whole appears one continued bone:——Nor is the perioſteum exempted from ſuch an induration (*a*).

To confirm this argument ſtill farther, we may obſerve, that bones begin their offification at the places where they are moſt expoſed to theſe cauſes, viz. in the cylindrical bones, from a middle ring; and in the broad ones, at or near their centre, from one or more diſtinct points. The reaſon of which is, That theſe parts are contiguous to the bellies of the muſcles annexed to the bones, where the ſwelling of theſe moving powers is greateſt. The effects of this may be ſeen in ſome of the bones, as the ſcapula and oſſa ilium, which are covered with muſcles on each ſide; how compact and thin they are in adults, where the bellies of the muſcles were lodged; whereas in children they are thicker. But this being the middle part of theſe bones, where the greateſt number of fibres are, this particular place would

(*y*) Riolan. Comment. de oſſib. cap. 32.—Bartholin. Hiſt. medic. cent. 1. hiſt. 50.——Ibid. cent. 2. hiſt. 45.

(*z*) Cheſelden's Anatomy, book 1. introd.——Garengeot, Hiſt. de l'acad. des ſciences, 1726.

(*a*) Peyer. Ephemerid. German. decur. 2. ann. 7. obſerv. 205.

would have been much thicker in adults, had not this forcible cauſe been applied, which has not had ſuch effects in children, whoſe muſcles have not been much exerciſed.—— Beſides, if we allow that all the parts of a bone are equally increaſed by the conſtant ſupply of new particles, each fibre, and every particle of a fibre, endeavours to make way for its own growth, by puſhing the one next to it; and conſequently by far the greateſt preſſure is on the middle, to make the particles firm, and therefore to begin their oſſification there. Laſtly, the pulſation of the medullary arteries, which enter the bones near to this middle part, may, as authors have alleged, contribute perhaps ſomewhat to this induration.

From the effects of preſſure only it is that we can account for the bones of old people having their ſides much thinner, yet more denſe and ſolid, while the cavities are much larger than in thoſe of young people; and for the prints of muſcles, veſſels, &c. being ſo much more ſtrongly marked on the ſurfaces of the former than of the latter, if they belong to people of near the ſame condition in life.—Preſſure muſt likewiſe be the cauſe which, in people of equal ages, make theſe prints ſtronger in the bones of thoſe who had much labour and exerciſe, than they are in people who have led an indolent and inactive life.

Perhaps both the cauſes of oſſification above mentioned may be aſſiſted by the nature of the climate people live in, and the food they uſe. Whence, in hot countries, the inhabitants ſooner come to their height of ſtature than in the northerly cold regions: And thence ſeems to have ariſen the common practice among the ladies, of making puppies drink brandy or ſpirit of wine, and of bathing them in theſe liquors, to prevent them growing big. Nay, it has been obſerved, that much uſe of ſuch ſpirits has occaſioned parts, naturally

turally ſoft, to petrify in ſome, and to oſſify in other people of no great age (*b*).

From the foregoing account of the ſtructure of bones, and of their oſſification, we may underſtand the reaſons of the following phenomena.

How the natural colour of bones may be changed by ſome ſorts of food (*c*).

Why the bones of ſome people are ſo long in hardening, and in others never completely indurate.

Why, in ſuch whoſe oſſification is ſlow, the bones are generally thicker in proportion to their lengths, eſpecially at their ends; as in the rickets.

How hard firm bones have become ſoft and pliable by diſeaſes (*d*).

Why, in ſome diſeaſes, epiphyſes ſeparate from bones (*e*), and the ends of fractured bones come aſunder many years after their fractures appeared to be cured (*f*).

How bones may waſte and diminiſh (*g*).

How bones may become ſolid all through, without any appearance of cancelli (*h*).

How nodes, tophi, and exoſtoſes, happen after the eroſion

(*b*) Littre, Hiſtoire de l'acad. des ſciences, 1706——Geoffry, Mem. de l'acad. des ſciences, 1706.

(*c*) Philoſoph. tranſact. n° 442. art. 8. n° 443. art. 2. n° 457. art. 4.——Mem. de l'acad. des ſciences, 1739, 1742.

(*d*) Hiſtoire de l'acad. des ſciences, 1704.——Mem. 1722. Gagliardi, Anatom. oſſium, cap. 2. obſerv. 3.——Ephem. Germ. decur. 1. ann. 1. obſ. 37. et ſchol. decur. 2. ann. 7. obſer. 212, 235. decur. 3. ann. 2. obſ. 3.——Philoſ. tranſ. n° 470. § 3. Ibid. vol. 48. § 4. and 44.

(*e*) Memoires de l'acad. des ſciences, 1699. Diemerbroek, lib. 9. cap. 19.—Cowper's Anat. Explic. tab. 96. fig. 1.

(*f*) Anſon's Voyage.

(*g*) Cheſelden's Anat. book 1. introd.——Hiſt. de l'acad. des ſciences, 1700.

(*h*) Ruyſch. Theſaur. 2. arc. 5. theſ. 3. loc. 1. n° 5. theſ. 9. n° 2. not. 3.—Bochmer. de callo oſſium.

ſion of the external plates of bones in the lues venerea, ſcurvy, rheumatiſm, and gout.

How bones exfoliate by the riſing of granulated fleſh from their ſurface.

How and from what callus is formed after a fracture (*i*).

Why callus appears to be rather the continued ſubſtance of the perioſteum than of the bone, while it remains ſoft and flexible; but ſeems continued with the bone after it oſſifies (*k*).

Why callus is ſenſible while it is ſoft, but becomes inſenſible when it hardens.

What occaſions ſometimes ſuch difficulty in curing fractured bones; or why they never re-unite, though they are reduced, and all proper means towards a cure are uſed (*l*) —— Are the bones of women with child more tedious in re-uniting than thoſe of other people (*m*).

Why calluſes, after fractures, are ſometimes very thick and protuberant.

What difference there ought to be in the application of bandages to fractures of the bones of old and of young patients.

How bones, remaining long unreduced after a luxation, may have their form ſo changed as to make their reduction very difficult, if not impoſſible (*n*).

Whoever is deſirous to know in what time and order each bone and its ſeveral parts begin to aſſume a bony na-

(*i*) Memoires de l'acad. des ſciences, 1741.—Dehtleef de oſſium callo.

(*k*) Mem. de l'acad. 1741.

(*l*) Meckren Obſerv. medico-chirurg. obſ. 71.—Ruyſch. Adverſ. dec. 2. § 2. Obſerv. anat.-chir. obſ. 4. —Van Swieten in Boerhaave Aphor. § 354.

(*m*) Hildan. centur. 5. obſ. 87. et cent. 6. obſ. 68.—Philoſoph. tranſact. n°. 494. § 21.

(*n*) Saltzman, Obſ. decur. obſ. 6.—Memoires de l'acad. de chirurgie, tom. 2. p. 155.—Bochmer Inſtit. oſteolog. §. 596.

ture, let him conſult Kerchingius (*o*), who gives us the delineations of abortions from three days after conception, and traces the oſſification of the bones from three weeks and a month, till the time of the birth: To whom ſhould be added Coiterus (*p*) and Eyſſonius (*q*). An account of this ſubject might alſo be collected out of Ruyſch's works, where ſome of the miſtakes committed by the former authors are corrected; and ſeveral more particulars, to make the hiſtory of the oſteogenea more accurate, have ſince been added by Neſbit (*r*) and Albinus (*s*).

I muſt refer to the authors now quoted for the more curious part of the human oſteogeny; not having preparations enough to give ſuch a full hiſtory of it as is done by them. But I ſhall endeavour to explain the more uſeful and neceſſary part of the oſteogeny, by ſubjoining to the deſcription of each bone of an adult, its condition in ripe children; that is, in ſuch as are born at the ordinary time; and ſhall point out what parts of each are afterwards joined in form of epiphyſes. This, with the following general rules, ſeem to me ſufficient for underſtanding as much of this ſubject as is neceſſary in the practice of phyſic and ſurgery.

1. Wherever I mention any parts being cartilaginous, or their being ſtill ſeparable from the other parts of the bone to which they belong, I would be underſtood to hint, that, about ſeven or eight years of age, ſuch parts are oſſified and united to their proper bones, unleſs when it is ſaid that they are afterwards formed into epiphyſes.

2. Such as become epiphyſes are generally oſſified at ſeven or eight years of age; but, being for the moſt part moiſtened

(*o*) Anthropograph. ichnograph. et oſteogenea fœtuum.

(*p*) De oſſibus fœtus abortivi.

(*q*) De oſſibus infant. cognoſcend. et curand.

(*r*) Human oſteogeny expiained.

(*s*) Icones oſſium fœtus humani; accedit oſteogeneæ brevis hiſtoria.

ed by ſynovia, their external ſurface is ſtill ſomewhat cartilaginous, and they are not yet united to their bones.

3. At eighteen or twenty years of age the epiphyſes are entirely oſſified, and have blended their fibres ſo with the body of the bone, as to make them inſeparable without violence.

The knowledge of this part of the oſteogény I think neceſſary to prevent dangerous miſtakes in the cure of ſeveral diſeaſes. As for example: Without this knowledge, the ſeparation of an epiphyſe might be miſtaken for a fracture or luxation.—The interſtice of two parts of a bone not yet joined might be judged to be a fiſſure.—A diaſtaſis, or ſeparation of ſuch disjointed pieces of a bone, might be thought a fracture.—The protruſion of one piece, or its overlopping any other, could be miſtaken for an excreſcence or exoſtoſis.—Such errors about the nature of a diſeaſe would give a perſon very different indications of cure from what he would have if he really underſtood his patient's caſe: And very often the knowledge of the different inequalities on the ſurfaces of bones, muſt direct us in the execution of what is proper to be done to cure ſeveral of their diſeaſes.

OF THE CONNECTION OF BONES.

HAVING thus conſidered the bones when ſingle, we ought next to ſhew the different manner of their conjunctions (*t*). To expreſs theſe, anatomiſts have contrived a great number of technical terms; about the meaning, propriety, and claſſing of which, there has unluckily been variety of opinions. Some of theſe terms, it is neceſſary to retain, ſince they

(*t*) Συνταξις, συνδεσις, συμβολη, ὁμιλια, Compoſitio, connexio, articulatio, conjunctio, nodus, commiſſura, ſtructura, compages.

they ſerve to expreſs the various circumſtances of the articulations, and to underſtand the writers on this ſubject.

The ARTICULATIONS are moſt commonly divided into three claſſes, *viz.* Symphyſis, Synarthroſis, and Diarthroſis.

I. SYMPHYSIS, which properly ſignifies the conception or growing together of parts, when uſed to expreſs the articulations of bones, does not ſeem to comprehend, under the meaning generally given to it, any thing relating to the form or motion of the conjoined bones; but by it moſt authors only denote the bones to be connected by ſome other ſubſtance; and as there are different ſubſtances which ſerve this purpoſe, therefore they divide it into the three following ſpecies.

1. *Synchondroſis* (*u*), when a cartilage is the connecting ſubſtance: thus the ribs are joined to the ſternum; thus the bodies of the vertebrae are connected to each other; as are likewiſe the oſſa pubis.

2. *Synneuroſis* or *ſyndeſmoſis*, when ligaments are the connecting bodies, as they are in all the moveable articulations.

3. *Syſſarcoſis*, when muſcles are ſtretched from one bone to another, as they muſt be where there are moveable joints.

II. The ſecond claſs of articulations, the SYNARTHROSIS, which is ſaid to be the general term by which the immoveable conjunction of bones is expreſſed, is divided into three kinds.

1. The *future* (*x*) is that articulation where two bones are mutually indented into each other, or as if they were ſewed

(*u*) Amphiarthroſis.

(*x*) Ραφη.

ſewed together; and is formed by the fibres of two bones meeting while they are yet flexible and yielding, and have not come to their full extent of growth; ſo that they mutually force themſelves into the interſtices of each other, till, meeting with ſuch reſiſtance as they are not able to overcome, they are ſtopped from ſprouting out farther, or are reflected; and therefore theſe indentations are very different both in figure and magnitude: thus the bones of the head are joined; thus epiphyſes are joined to the bones, before their full connection and union with them.

Under this title of *ſuture*, the *harmonia* of the ancients may be comprehended; ſcarce any unmoved bones being joined by plain ſurfaces (*y*).

2. *Gomphoſis* (*z*) is the fixing one bone into another, as a nail is fixed in a board: thus the teeth are ſecured in their ſockets.

3. *Schindyleſis* or *ploughing* (*a*), when a thin lamella of one bone is received into a long narrow furrow of another: thus the proceſſus azygos of the ſphenoid, and the naſal proceſs of the ethmoid bone, are received by the vomer.

III. The third claſs, or DIARTHROSIS (*b*), is the articulation where the bones are ſo looſely connected as to allow large motion. This is ſubdivided into three kinds.

1. *Enarthroſis*, or the ball and ſocket, when a large head is received into a deep cavity; as the head of the os femoris is into the acetabulum coxendicis.

2. *Arthrodia*, when a round head is received into a ſuperficial cavity; as in the articulation of the arm bone and ſcapula. Theſe two ſpecies of diarthroſis allow motion to all ſides.

3. *Gin-*

(*y*) Veſal. obſerv. Fallop. examen.

(*z*) Conclavatio.

(*a*) Keil's Anat. chap. 5.

(*b*) Απαρθρωσις, Dearticulatio, abarticulatio.

3. *Ginglimus* (*c*), which properly ſignifies the hinge of a door or window; in it the parts of the bones mutually receive and are received, and allow of motion two ways: Workmen call it *charnal.*

The *ginglimus* is generally divided into three kinds; to which ſome (*d*) give the name of *contiguous* (*e*), *diſtant* (*f*), and *compound* (*g*).

The firſt kind of ginglimus is when a bone has ſeveral protuberances and cavities, which anſwer to as many cavities and proceſſes of the other bone with which it is articulated; as in the conjunction of the femur with the tibia.

The ſecond ſpecies is, when a bone receives another at one end, and is received by the ſame bone at the other end; as in the radius and ulna.

The laſt ſort is, when a bone receives another, and is received by a third; as in the oblique proceſſes of the vertebrae.

When I firſt mentioned the articulations of bones, I ſaid there were different opinions concerning the uſe of their technical names; *e. g.* It has been ſaid, that *ſymphyſis* ſhould be the name for the immoveable articulations, and *ſynarthroſis* ſhould be underſtood to be the conjunction of bones by ſome connecting medium.——Thoſe who have taken *ſymphyſis* in the ſenſe I did, of its expreſſing the conjunction of bones with a connecting ſubſtance, have diſagreed in their definition of it; ſome inſerting, and others leaving out, its allowing motion. When they have agreed in their definition, they have not been of the ſame mind concerning the ſpecies of it; for ſeveral think the *ſyſſarcoſis* and *ſyndeſmoſis* applicable to ſo many joints which are univerſally allowed to be claſſed

(*c*) Articulatio mutua.
(*d*) Baker, Curſ. Oſteolog. demonſt. 1.
(*e*) Proximus.
(*f*) Longus.
(*g*) Compoſitus.

claſſed under the *diarthroſis*, that it muſt create confuſion to name them by any ſpecies of the *ſymphyſis*. Few keep to ſuch a general definition of the ſynchondroſis as I have done; and whether they determine it to allow no motion, or an obſcure or a manifeſt one, they bring themſelves into difficulties, becauſe there are examples of all theſe three kinds.——Some again, by too nicely diſtinguiſhing obſcure and manifeſt motions of bones, have blended the *ſynarthroſis* and *diarthroſis*; and from thence have branched out the different compound ſpecies of articulations that may be formed of them, ſo far, that they could find no examples in the body to illuſtrate them by.—It would be tedious to enumerate more of the jarring opinions, and it would be far more ſo to give a detail of the arguments uſed by the diſputants. It is ſufficient for my purpoſe that it is underſtood in what ſenſe I take theſe technical terms; which I do in the following manner.

When I mention the ſymphyſis or ſynarthroſis, or any ſpecies of them, I ſhall always underſtand them according to the explication already given of them. But though the preceding account of the diarthroſis, or articulation of moveable bones, has been almoſt univerſally received; yet as it does not comprehend all the moveable articulations of the body, and one of its ſpecies does not anſwer to any notion we can have of the conjunction of two bones, I muſt beg leave to change the definitions and kinds of theſe joints.

I would call *diarthroſis*, That conjunction of bones whereby they are fitted for motion, being each covered with a ſmooth cartilage, connected by one or more common ligaments, and lubricated with liquor at the conjoined parts: In which definition I have no regard to the quantity of motion which they really do perform; the motion being often confined or enlarged by ſome other cauſe not immediately

diately depending on the frame of the two ſurfaces of the bones forming the particular joint, which then is conſidered.

The firſt ſpecies of the diarthroſis, viz. the *enarthroſis*, or ball and ſocket, I would define more generally than above, That articulation where a round head of one bone is received into a cavity of another; and conſequently, without ſome foreign impediment, is capable of motion to all ſides. Examples of this kind are to be ſeen in the articulation of the thigh-bone and oſſa innominata; arm-bone and ſcapula; aſtralagus and os naviculare; magnum of the wriſt with the ſcaphoides and lunare; firſt bone of the thumb with the ſecond, &c.

The ſecond ſort, or the *arthrodia*, differing from the enarthroſis, in the preceding account, only in the cavity's being more ſuperficial, (which makes no eſſential difference, eſpecially that, in the recent ſubject, cartilages or ligaments ſupply the deficiency of bone), ought, in my opinion, to be called with Veſalius (*b*), that articulation of two bones adapted for motion, where it is not at firſt ſight obvious which of the two has the head or cavity, or where they are joined by plain ſurfaces, or nearly ſo; ſuch is the conjunction of the clavicle with the ſcapula, oſſa cuneiformia with the os naviculare, metatarſal bones with the oſſa cuneiformia, &c. From the nature of this ſort of joint it is plain, that very great motion cannot be allowed, without the bones going further out of their natural ſituation than is convenient or ſafe.

Ginglimus, I would reckon that articulation by the form of which the motion of the joined bones muſt be chiefly confined to two directions, as hinges of doors are.

The firſt ſpecies of this is the *trochoides*, when one bone turns

(*b*) De corp. human. fabrica, lib. 1. cap. 4.

turns on another, as a wheel does on its axis. Thus the firſt vertebra of the neck moves on the tooth-like proceſs of the ſecond. This is the moſt proper kind of ginglimus.

The ſecond ſpecies ſhould be eſteemed the articulation where ſeveral prominent and hollow ſurfaces of two bones move on each other within the ſame common ligament; as in the knee, &c.

The third ſort of ginglimus is, when two bones are articulated to each other at different parts, with a diſtinct apparatus of the motory machines at each; ſuch is the articulation of the os occipitis with the firſt vertebra of the neck; of any two contiguous vertebrae by their oblique proceſſes; of the ribs with the bodies of tranſverſe proceſſes of the vertebrae; of the radius with the ulna, tibia with the fibula, aſtragalus calcaneum, &c.

I would entirely throw out what is commonly called the third kind of ginglimus; for in examining the conjunction of a bone with two others, as in the common example of a vertebra joined with the one above and below, the connection of the middle one with each of the other two ought to be conſidered ſeparately; otherwiſe we might, with the ſame propriety, eſteem the articulations that the long bones, the femur, tibia, humerus, &c. have at their different ends, as one articulation; which is abſurd.

If the moveable bones are not connected and kept firm by ſome ſtrong ſubſtance, they would be luxated at every motion of the joints; and if their hard rough unequal ſurfaces were to play on each other, their motion would not only be difficult, but the loſs of ſubſtance from attrition would be great. Therefore ligaments are made to obviate the firſt, and cartilages to prevent the other inconveniency. But becauſe ligaments and cartilages turn rigid, inflexible, and rough, unleſs they are kept moiſt, a ſufficient quantity of proper liquors is ſupplied for their lubrication, and to

preſerve them in a flexible ſtate. Seeing, then, theſe parts are ſo neceſſary to the articulations, I ſhall next conſider their ſtructure, ſituation, and uſes, ſo far as they are ſubſervient to the bones and their motions.

OF THE LIGAMENTS.

Ligaments (*i*) are white flexible bodies, thicker and firmer than membranes, and not ſo hard or firm as cartilages, without any remarkable cavity in their ſubſtance, difficultly ſtretched, and with little elaſticity; ſerving to connect one part to another, or to prevent the parts to which they are fixed from being removed out of that ſituation which is uſeful and ſafe.

After maceration in water, the ligaments can eaſily be divided; and each ligamentous layer appears compoſed of fibres, the largeſt of which are diſpoſed in a longitudinal direction.

The arteries of ligaments are very conſpicuous after a tolerable injection, and the larger trunks of their veins are ſometimes to be ſeen full of blood.

Such ligaments as form the ſides of cavities, have numerous orifices of their arteries opening upon their internal ſurface, which keep it always moiſt. If we rub off that moiſture, and then preſs the ligament, we can ſee the liquor oozing out from ſmall pores; and we can force thin liquors injected by the arteries into the cavities formed by ligaments.

Theſe exhalent arteries muſt have correſponding abſorbents, otherwiſe the cavities would ſoon be too full of liquor.

Ligaments, then, muſt be ſubject to the diſeaſes common to other parts, where there is a circulation of fluids, allowance

(*i*) Συνδεσμοι, νευρα, copulae, vincula.

allowance always being made for the ſize of veſſels, the nature of the fluids, and the firmneſs of the texture of each part.

Authors generally ſay that ligaments are inſenſible; and conſequently it may be inferred, that they have no nerves beſtowed on them. But the violent racking pain felt on the leaſt motion of a joint labouring under a rheumatiſm, the ſeat of which diſeaſe ſeems often to be in the ligaments, and the inſufferable torture occaſioned by inciſions of ligaments, and by a collection of acrid matter in a joint, or by tophi in the gout, would perſuade us that they are abundantly ſupplied with nerves.

The ligaments which connect the moveable bones commonly riſe from the conjunction of the epiphyſes of the one bone, and are inſerted into the ſame place of the other; or, where epiphyſes are not, they come out from the cervix, and beyond the ſupercilia of the articulated bones; and after ſuch a manner, in both caſes, as to include the articulation in a purſe or bag; with this difference, depending on their different motions, that where the motion is only to be in two directions, the ligaments are ſtrongeſt on thoſe ſides towards which the bones are not moved; and when a great variety of motion is deſigned to be allowed, the ligaments are weaker than in the former ſort of articulations, and are nearly of the ſame ſtrength all round.

Part of the capſular ligaments is compoſed of the perioſteum, continued from one bone to another, as was obſerved, p. 6. and their internal layer is continued on the parts of the bone or cartilage which the ligament includes (*k*).

Beſides theſe common capſular ligaments of the joints, there are particular ones in ſeveral places, either for the firmer connection of the articulated bones, or for reſtraining

(*k*) Neſbit, Oſteogen.—Philoſ. tranſact. N°. 470. § 6.

ing and confining the motion to ſome one ſide; ſuch are the *croſs* and *lateral* ligaments of the knee, the *round* one of the thigh, &c.

From this account of the ligaments, we may conclude, that, *cæteris paribus*, in whatever articulation the ligaments are few, long, and weak, the motion is more free and quick; but luxations happen frequently: and, on the contrary, where the ligaments are numerous, ſhort, and ſtrong, the motion is more confined; but ſuch a joint is leſs expoſed to luxations (*l*).——Whence we may judge how neceſſary it is to attend to the different ligaments, and the changes which have been made on them by a luxation, when it is to be reduced.

Ligaments alſo ſupply the place of bones in ſeveral caſes to advantage. Thus the parts in the pelvis are more ſafely ſupported below by ligaments than they could have been by bone.——The ligaments placed in the great holes of the oſſa innominata and between the bones of the fore-arm and leg, afford convenient origin to muſcles.——Immoveable bones are firmly connected by them; of which the conjunction of the os ſacrum and innominatum is an example.——They afford a ſocket for moveable bones to play in, as we ſee part of the aſtragalus does on the ligament ſtretched from the heel-bone to the ſcaphoid.

Numerous inconveniencies may ariſe from too long or ſhort, ſtrong or weak, lax or rigid ligaments.

OF THE CARTILAGES.

CARTILAGES (*m*) are ſolid, ſmooth, white, elaſtic ſubſtances, between the hardneſs of bones and ligaments, and covered with a membrane named *perichondrium*; which is of

(*l*) Fabric. ab. Aquapend. de articul. part. utilit. pars 3.

(*m*) Χονδροι.

of the ſame ſtructure and uſe to them as the perioſteum is to the bones.

Cartilages are compoſed of plates, which are formed of fibres, diſpoſed much in the ſame way as thoſe of bones are; as might be reaſonably concluded from obſerving bones in a cartilaginous ſtate before they oſſify, and from ſeeing, on the other hand, ſo many cartilages become bony. This may be ſtill further confirmed by the exfoliation which cartilages are ſubject to as well as bones.

The perichondrium of ſeveral cartilages, for example thoſe of the ribs and larynx, has arteries, which can be equally well injected with thoſe of the perioſteum: but the veſſels of that membrane in other parts, *e. g.* the articular cartilages, are ſmaller, and in none of them does injection enter deep into the ſubſtance of the cartilages; nay madder, mixed with the food of animals, does not change the colour of cartilages as it does that of bones (*n*).

The granulated fleſh which riſes from the ends of metacarpal or metatarſal bones, when the cartilage exfoliates, after a finger or toe has been taken off at the firſt joint, is very ſenſible; from which the exiſtence of nerves in cartilages may be inferred.

While cartilages are in a natural ſtate, it is to be remarked, firſt, That they have no cavity in their middle for marrow. Secondly, That their outer ſurface is ſofteſt, which renders them more flexible. Thirdly, That they do not appear to change their texture near ſo much by acids as bones do. And, laſtly, That as the ſpecific gravity of cartilages is near a third leſs than that of bones; ſo the coheſion of their ſeveral plates is not ſo ſtrong as in bones: Whence cartilages laid bare in wounds or ulcers, are not only more liable to corrupt, but exfoliate much ſooner than bones do.

Cartilages

(*n*) Philoſ. Tranſact. N° 442. art. 8. N° 443. art. 2. N° 457. art. 4. Mem. de l'acad. des ſciences, 1739 et 1742.—Dehtleef de oſſium callo.

Cartilages ſeem to be principally kept from oſſifying, either by being ſubjected to alternate motions of flexion and extenſion, the effects of which are very different from any kind of ſimple preſſure; or by being conſtantly moiſtened (*o*). Thus the cartilages on the articulated ends of the great bones of the limbs, and the moveable ones placed between the moving bones in ſome articulations, which are obliged to ſuffer many and different flexions, and are plentifully moiſtened, ſcarce ever change into bone; while thoſe of the ribs and larynx are often oſſified. The middle angular part of the cartilages of the ribs, which is conſtantly in an alternate ſtate of flexion and extenſion, by being moved in reſpiration, is always the laſt of becoming bony. In the larynx, the epiglottis, which is oftener bended and more moiſtened than the other four cartilages, is ſeldom oſſified, while the others as ſeldom eſcape it in adults.

The cartilages ſubſervient to bones are ſometimes found on the ends of bones which are joined to no other; but are never wanting on the ends and in the cavities of ſuch bones as are deſigned for motion (*p*). Cartilages alſo are interpoſed between ſuch other cartilages as cover the heads and cavities of articulated bones; nay, they are alſo placed between immoveable bones.

The *uſes* of cartilages, ſo far as they regard bones, are, to allow, by their ſmoothneſs, ſuch bones as are deſigned for motion, to ſlide eaſily, and without detrition, while, by their flexibility, they accommodate themſelves to the ſeveral figures neceſſary in different motions, and, by their elaſticity, they recover their natural poſition and ſhape as ſoon as the preſſure is removed.——This ſpringy force may alſo render the motion of the joints more expeditious, and may likewiſe leſſen the ſhocks in running, jumping, &c.——To theſe cartilages

(*o*) Havers Oſteolog. Nov.

(*p*) Celſ. de re medic. lib. 8. cap. 1.

tilages we chiefly owe the ſecurity of the moveable articulations; for, without them, the bony fibres would ſprout out, and intimately coaleſce with the adjoining bone; whence a true anchyloſis muſt neceſſarily follow; which always happens when the cartilages are eroded by acrid matter, or oſſified from want of motion or defect of liquor, as is frequently the caſe after wounds of the joints, paidarthrocace, ſcrophula, and ſpina ventoſa, or from old age, and long immobility of joints (*q*). Hence we may know what the annihilation is, which is ſaid to be made of the head of a bone, and of the cavity for lodging it, after an unreduced fracture (*r*). The moveable cartilages interpoſed in joints ſerve to make the motions both freer and more ſafe than they would otherwiſe be. Thoſe placed on the ends of bones that are not articulated, as on the ſpine of the os ilium, baſe of the ſcapula, &c. ſerve to prevent the bony fibres from growing out too far. Cartilages ſometimes ſerve as ligaments, either to faſten together bones that are immoveably joined, ſuch are the cartilages between the os ſacrum and oſſa ilium, the oſſa pubis, &c. or to connect bones that enjoy manifeſt motion, as thoſe do which are placed between the bodies of the true vertebrae, &c. Cartilages very often do the office of bones to greater advantage than theſe laſt could; as in the cartilages of the ribs, thoſe which ſupply brims to cavities, &c.

Too great thickneſs or thinneſs, length or ſhortneſs, hardneſs or ſuppleneſs, of cartilages, may therefore cauſe great diſorders in the body.

OF

(*q*) Columb. de re anat. lib. 15.—Deſlandes, Hiſt. de l'acad. des ſciences, 1716.—Phil. Tranſ. N° 215.—Ibid. N° 461. ſect. 16.

(*r*) Hildan. de ichor. et melicer. acri Celſi, cap. 5.——Ruyſch. Theſ. 8. N° 103.——Saltzman, in Act. Petropolit. tom. 3. p. 275.

OF THE SYNOVIA.

THE liquor which principally ſerves to moiſten the ligaments and cartilages of the articulations is ſupplied by glands, which are commonly ſituated in the joint, after ſuch a manner as to be gently preſſed, but not deſtroyed by its motion. By this means, when there is the greateſt neceſſity for this liquor, that is, when the moſt frequent motions are performed, the greateſt quantity of it muſt be ſeparated. Theſe glands are ſoft and pappy, but not friable: in ſome of the large joints they are of the conglomerate kind, or a great number of ſmall glandules are wrapt up in one common membrane. Their excretory ducts are long, and hang looſe, like ſo many fringes, within the articulation; which, by its motion and preſſure, prevents obſtructions in the body of the gland or its excretories, and promotes the return of this liquor, when fit to be taken up by the abſorbent veſſels, which muſt be in the joints, as well as in the other cavities in the body; and, at the ſame time, the preſſure on the excretory ducts hinders a ſuperfluous unneceſſary ſecretion, while the fimbriated diſpoſition of theſe excretories does not allow any of the ſecreted liquor to be puſhed back again by theſe canals towards the glands (*s*).

Very often theſe fountains of ſlimy liquor appear only as a net-work of veſſels.—Frequently they are almoſt concealed by cellular membranes containing the fat—and ſometimes ſmall ſimple mucous folliculi may be ſeen (*t*).

The different joints have theſe organs in different numbers and ſizes: the conglomerate ones do not vary much, eſpecially as to ſituation, in the ſimilar joints of different bodies; but the others are more uncertain.

Upon

(*s*) Cowper, Anatom. explicat. tab. 79. lit. E. E.

(*t*) Morgagn. Adverſar. 2. animad. 23.

Upon preſſing any of theſe glands with the finger, a mucilaginous liquor may be ſqueezed out of their excretories, which ſomewhat reſembles the white of an egg or ſerum of the blood; but it is manifeſtly ſalt to the taſte. It does not coagulate by acids nor by heat, as the ſerum does; but by the latter turns firſt thinner, and, when evaporated, leaves only a thin ſalt film.

The quantity of this mucilage, conſtantly ſupplied, muſt be very conſiderable, ſince we ſee what a plentiful troubleſome diſcharge of glary matter follows a wound or ulcer of any joint: of which liquor the mucilage is a conſiderable part.

The veſſels which ſupply liquors for making the ſecretion of this mucilage, and the veins which bring back the blood remaining after the ſecretion, may be ſeen without any preparation; and, after a tolerable injection of the arteries, the glands are covered with them.

In a ſound ſtate, we are not conſcious of any ſenſibility in thoſe glands: but, in ſome caſes which I have ſeen, when they inflame and ſuppurate, the moſt racking pain is felt in them; a melancholy, though a ſure, proof that they have nerves.

Theſe mucilaginous glands are commonly lodged in a cellular ſubſtance; which is alſo to be obſerved in other parts of the bag formed by the ligaments of the articulation; and which contains a fatty matter, that muſt neceſſarily be attenuated, and forced through the including membranes into the cavity of the joint, by the preſſure which it ſuffers from the moving bones.

If, then, the oil is conveyed from this cellular ſubſtance; and if the attenuated marrow paſſes from the cancelli of the bones by the large pores near their ends, or in their cavities, and ſweats through the cartilages there into the articulations; which it may, when aſſiſted by the conſtant

heat and action of the body, more eaſily do, than when it eſcapes through the compact ſubſtance of the bones in a ſkeleton: If, I ſay, this oil is ſent to a joint, and is incorporated with the mucilage, and with the fine lymph that is conſtantly oozing out at the extremities of the ſmall arteries diſtributed to the ligaments, one of the fitteſt liniments imaginable muſt be produced; for the mucus diluted by the lymph contributes greatly to its lubricity, and the oil preſerves it from hardening. How well ſuch a mixture ſerves the purpoſe it is deſigned for, Boyle (*u*) tells us he experienced in working his air-pump; for the ſucker could be moved with much leſs force after being moiſtened with water and oil, than when he uſed either of theſe liquors ſingly. And I believe every one, at firſt view, will allow the diluted mucilage to be much preferable to ſimple water. The ſynovia (*x*), as this liquor compoſed of oil, mucilage, and lymph, is commonly now called, while in a ſound ſtate, effectually preſerves all the parts concerned in the articulations ſoft and flexible, and makes them ſlide eaſily on each other, by which their mutual detrition and overheating is prevented, in the manner daily practiſed in coach and cart wheels by beſmearing them with greaſe and tar.

After the liquor of the articulations becomes too thin and unſerviceable, by being conſtantly pounded and rubbed between the moving bones, it is reaſſumed into the maſs of blood by the abſorbent veſſels.

When the ſynovia is not rubbed betwixt the bones, it inſpiſſates. And ſometimes, when the head of bone has been long out of its cavity, this liquor is ſaid to fill up the place of the bone, and hinder its reduction; or if a joint continues long unmoved, it is alſo ſaid to cement the bones, and

(*u*) Phyſico-mechanic. experim.

(*x*) Μυξα, mucus, axungia.

and occaſion a true anchyloſis (*y*).——If the ſynovia becomes too acrid, it erodes the cartilages and bones; as frequently happens to thoſe who labour under the lues venerea, ſcurvy, ſcrophula, or ſpina ventoſa.——If this liquor is ſeparated in too ſmall a quantity, the joint becomes ſtiff; and, when with difficulty it is moved, a crackling noiſe is heard, as people advanced in years frequently experience (*z*).——If the mucilage and lymph are depoſited in too great a quantity, and the abſorbent veſſels do not perform their office ſufficiently, they may occaſion a dropſy of the joints (*a*).——From this ſame cauſe alſo the ligaments are often ſo much relaxed, as to make the conjunction of the bones very weak: Thence ariſe the luxations from an internal cauſe, which are eaſily reduced, but difficultly cured (*b*). Frequently, when ſuch a ſuperfluous quantity of this liquor is pent up, it becomes very acrid, and occaſions a great train of bad ſymptoms; ſuch as ſwelling and pain of the joints, long ſinous ulcers, and fiſtulae, rotten bones, immobility of the joints, marcor and atrophia of the whole body, hectic fevers, &c. (*c*).——From a depravity in the blood, or diſeaſes in the organs that furniſh the ſynovia of the joints, it may be greatly changed from its natural ſtate; it may be purulent after inflammation, mucous in the white ſwellings, gelatinous in the rheumatiſm, chalky from the gout, &c.: hence a great variety of diſorders in the joints (*d*).

CHAP.

(*y*) Paré, Chirurgie, livre 15. chap. 18. et livre 16. chap. 5.

(*z*) Galen. de uſu part. lib. 12. cap. 2.——Fabric. ab Aquapend. de articul. part. utilitat. pars 3.——Bartholin. hiſt. medic. cent. 3. hiſt. 11.

(*a*) Hildan. de ichore et melicer. acri Celſi.

(*b*) Hippocrat. de locis in homine, ſect. 14. et de articul.

(*c*) Hildan. de ichore et meliceria acri Celſi.

(*d*) See Reimar, Diſſert. de fungo articulor.

CHAP. II.

Of the SKELETON.

THOUGH any dry ſubſtance may be called *ſkeleton*, yet, among anatomiſts, this word is univerſally underſtood to ſignify the bones of animals connected together, after the teguments, muſcles, bowels, glands, nerves, and veſſels, are taken away (*a*).

A ſkeleton is ſaid to be a *natural* one when the bones are kept together by their own ligaments; and it is called *artificial* when the bones are joined with wire, or any other ſubſtance which is not part of the creature to which they belonged. Small ſubjects, and ſuch whoſe bones are not fully oſſified, are commonly prepared the firſt way; becauſe, were all their parts divided, the niceſt artiſt could not rejoin them, by reaſon of their ſmallneſs, and of the ſeparation of their unoſſified parts; whereas the bones of large adult animals are ſooneſt and moſt conveniently cleaned when ſingle, and are eaſily reſtored to and kept in their natural ſituation.——Sometimes the ſkeleton of the ſame animal is prepared in both theſe ways; that is, the ſmaller bones are kept together by their natural ligaments, and the larger ones are connected by wires or ſome ſuch ſubſtances.

Before we proceed to the diviſion and particular deſcription of the ſkeleton, it is to be obſerved, that, when the bones are put into their natural ſituation, ſcarce any one of them is placed in a perpendicular bearing to another; though the fabric compoſed of them is ſo contrived, that, in an erect poſture, a perpendicular line, for their common centre of gravity, falls in the middle of their common

(*a*) Cadaveris crates.

mon baſe (*b*). On this account, we can ſupport ourſelves as firm as if the axis of all the bones had been a ſtraight line perpendicular to the horizon; and we have much greater quickneſs, eaſe, and ſtrength, in ſeveral of the moſt neceſſary motions we perform. It is true, indeed, that wherever the bones, on which any part of our body is ſuſtained, decline from a ſtraight line, the force required in the muſcles to counteract the gravity of that part is greater than was otherwiſe neceſſary: But then this is effectually provided for in ſuch places, by the number and ſtrength of the muſcles. So long, therefore, as we remain in the ſame poſture, a conſiderable number of muſcles muſt be in a conſtant ſtate of contraction; which we know, both from reaſon and experience, muſt ſoon create an uneaſy ſenſation. This we call *being weary of one poſture:* An inconvenience that we ſhould not have had in ſtanding erect, if the bearing of all the bones to each other had been perpendicular; but it is more than compenſated by the advantages above-mentioned.

The human ſkeleton is generally divided into the HEAD, the TRUNK, the SUPERIOR and the INFERIOR EXTREMITIES.

SECT. I. OF THE HEAD.

BY the HEAD is meant all that ſpheroidal part which is placed above the firſt bone of the neck. It therefore comprehends the cranium and bones of the face.

§ 1. *The* CRANIUM.

THE cranium (*c*), helmet, or brain-caſe, conſiſts of ſeveral pieces, which form a vaulted cavity, for lodging and defending

(*b*) Cowper, Anat. of human bodies, explic. of tab. 87. 88.

(*c*) Κολχος, κυτος, κωδιξ, εγκαφιον, calva, calvaria, cerebri galea, theca, et olla capitis, teſta capitis, ſcutella capitis.

defending the brain and cerebellum, with their membranes, veſſels, and nerves.

The cavity of the cranium is proportioned to its contents. Hence ſuch a variety of its ſize is obſerved in different ſubjects; and hence it is neither ſo broad nor ſo deep at its fore-part, in which the anterior lobes of the brain are lodged, as it is behind, where the large poſterior lobes of the brain, and the whole cerebellum, are contained.

The roundiſh figure of the ſkull, which makes it more capacious, and better able to defend its contents from external injuries, is chiefly owing to the equal preſſure of theſe contained parts as they grow and increaſe before it is entirely oſſified.——It is to be obſerved, however, that the ſides of the cranium are depreſſed below a ſpherical ſurface by the ſtrong temporal muſcles, whoſe action hinders here the uniform protruſion of the bones, which is more equally performed in other parts where no ſuch large muſcles are. In children whoſe muſcles have not acted much, and conſequently have not had great effects on the bones, this depreſſion is not ſo remarkable; and therefore their heads are much rounder than in adults. Theſe natural cauſes, differently diſpoſed in different people, produce a great variety in the ſhapes of ſkulls, which is ſtill increaſed by the different management of the heads of children when very young: So that one may know a Turk's ſkull by its globular figure, a German's by its breadth and flatneſs of the occiput, Dutch and Engliſh by their oblong ſhapes, &c. (*d*). Two advantages are reaped from this flatneſs of the ſides of the cranium, viz. the enlargement of our ſphere of viſion, and more advantageous ſituation of our ears for receiving a greater quantity of ſound, and for being leſs expoſed to injuries.

The

(*d*) Veſal. lib. 1. c. 5.

The external ſurface of the upper part of the cranium is very ſmooth and equal, being only covered with the perioſteum (common to all the bones, but in the ſkull diſtinguiſhed by the name of *pericranium*), the thin frontal and occipital muſcles, their tendinous aponeuroſis, and with the common teguments of the body; while the external ſurface of its lower part has numerous riſings, depreſſions, and holes, which afford convenient origin and inſertion to the muſcles that are connected to it, and allow ſafe paſſage for the veſſels and nerves that run through and near it.

The internal ſurface of the upper part of the ſkull is commonly ſmooth, except where the veſſels of the dura mater have made furrows in it, while the bones were ſoft.—— Surgeons ſhould be cautious when they trepan here, leſt in ſawing or raiſing the bone where ſuch furrows are, they wound theſe veſſels.——In the upper part of the internal ſurface of ſeveral ſkulls, there are likewiſe pits of different magnitudes and figures, which ſeem to be formed by ſome parts of the brain being more luxuriant and prominent than others. In theſe pits, the ſkull is ſo much thinner than any where elſe, that it is often rendered diaphanous, the two tables being cloſely compacted without a diploë; the want of which is ſupplied by veſſels going from the dura mater into a great many ſmall holes obſervable in the pits. Theſe veſſels are larger, and much more conſpicuous than any others that are ſent from the dura mater to the ſkull; as evidently appears from the drops of blood they pour out, when the ſkull is raiſed from the dura mater in a recent ſubject; and therefore they may furniſh a ſufficient quantity of liquors neceſſary to prevent the brittleneſs of this thin part.—The knowledge of theſe pits ſhould teach ſurgeons to ſaw cautiouſly and ſlowly through the external table of the ſkull, when they are performing the operation of the trepan; ſince,

ſince, in a patient whoſe cranium has theſe pits, the dura mater and brain may be injured, before the inſtrument has pierced near the ordinary thickneſs of a table of the ſkull.—The internal baſe of the ſkull is extremely unequal, for lodging the ſeveral parts and appendices of the brain and cerebellum, and allowing paſſage and defence to the veſſels and nerves that go into or come out from theſe parts.

The bones of the cranium are compoſed of two tables, and intermediate cancelli, commonly called their *diploë* (*e*). The external table is thickeſt; the inner, from its thinneſs and conſequent brittleneſs, has got the name of *vitrea*. Whence we may ſee the reaſon of thoſe miſchievous conſequences which ſo often attend a collection of matter in the diploë, either from an external or internal cauſe, before any ſign of ſuch a collection appears in the teguments which cover that part of the ſkull where it is lodged (*f*).

The diploë has nearly the ſame texture and uſes in the ſkull with the cancelli in other bones.

The diploë of ſeveral old ſubjects is ſo obliterated, that ſcarce any veſtige of it can be ſeen; neither is it obſervable in ſome of the hard craggy bones at the baſe of the ſkull. Hence an uſeful caution to ſurgeons who truſt to the bleeding, want of reſiſtance, and change of ſound, as certain marks, in the operation of the trepan, that their inſtrument has ſawed through the firſt table, and reached the diploë (*g*). In other people, the diploë becomes of a monſtrous thickneſs, while the tables of the ſkull are thinner than paper.

The cranium conſiſts of eight bones, ſix of which are ſaid to be proper, and the other two are reckoned common to

(*e*) Meditulium, commiſſura.

(*f*) Boneti Sepulchret. anat. lib. 1. § 1. obſ. 96.—103.

(*g*) Bartholin. Anat. reform. lib. 4. cap. 4.

to it and to the face.—The ſix proper are the *os frontis*, two *oſſa parietalia*, two *oſſa temporum*, and the *os occipitis*.—The common are the *os ethmoides* and *ſphenoides*.

The *os frontis* forms the whole fore-part of the vault; the two *oſſa parietalia* form the upper and middle part of it; the *oſſa temporum* compoſe the lower part of the ſides; the *os occipitis* makes the whole hinder part, and ſome of the baſe; the *os ethmoides* is placed in the fore-part of the baſe, and the *os ſphenoides* in the middle of it.

THE SUTURES.

THE above bones are joined to each other by five ſutures; the names of which are the *coronal*, *lambdoidal*, *ſagittal*, and two *ſquamous*.

The *coronal* (*h*) ſuture is extended over the head, from within about an inch of the external canthus of one eye, to the like diſtance from the other; which being near the place where the ancients wore their vittæ, coronæ, or garlands, this ſuture has thence got its name.—Though the indentations of this ſuture are conſpicuous in its upper part, yet an inch or more of its end on each ſide has none, but is ſquamous and ſmooth.

The *lambdoidal* (*i*) ſuture begins ſome way below, and farther back than the vertex or crown of the head, whence its two legs are ſtretched obliquely downwards and to each ſide in form of Greek letter Λ, and are now generally ſaid to extend themſelves to the baſe of the ſkull: but formerly anatomiſts (*k*) reckoned the proper lambdoidal ſuture to terminate at the ſquamous ſutures; and what is extended at an angle down from that on each ſide, where the indentations

(*h*) Στεφανιαια, Arcualis, puppis.

(*i*) Laudæ, proræ, hypſyloides.

(*k*) Veſal. Anat. lib. I. cap. 6.

are leſs conſpicuous than in the upper part of the future, they called *additamentum futuræ lambdoidis* (*l*).

This future is ſometimes very irregular, being made up of a great many ſmall futures, which ſurround ſo many little bones that are generally larger and more conſpicuous on the external ſurface of the ſkull than internally. Theſe bones are commonly called *triquetra* or *Wormiana:* but ſome other name ought to be given them, for they are not always of a triangular figure, and older anatomiſts (*m*) than Olaus Wormius (*n*) have deſcribed them.—The ſpecific virtue which theſe bones were once thought to have in the cure of the epilepſy (*o*), is not aſcribed to them now; and anatomiſts generally agree, that their formation is owing to a greater than ordinary number of points of oſſification in the ſkull, or to the ordinary bones of the cranium not extending their oſſification far enough or ſoon enough; in which caſe, the unoſſified interſtice between ſuch bones begins a ſeparate oſſification in one or more points; from which the oſſification is extended to form as many diſtinct bones as there were points, that are indented into the large ordinary bones, and into each other.—Probably thoſe children who have a large opening in this place at their birth, will have the largeſt oſſa triquetra.—To confirm this account of the formation of theſe little bones, we may remark, that ſuch bones are ſometimes ſeen in other ſutures, as well as in the lambdoidal (*p*); and

(*l*) Lambdoides harmonialis, lambdoides inferior, occipitis corona.

(*m*) Euſtach. Oſſium examen.——Bauhin. Theat. Anat. lib. 3. cap. 5.——Paaw in Hippocrat. de vulner. capit. p. 50.

(*n*) Muſæum, lib. 3. c. 26.

(*o*) Bauhin. et Paaw ibid.—Bartholin. Anat. reform. lib. 4. c. 5.—Hildan. Epiſtol. 65.

(*p*) See examples in Veſal. lib. 1. cap. 6. fig. 4.—Paaw in Hippocrat. de cap. vuln.—Bartholin. Hiſt. Anat. cent. 1. Hiſt. 51.—Ruyſch. Muſ. Anat.—Sue Trad. d'oſteolog. p. 47.

and they are ſometimes in one table of the ſkull, and not in the other (*q*).

The *ſagittal* future (*r*) is placed longitudinally in the middle of the upper part of the ſkull, and commonly terminates at the middle of the coronal and of the lambdoidal ſutures; between which it is ſaid to be placed, as an arrow is between the ſtring and the bow.—However, this ſuture is frequently continued through the middle of the os frontis down to the root of the noſe; which, ſome (*s*) ſay, happens oftener in women than men; but others (*t*) allege, that it is to be met with more frequently in male ſkulls than in female: Among the ſkulls which I have ſeen thus divided, the female are the moſt numerous.——Several (*u*) have delineated and deſcribed the ſagittal ſuture, ſometimes dividing the occipital bone, as far down as the great hole through which the medulla ſpinalis paſſes. This I never ſaw.

In ſome old ſculls that are in my poſſeſſion, there is ſcarce a veſtige of any of the three ſutures which I have now deſcribed. In other heads, one or two of the ſutures only diſappear; but I never could diſcover any reaſon for thinking them diſpoſed in ſuch different manners in ſkulls of different ſhapes, as ſome ancients allege they are (*x*).

The *ſquamous agglutinations*, or *falſe ſutures* (*y*), are one on

(*q*) Hunald in Mem. de l'acad. des ſciences, 1730.

(*r*) Ῥαβδοειδης, οξελαια, επιζευγνυσα, Inſtar virgae, nervalis, inſtar teli, inſtar veru, ſecundum capitis longitudinem prorepens, conjungens, columnalis, recta, arcualis.

(*s*) Riolan. Comment. de oſſib. cap. 8.

(*t*) Veſal. lib. 1. cap. 6. et in epitome.

(*u*) Veſal. lib. 1. cap. 5. fig. 3. 4. et in text. cap. 6.———Paaw in Celſ. de re medic. cap. 1.———Laurent. Hiſt. Anat. lib. 2. cap. 16.

(*x*) Hippocrat. de vulner. capitis, § 1.———Galen. de oſſib. et de uſu part. lib. 9. cap. 17.

(*y*) Λεπιδοειδη, προσκολληματα, κροταφιαι. Temporales, corticales, mendoſae; harmonialis, commiſſurae in unguem.

on each ſide, a little above the ear, of a ſemicircular figure, formed by the overlapping (like one ſcale upon another) of the upper part of the temporal bones on the lower part of the parietal, where, in both bones, there are a great many ſmall riſings and furrows, which are indented into each other; though theſe inequalities do not appear till the bones are ſeparated. In ſome ſkulls, indeed, the indentations here are as conſpicuous externally as in other ſutures (z); and what is commonly called the poſterior part of this ſquamous ſuture, always has the evident ſerrated form; and therefore is reckoned by ſome (*a*) a diſtinct ſuture, under the name of *additamentum poſterius ſuturae ſquamoſae.*—I have ſeen two ſquamous ſutures on the ſame temple, with a ſemicircular piece of bone between them (*b*).

We ought here to remark, that the true ſquamous ſort of ſuture is not confined to the conjunction of the temporal and parietal bones, but is made uſe of to join all the edges of the bones on which each temporal muſcle is placed (*c*): For the two parts of the ſphenoidal ſuture which are continued from the anterior end of the common ſquamous ſuture juſt now deſcribed, of which one runs perpendicularly downwards, and the other horizontally forwards, and alſo the lower part of the coronal ſuture already taken notice of, may all be juſtly ſaid to pertain to the ſquamous ſuture.——The manner how, I imagine, this ſort of ſuture is formed at theſe places is, that, by the action of the ſtrong temporal muſcles on one ſide, and by the preſſure of the brain on the other, the bones are made ſo

(z) Columb. de re anat. lib. 1. cap. 4.——Dionis, Anat. 3. demonſtrat. des os.

(*a*) Albin. de oſſib. § 54.

(*b*) Sue Trad. d'oſteol. p. 48.

(*c*) Veſal. Anat. lib. 1. cap. 6.--Winſlow, Mem. de l'acad. des ſciences, 1720.

ſo thin, that their edges oppoſed to each other are not ſufficiently thick to ſtop the extenſion of their fibres in length, and thus to cauſe the common ſerrated appearance of ſutures explained in p. 36.; but the narrow edge of the one bone ſlides over the other. The ſquamous form is alſo more convenient here; becauſe ſuch thin edges of bones, when accurately applied one to another, have ſcarce any rough ſurface, to obſtruct or hurt the muſcle in its contraction; which is ſtill further provided for, by the manner of laying theſe edges on each other; for, in viewing their outſide, we ſee the temporal bones covering the ſphenoidal and parietal, and this laſt ſupporting the ſphenoidal, while both mount on the frontal: From which diſpoſition it is evident, that, while the temporal muſcle is contracting, which is the only time it preſſes ſtrongly in its motion on the bones, its fibres ſlide eaſily over the external edges. Another advantage of this ſtructure is, that the whole part is made ſtronger by the bones thus ſupporting each other.

The bones of the ſkull are joined to thoſe of the face, by ſchindyleſis and ſutures.——The ſchindyleſis is in the partition of the noſe.——The ſutures ſaid to be common to the cranium and face are five; viz. the ethmoidal, ſphenoidal, tranſverſe, and two zygomatic.———Parts, however, of theſe ſutures are only at the junction of the bones of the ſkull.

The *ethmoidal* and *ſphenoidal* ſutures ſurround the bones of theſe names; and in ſome places help to make up other ſutures, particularly the *ſquamous* and *tranſverſe*; and in other parts there is but one ſuture common to theſe two bones.

The *tranſverſe* ſuture is extended quite croſs the face, from the external canthus of one orbit to the ſame place of the other, by ſinking from the canthus down the outſide of the

the orbit to its bottom; then mounting upon its inside, it is continued by the root of the nose down the internal part of the other orbit, and rises up again on its outside to the other canthus. It may be here remarked, that there are some interruptions of this suture in the course I have described; for the bones are not every where contiguous, but are separated, to leave holes and apertures, as shall be mentioned hereafter.

The *zygomatic* sutures are one on each side, being short and slanting from above obliquely downwards and backwards, to join a process of the cheek bone to one of the temporal bones, which advance towards the face; so that the two processes thus united, form a sort of bridge or jugum, under which the temporal muscle passes, on which account the processes, and suture joining them, have been called *zygomatic*.

It must be observed, that the indentations of the sutures are not so strongly marked on the inside as on the outside of the cranium; and sometimes the bones seem to be joined by a straight line: Nay, in some skulls, the internal surface is found entire, while the sutures are manifest without; which may possibly be owing to the less extent of the concave than of the convex surface of the cranium, whereby the fibres of the internal side would be stretched farther out at the edges of the bones than the exterior ones, if they were not resisted. The resistances are, the fibres of the opposite bone, the parts within the skull, and the diploë: the last of which being the weakest, the most advanced fibres or serrae run into it, and leave the contiguous edges equal, and more ready to unite; whereas the serrae of the external table have space enough for their admission between the fibres of the opposite bone, and therefore remain of the indented form, and are less liable to the concretion whereby the sutures are obliterated (*d*).——By this mechanism,

(*d*) Hunald. Memoires de l'acad des sciences, 1730.

mechaniſm, there is no riſk of the ſharp points of the bones growing inwards, ſince the external ſerræ of each of the conjoined bones reſt upon the internal ſmooth-edged table of the other; and external forces applied to theſe parts are ſtrongly reſiſted, becauſe the ſutures cannot yield, unleſs the ſerrated edges of the one bone, and the plain internal plate of the other, are broken (*e*).

The advantages of the ſutures of the cranium are theſe: 1. That this capſula is more eaſily formed and extended into a ſpherical figure, than if it had been one continued bone. 2. That the bones which are at ſome diſtance from each other at birth, might then yield, and allow to the head a change of ſhape, accommodated to the paſſage it is engaged in. Whence, in difficult parturition, the bones of the cranium, inſtead of being only brought into contact, are ſometimes made to mount one upon the other. 3. It is alledged, that, through the ſutures there is a tranſpiration of ſteams from the brain, which was the old doctrine; or ſome communication of the veſſels without, and of thoſe within the ſkull, larger here than in any other part of the cranium, according to ſome moderns; and therefore cucuphae, fomentations, cataplaſms, cephalic plaſters, bliſters, are applied, and iſſues are eroded or cut in the head, at thoſe places where the ſutures are longeſt in forming, and where the connection of the bones is afterwards looſeſt, for the cure of a phrenitis, mania, inveterate headach, epilepſy, apoplexy, and other diſeaſes of the head. The favourers of the doctrine of tranſpiration, or communication of veſſels at the ſutures, endeavour to ſupport it by obſervations of perſons ſubject to headachs which cauſed death, from the ſutures being too cloſely united (*f*). 4. That the dura mater may be more firmly ſuſpended by its proceſſes, which inſinuate themſelves into this

(*e*) Winſlow, Memoires de l'acad. des ſciences, 1720.

(*f*) Columb. de re. anat. lib. 1. cap. 5.—Verduc. Nouvelle Oſteologie, chap. 14.—Dionis, Anat. 3. demonſt. des os.

this conjunction of the bones: for doing this equally, and where the greateſt neceſſity of adheſion is, the ſutures are diſpoſed at nearly equal diſtances; and the larger reſervoirs of blood, the ſinuſes, are under or near them. 5. That fractures might be prevented from reaching ſo far as they would in a continued bony ſubſtance. 6. That the connection at the ſutures being capable of yielding, the bones might be allowed to ſeparate; which has given great relief to patients from the violent ſymptoms which they had before this ſeparation happened (*g*). And it ſeems reaſonable to believe, that the opening of the ſutures was of great benefit to ſeveral others who were rather thought to have been hurt by it (*h*): for the conſequences of ſuch a force acting upon the brain, as was capable of thruſting the bones aſunder, muſt have been fatal, unleſs it had been thus yielded to.

Having gone through the general ſtructure of the cranium, I now proceed to examine each bone of which that brain caſe conſiſts, in the order in which I firſt named them.

OS FRONTIS.

THE os frontis (*i*) has its name from its being the only bone of that part of the face we call the *forehead*, though it reaches a good deal farther. It has ſome reſemblance in ſhape to the ſhell of the concha bivalvis, commonly called the *cockle*: for the greateſt part of it is convex externally, and concave internally, with a ſerrated circular edge; while the

(*g*) Ephemerid. Germanic. dec. 1. ann. 4. et 5. obſerv. 33.

(*h*) Ephemerid. Germ. dec. 2. ann. 9. obſ. 230. Ibid. cent. 10. obſ. 31.—Vander Linden Medicin. phyſ. cap. 8. art. 4. § 16.—Hildan. Obſerv. cent. 1. obſ. 1. cent. 2. obſ. 7.—Bauhin. Theat. Anat. lib. 3. cap. 6.—Pechlin. Obſerv. lib. 2. obſ. 39.

(*i*) Μετωπη, Βρεγμα, Coronale, inverecundum, puppis, ſenſus communis, ſincipitis.

the ſmaller part has proceſſes, and depreſſions, which make it of an irregular figure.

The external ſurface of the os frontis is ſmooth at its upper convex part; but ſeveral proceſſes and cavities are obſervable below: for at each angle of each orbit, the bone juts out to form four proceſſes, two internal, and as many external; which, from this ſituation, may well enough be named *angular*. Between the internal and external angular proceſſes on each ſide, an arched ridge is extended, on which the eye-brows are placed.——Very little above the internal end of each of theſe ſuperciliarly ridges a protuberance may be remarked, in moſt ſkulls, where there are large cavities, called *ſinuſes*, within the bone; of which hereafter.——Between the internal angular proceſſes a ſmall proceſs riſes, which forms a ſmall part of the noſe, and thence is named *naſal*.—Some obſerve a protuberant part on the edge of the bone behind each external angular proceſs, which they call *temporal* proceſſes; but theſe are inconſiderable.——From the under part of the ſuperciliary ridges, the frontal bone runs a great way backwards: theſe parts may juſtly enough be called *orbitar* proceſſes, which, contrary to the reſt of this bone, are concave externally, for receiving the globes of the eyes, with their muſcles, fat, &c.

In each of the orbitar proceſſes, behind the middle of the ſuperciliary ridges, a conſiderable ſinuoſity is obſerved, where the glandula innominata Galeni, or lacrymalis, is lodged.——Behind each internal angular proceſs, a ſmall pit may be remarked, where the cartilaginous pully of the muſculus obliquus major of the eye is fixed.—Between the two orbitar proceſſes, there is a large diſcontinuation of the bone, in which the cribriform part of the os ethmoides is incaſed.——The frontal bone has frequently little caverns formed in it where it is joined to the ethmoid bone.—

Behind each external angular proceſs, the ſurface of the frontal bone is conſiderably depreſſed where part of the temporal muſcle is placed.

The *foramina*, or holes, obſervable on the external ſurface of the frontal bone, are three in each ſide.——One in each ſuperciliary ridge, a little removed from its middle towards the noſe; through which a twig of the ophthalmic branch of the fifth pair of nerves paſſes out of the orbit, with a ſmall artery from the internal carotid, to be diſtributed to the teguments and muſcles of the forehead. —Theſe veſſels, in ſome ſkulls, make furrows in the os frontis, eſpecially in the bones of children, as has alſo been obſerved of another conſiderable veſſel of this bone near its middle (*k*); and therefore we ought to beware of tranſverſe inciſions on either ſide of the os frontis, which might either open theſe veſſels, or hurt the nerves, while they are yet in part within the bone: for when veſſels are thus wounded, it is difficult to ſtop the haemorrhagy, becauſe the adheſion of a part of the artery to the bone hinders its contraction, and conſequently ſtyptics can have little effect; the ſides of the furrow keep off compreſſing ſubſtances from the artery; and we would wiſh to ſhun cauteries or eſcharotics, becauſe they make the bone carious; and nerves, when thus hurt, ſometimes produce violent ſymptoms.——But to return to the ſuperciliary foramina, we muſt remark, that often, inſtead of a hole, a notch only is to be ſeen: nay, in ſome ſkulls, ſcarce a veſtige even of this is left; in others, both hole and notch are obſervable, when the nerve and artery run ſeparately. Frequently a hole is found on the one ſide, and a notch on the other; at other times we ſee two holes, or there is a common hole without, and two diſtinct entries internally. The reaſon of this variety of a hole, notch, depreſſion, or ſmoothneſs, in

(*k*) Ruyſch. Muſ. Anat. theca D, repoſit. 4. N° 3.

in the ſuperciliary ridge, is the different length and tenſion of the nerves and veſſels; the ſhorter they are, the more they are ſunk into the bone as it grows.—Near the middle of the inſide of each orbit, hard by or in the tranſverſe ſuture, there is a ſmall hole for the paſſage of the naſal twig of the firſt branch of the fifth pair of nerves, and of a branch of the ophthalmic artery. This hole is ſometimes entirely formed in the os frontis; in other ſkulls, the ſides of it are compoſed of this laſt bone and of the os planum. It is commonly known by the name of *orbitarium internum*, though *anterius* ſhould be added, becauſe of the next, which is commonly omitted.—This, which may be called *orbitarium internum poſterius*, is ſuch another as the former; only ſmaller, and about an inch deeper in the orbit: Through it a ſmall branch of the ocular artery paſſes to the noſe.—Beſides theſe ſix, there are a great number of ſmall holes obſervable on the outer ſurface of this bone, particularly in the two protuberances above the eye brows. Moſt of them penetrate no further than the ſinuſes, or than the diploë if the ſinuſes are wanting; though ſometimes I have ſeen this bone ſo perforated by a vaſt number of theſe ſmall holes, that, placed between the eye and a clear light, it appeared like a ſieve.—In the orbit of the generality of ſkeletons, we may obſerve one, two, or more holes, which allow a paſſage to a hog's briſtle through the ſkull. The place, ſize, and number of theſe, are however uncertain: they generally ſerve for the tranſmiſſion of ſmall arteries or nerves.

The internal ſurface of the os frontis is concave, except at the orbitar proceſſes, which are convex, to ſupport the anterior lobes of the brain. This ſurface is not ſo ſmooth as the external; for the larger branches of the arteries of the dura mater make ſome furrows in its ſides and back parts. The ſinuoſities from the luxuriant riſings of the brain, mentioned

tioned when describing the general structure of the cranium, are often very observable on its upper part; and its lower and fore parts are marked with the contortions of the anterior lobes of the brain.—Through the middle of this internal surface, where always in children, and sometimes in old people, the bone is divided, either a ridge stands out, to which the upper edge of the falx is fastened, or a furrow runs, in which the upper side of the superior longitudinal sinus is lodged; on both these accounts, chirurgical authors justly forbid the application of the trepan here.—The reason of this difference in skulls is alleged by some authors to be this, That in thin skulls the ridge strengthens the bones, and in thick ones there is no occasion for it. To this way of accounting for the phenomenon, it may justly be objected, that, generally, very thick skulls have a large spine here, and frequently thin ones have only a furrow. Perhaps this variety may be owing to the different times of complete ossification of those parts in different subjects; for if the two sides of this bone meet before they arrive at their utmost extent of growth, they unite very firmly, and all their fibres endeavour to stretch themselves out where there is the least resistance, that is, between the hemispheres of the brain. To support this reasoning, we may remark, that those adults whose frontal bone is divided by the sagittal future, never have a ridge in this place.

Immediately at the root of this ridge or furrow, there is a small hole, which sometimes pierces through the first table, and, in other skulls, opens into the superior sinus of the ethmoid bone within the nose. In it a little process of the falx is lodged; and a small artery, and sometimes a vein, runs (*l*); and the superior longitudinal sinus begins here.—This hole, however, is often not entirely proper to the os frontis: for, in several skulls, the lower part is formed in the upper part of

(*l*) Morgagn. Adversar. 6. animad. 32.

of the baſe of the criſta galli, which is a proceſs of the ethmoid bone (*m*).

The os frontis is compoſed of two tables, and an intermediate diploë, as the other bones of the cranium are: It is of a mean thickneſs between the os occipitis and the parietal bones; and is nearly equally denſe throughout, except at the orbitar proceſſes, where, by the action of the eye on one ſide, and preſſure of the lobes of the brain on the other, it is made extremely thin and diaphanous, and the meditullium is entirely obliterated. In this place there is ſo weak a defence for the brain, that fencers eſteem a puſh in the eye mortal (*n*).

The diploë is alſo exhauſted in that part above the eyebrows, where the two tables of the bone ſeparate, by the external being protruded outwards, to form two large cavities, called *ſinus frontales*.—Theſe are divided by a middle perpendicular bony partition.—Their capacities in the ſame ſubject are ſeldom equal; in ſome the right, in others the left, is largeſt.—And in different bones their ſize is as inconſtant: nay, I have examined ſome where they were entirely wanting; which oftener happens in ſuch as have a flat forehead, and whoſe ſagittal ſuture is continued down to the noſe, than in others (*o*).—In ſome ſkulls, beſides the large perpendicular *ſeptum*, ſeveral bony pillars, or ſhort partitions, are found in each ſinus: in others theſe are wanting.—For the moſt part the *ſeptum* is entire; at other times it is diſcontinued, and the two *ſinuſes* communicate.—When the ſinuſes are ſeen in ſuch ſkulls as have the frontal bone divided by the ſagittal ſuture, the partition dividing theſe cavities is evidently compoſed of two plates, which eaſily ſeparate.—Each ſinus commonly opens by a roundiſh ſmall hole, at the inner

(*m*) Ingraff. Comment. in Galen. de oſſib. cap. 1. comment. 8.

(*n*) Ruyſch. Obſerv. Anat.-chir. obſerv. 54.—Diemerbroek, Anat. lib. 3. cap. 10.—Bonet. Sepulch. Anat. lib. 4. § 3. obſerv. 17.

(*o*) Fallop. Expoſit. de oſſib. cap. 13.

inner and lower part of the internal angular proceſſes, into a ſinus formed in the noſe, at the upper and back part of the os unguis; near to which there are alſo ſome other ſmall ſinuſes of this bone (*p*), the greater part of which open ſeparately near the ſeptum narium, and often terminate in the ſame common canal with the large ones.

In a natural and ſound ſtate, theſe cavities are of conſiderable advantage: for the organ of ſmelling being thus enlarged, the effluvia of odorous bodies more difficultly eſcape it; and their impreſſions being more numerous, are therefore ſtronger, and affect the organ more. That odorous particles may be applied to the membrane of the ſinuſes, is evident from the pain felt in this part of the forehead, when the effluvia of volative ſpirits, or of ſtrong aromatics, are drawn up into the noſe by a quick inſpiration.—Theſe, and the other cavities which open into the noſe, increaſe the ſound of our voice, and render it more melodious, by ſerving as ſo many vaults to reſound the notes. Hence people labouring under a coryza, or ſtoppage of the noſe from any other cauſe, when they are by the vulgar, though falſely, ſaid to ſpeak through the noſe, have ſuch a diſagreeable harſh voice.—The liquor ſeparated in the membrane of theſe ſinuſes runs down upon the membrane of the noſe to keep it moiſt.

From the deſcription of theſe ſinuſes, it is evident, how uſeleſs, nay, how pernicious, it muſt be to apply a trepan on this part of the ſkull: for this inſtrument, inſtead of piercing into the cavity of the cranium, would reach no further than the ſinuſes; or, if the inner table was perforated, any extravaſated blood that happened to be within the ſkull, would not be diſcharged outwardly, but would fall into the ſinuſes, and there ſtagnate, corrupt, and ſtimulate

(*p*) Cowper in Drake's Anthrolopog. book 3. chap. 10.

ſtimulate the ſenſible membranes; from which alſo there would be ſuch a conſtant flow of glairy mucus, as would retard, if not hinder a cure, and would make the ſore degenerate into an incurable fiſtula. Beſides, as it would be almoſt impoſſible in this caſe to prevent the air, paſſing through the noſe, from having conſtant acceſs to the dura mater or brain; ſuch a corruption would be brought on theſe parts as would be attended with great danger. Farther, in reſpiration, the air ruſhing violently into theſe cavities of the os frontis, and paſſing through the external orifice, whenever it was not well covered and defended, would not only prevent the cloſing up of the external orifice, but might otherwiſe bring on bad conſequences (*q*).—The membrane lining theſe ſinuſes is ſo ſenſible, that inflammations of it muſt create violent torture (*r*); and worms, or other inſects crawling there, muſt give great uneaſineſs (*s*).

The upper circular part of the os frontis is joined to the oſſa parietalia, from one temple to the other, by the coronal ſuture. From the termination of the coronal ſuture to the external angular proceſſes, this bone is connected to the ſphenoid by the ſphenoidal ſuture. At the external canthi of the eyes, its angular proceſſes are joined by the tranſverſe ſuture to the oſſa malarum, to which it adheres one-third down the outſide of the orbits; whence to the bottom of theſe cavities, and a little upon their internal ſides, theſe orbitar proceſſes are connected to the ſphenoidal bone by that ſame ſuture.——In ſome few ſkulls, however, a diſcontinuation of theſe two bones appears at the upper part of the long ſlit, near the bottom of the orbit.—On the inſide of each orbit, the orbitar proceſs is indented

(*q*) Paaw de oſſibus, pars 1. cap. 7.——Palfyne Anatom. chir. traite 4. chap. 15. Nouvelle Oſteologie, partie 2. chap. 3.

(*r*) Fernel Pathalog. lib. 5. cap. 7.——Saltzman Decur. obſerv. 10.

(*s*) Fernel Patholog. lib. 5. cap. 7.——Bartholin. Epiſtol. Medic. cent. 2. epiſt. 74.—Hiſt. de l'acad. des ſciences, 1708 &. 1733.

dented between the cribriform part of the ethmoid bone and the os planum and unguis.—The tranſverſe ſuture afterwards joins the frontal bone to the ſuperior naſal proceſſes of the oſſa maxillaria ſuperiora, and to the naſal bones. And, laſtly, its naſal proceſs is connected to the naſal lamella of the ethmoid bone.

The frontal bone ſerves to defend and ſupport the anterior lobes of the brain. It forms a conſiderable part of the cavities that contain the globes of the eyes, helps to make up the ſeptum narium, organ of ſmelling, &c. From the deſcription of the ſeveral parts, the other uſes of this bone are evident.

In a ripe child, the frontal bone is divided through the middle; the ſuperciliary holes are not formed; often a ſmall round piece of each orbitar proceſs, behind the ſuperciliary ridge, is not oſſified; and there is no ſinus to be ſeen within its ſubſtance.

OSSA PARIETALIA.

Each of the two Oſſa Parietalia (*t*), or bones ſerving as walls to the encephalon, is an irregular ſquare; its upper and fore ſides being longer than the one behind or below. The inferior ſide is a concave arch; the middle part receiving the upper round part of the temporal bone.——The angle formed by this upper ſide and the fore one, is ſo extended, as to have the appearance of a proceſs.

The external ſurface of each os parietale is convex. Upon it, ſomewhat below the middle height of the bone, there is a tranſverſe arched ridge, generally of a whiter colour than any other part of the bone; from which in bones that have ſtrong prints of muſcles, we ſee a great many converging furrows, like ſo many radii drawn from a circumference

(*t*) Κορυφης, Paria ſyncipitis, verticis, arcualia, nervalia, cogitationis, rationis, bregmatis, madefactiones.

rence towards a centre. From this ridge of each bone the temporal muſcle riſes: and, by the preſſure of its fibres, occaſions the furrows juſt now mentioned.—Below theſe we obſerve, near the ſemicircular edges, a great many riſings and depreſſions, which are joined to like inequalities on the inſide of the temporal bone, and form the ſquamous ſuture. The temporal bone may therefore ſerve here as a buttreſs, to prevent the lower ſide of the parietal from ſtarting outwards when its upper part is preſſed or ſtruck (*u*).

Near the upper ſides of theſe bones, towards the hind part, is a ſmall hole in each, through which a vein paſſes from the teguments of the head to the longitudinal ſinus. Sometimes I have ſeen a branch of the temporal artery paſs through this hole, to be diſtributed to the upper part of the falx, and to the dura mater at its ſides, where it had frequent anaſtomoſes with the branches of the arteries derived from the external carotids, which commonly have the name of the arteries of the dura mater, and with the branches of the internal carotids which go to the falx.—— In ſeveral ſkulls, one of the oſſa parietalia has not this hole: In others, there are two in one bone; and in ſome, not one in either. Moſt frequently this hole is through both tables; at other times the external table only is perforated. ——The knowledge of the courſe of theſe veſſels may be of uſe to ſurgeons when they make any inciſion near this part of the head; leſt, if the veſſels are raſhly cut near the hole, they ſhrink within the ſubſtance of the bone, and ſo cauſe an obſtinate hæmorrhagy, which neither ligatures nor medicines can ſtop.

On the inner concave ſurface of the parietal bones, we ſee a great many deep furrows, diſpoſed ſomewhat like the branches of trees: the furrows are largeſt and deepeſt at

(*u*) Hunauld. in Mem. de l'acad. des ſciences, 1730.

the lower edge of each os parietale, eſpecially near its anterior angle, where a complete canal is ſometimes formed. They afterwards divide into ſmall furrows, in their progreſs upwards.—In ſome ſkulls a large furrow begins at the hole near the uper edge, and divides into branches, which join with thoſe that come upwards; ſhewing the communications of the upper and lower veſſels of the dura mater.—In theſe furrows we frequently ſee paſſages into the diploë; and ſometimes I have obſerved canals going off, which allowed a ſmall probe to paſs a few inches into the bony ſubſtance. Some (*x*) authors tell us, that they have obſerved theſe canals piercing the bone towards the occiput.—On the inſide of the upper edge of the oſſa parietalia, there is a large ſinuoſity, frequently larger in the bone of one ſide than of the other, where the upper part of the falx is faſtened, and the ſuperior longitudinal ſinus is lodged.—Part of the lateral ſinuſes generally makes a depreſſion near the angle, formed by the lower and poſterior ſides of theſe bones; and the pits made by the prominent parts of the brain are to be ſeen in no part of the ſkull more frequent or more conſiderable, than in the internal ſurface of theſe bones.

The oſſa parietalia are the moſt equal and ſmooth, and are among the thinneſt bones of the cranium; but they enjoy the general ſtructures of two tables and diploë the compleſeſt.

Theſe bones are *joined* at their fore-ſide to the os frontis by the coronal ſuture; at their long inferior angles, to the ſphenoid bone, by part of the ſuture of this name; at their lower edge, to the oſſa temporum, by the ſquamous ſuture, and its poſterior additamentum; behind, to the os occipitis, or oſſa triquetra, by the lambdoid ſuture; and above, to one another, by the ſagittal ſuture.

They

(*x*) Cowper. Anatom. explic. of tab. 90. fig. 2.

They have no particular *uſes* beſides thoſe mentioned in the deſcription of their ſeveral parts, except what are included in the account of the general ſtructure of the cranium.

In a child born at the full time, none of the ſides of this bone are completed; and there never is a hole in the oſſified part of it near the ſagittal ſuture.

The large unoſſified ligamentous part of the cranium, obſervable between the parietal bones and the middle of the divided os frontis of new-born children, called by the vulgar the *open of the head*, was imagined by the ancients to ſerve for the evacuation of the ſuperfluous moiſture of the brain: and therefore they named it *bregma* (*y*), or the fountain; ſometimes adding the epithet *pulſatilis*, or beating, on account of the pulſation of the brain felt through this flexible ligamento-cartilaginous ſubſtance. Hence the parietal bones are very frequently called *oſſa bregmatis*.

The upper middle part of the head of a child, in a natural birth, being what preſents itſelf firſt at the os uteri (*z*), an accoucheur may reach the bregma with his finger, when the os uteri is a little opened. If the bregma is ſtretched, and the pulſation of the brain is felt through it, the child is certainly alive: but if it is ſhrivelled and flaccid, and without any obſervable pulſation in it, there is ſome reaſon to ſuſpect the child to be very weak, or dead. Thoſe who practiſe midwifery ſhould therefore examine the ſtate of the bregma accurately.

All the bregma is generally oſſified before ſeven years of age. Several authors (*a*) ſay, they have obſerved it unoſſified in adults; and phyſicians, who order the application of medicines at the meeting of the coronal and ſagittal ſutures, ſeems yet

(*y*) Pulpitans vertex, foliolum, folium, triangularis lacuna.

(*z*) Burton's Midwifery, § 51.—Smellie's Midwifery, book 1. chap. 1. § 5.

(*a*) Bartholin. Anat. reform. lib. 4. cap. 6.—Diemerboeck, Anat. lib. 9. cap. 6.—Kerkring Oſteogen. cap. 2.

yet to think that a derivation of noxious humours from the encephalon is more eaſily procured at this, than at any other part of the ſkull, and that medicines have a greater effect here than elſewere in the internal diſorders of the head.

OSSA TEMPORUM.

Ossa Temporum (*b*), ſo named, ſay authors, from the hair's firſt becoming grey on the temples, and thus diſcovering peoples ages, are each of them equal and ſmooth above, with a very thin ſemicircular edge; which, from the manner of its connection with the neighbouring bones, is diſtinguiſhed by the name of *os ſquamoſum*.——Behind this, the upper part of the temporal bone is thicker, and more unequal; and is ſometimes deſcribed as a diſtinct part, under the name of *pars mamillaris* (*c*).——Towards the baſe of the ſkull, the temporal bone appears very irregular and unequal; and this part, inſtead of being broad, and placed perpendicularly, as the others are, is contracted into an oblong very hard ſubſtance, extended horizontally forwards and inwards, which in its progreſs becomes ſmaller, and is commonly called *os petroſum*.

Three external proceſſes of each temporal bone are generally deſcribed.—The firſt, placed at the lower and hind part of the bone, from its reſemblance to a nipple, is called *maſtoides* or *mamillaris*. It is not ſolid; but within is compoſed of cancelli, or ſmall cells, which have a communication with the large cavity of the ear, the drum; and therefore ſounds, being multiplied in this vaulted labyrinth, are increaſed before they are applied to the immediate organ

(*b*) Κροταφων, κορσων, κορρων, λεπιδοειδη, πολυειδη, λιθοειδη, Temporalia, lapidoſa, mendoſa, dura, arcualia, tympanum, armalia, ſaxea, parietalia.

(*c*) Albin. de oſſib. ſect. 26.

gan of hearing. Into the maſtoid proceſs the ſterno maſtoideus muſcle is inſerted; and to its back part, where the ſurface is rough, the trachelo-maſtoideus and part of the ſplenius are fixed. About an inch farther forward, the ſecond proceſs begins to riſe out from the bone; and having its origin continued obliquely downwards and forwards for ſome way, it becomes ſmaller, and is ſtretched forwards to join with the os malae; they together forming the bony jugum, under which the temporal muſcle paſſes. Hence this proceſs has been named *zygomatic* (*d*). Its upper edge has the ſtrong aponeuroſis of the temporal muſcle fixed into it; and its lower part gives riſe to a ſhare of the maſſeter.—The fore-part of the baſe of this proceſs is an oblong tubercle, which, in a recent ſubject, is covered with a ſmooth poliſhed cartilage, continued from that which lines the cavity immediately behind this tubercle.——From under the craggy part of the os temporum, the third proceſs ſtands out obliquely forwards. The ſhape of it is generally ſaid to reſemble the ancient *ſtylus ſcriptorius*; and therefore it is called the *ſtyloid proceſs* (*e*) Some authors (*f*) however contend, that it ought to be named *ſteloid*, from its being more like a pillar. Several muſcles have their origin from this proceſs, and borrow one half of their name from it; as *ſtylo-gloſſus*, *ſtylo-hyoides*, *ſtylo-pharyngeus*: a ligament of the os hyoides is ſometimes fixed to it; and another is extended from it to the inſide of the angle of the lower jaw. This proceſs is often, even in adults, not entirely oſſified, but is ligamentous at its root, and is ſometimes compoſed of two or three diſtinct pieces.——Round the root of it, eſpecially at the forepart, there

(*d*) Paris, anſae oſſium temporum, oſſa arcualia, paria, jugalia, conjugalia.

(*e*) Γραφοειδη, βελονοειδη, πληκτρον, Os calaminum, ſagittale, clavale, acuale calcar capitis.

(*f*) Galen, de uſu part. lib. 2. cap. 4.—Fallop. Obſerv. anat.

there is a rising of the os petrosum, which some authors have esteemed a process: and, from the appearance it makes with the styliform, have named it *vaginalis*.—Others again have, under the name of *auditory* process, reckoned among the external processes, that semicircular ridge, which running between the root of the mastoid and zygomatic processes, forms the under part of the external *meatus auditorius*.

The *sinuosities* or depressions on the external surface of each temporal bone are these:——A long fossa at the inner and back part of the root of the mammary process, where the posterior head of the digastric muscle has its origin.——Immediately before the root of the zygomatic process, a considerable hollow is left for lodging the crotaphite muscle.——Between the zygomatic, auditory, and vaginal processes, a large cavity is formed; through the middle of which, from top to bottom, a fissure is observable, into which part of the ligament that secures the articulation of the lower jaw with this bone is fixed. The fore-part of the cavity being lined with the same cartilage which covers the tubercle before it, receives the condyle of the jaw; and in the back-part a small share of the parotid gland, and a cellular fatty substance, are lodged.—At the inside of the root of the styloid apophyse, there is a thimble-like cavity, where the beginning of the internal jugular vein, or end of the lateral sinus is lodged.—And as the sinuses of the two sides are frequently of unequal size, so one of these cavities is as often larger than the other (g).——Round the external meatus auditorius, several sinuosities are formed for receiving the cartilages and ligaments of the ear, and for their firm adhesion.

The *holes* that commonly appear on the outside of each of these bones, and are proper to each of them, are five.

——The

(g) Hunauld. in Mem. de l'acad. des sciences, 1730.

——The firſt, ſituated between the zygomatic and maſtoid proceſſes, is the orifice of a large funnel-like canal, which leads to the organ of hearing; and is therefore called *meatus auditorius externus* (*h*).—The ſecond gives paſſage to the portio dura of the ſeventh pair of nerves; and from its ſituation between the maſtoid and ſtyloid proceſſes, is called *foramen ſtylo-maſtoideum* (*i*).—Some way before, and to the inſide of the ſtyloid proceſs, is the third hole: the canal from which runs firſt upwards, then forwards, and receives into it the internal carotid artery, and the beginning of the intercoſtal nerve; where this canal is about to make the turn forwards, one, or ſometimes two, very ſmall holes go off towards the cavity of the ear, called *tympanum*: Through theſe Valſalva (*k*) affirms the proper artery or arteries of that cavity are ſent.—On the anterior edge of this bone, near the former, a fourth hole is obſervable, being the orifice of a canal which runs outwards and backwards, in a horizontal direction, till it terminates in the tympanum. This, in the recent ſubject, is continued forward and inward, from the parts which I mentioned juſt now as its orifice in the ſkeleton, to the ſide of the noſtrils; being partly cartilaginous, and partly ligamentous. The whole canal is named *Iter a palato ad aurem*, or *Euſtachian tube*.——On the external ſide of the bony part of this canal, and on the top of the chink, in the cavity that receives the condyle of the lower jaw, is the courſe of the little nerve, called *chorda tympani*, and commonly ſaid to be reflected from the lingual branch of the fifth pair, till it enters the tympanum, to run acroſs this cavity.——The fifth hole is very uncertain, appearing ſometimes behind the maſtoid proceſs; ſometimes it is common to the temporal and occipital

(*h*) Πυλης της ακοης, οτη των ωτων, feneſtra aurium.

(*i*) Aquaeductus Fallopii.

(*k*) De aure humana, cap. 2. § 22. et tab. 7. fig. 1.

occipital bones; and in ſeveral ſkulls there is no ſuch hole. The uſe of it, when found, is for the tranſmiſſion of a vein from the external teguments to the lateral ſinus: But, in ſome ſubjects, a branch of the occipital artery paſſes through this hole, to ſerve the back-part of the dura mater; in others, I have ſeen two or three ſuch holes: but they are oftener wanting than found. And we may once for all, in general remark, that the largeneſs, number, ſituation, and exiſtence of all ſuch holes as for the moſt part allow only a paſſage for veins from without to the internal receptacles, are very uncertain.

The internal ſurface of the oſſa temporum is unequal; the upper circular edge of the ſquamous part having numerous ſmall ridges and furrows for its conjunction with the parietal bones; and the reſt of it is irregularly marked with the convolutions of the middle part of the brain, and with furrows made by the branches of the arteries of the dura mater.

From the under part of this internal ſurface, a larger tranſverſe hard craggy protuberance runs horizontally inwards and forwards, with a ſharp edge above, and two flat ſides, one facing obliquely forwards and outwards, and the other as much backwards and inwards. To the ridge between theſe two ſides, the large lateral proceſs of the dura mater is fixed.

Sometimes a ſmall bone, like the ſeſamoid, is found between the ſmall end of this petrous proceſs and the ſphenoid bone (*l*).

Towards the back-part of the inſide of the temporal bone, a large deep foſſa is conſpicuous, where the lateral ſinus lies; and frequently on the top of the petrous ridge, a furrow may be obſerved, where a ſmall ſinus is ſituated.

The

(*l*) Riolan. Comment. de oſſib. cap. 32.——Winſlow, Expoſition anatomique de corps humain. trait. des os ſecs, § 266.

The internal proper *foramina* of each of theſe bones are, firſt, the internal meatus auditorius in the poſterior plain ſide of the petrous proceſs. This hole ſoon divides into two; one of which is the beginning of the aqueduct of Fallopius; the other ends in ſeveral very ſmall canals (*m*) that allow a paſſage to the branches of the portio mollis of the ſeventh pair of nerves, into the veſtibule and cochlea. Through it alſo an artery is ſent, to be diſtributed to the organ of hearing.—The ſecond hole, which is on the anterior plain ſide of the craggy proceſs, gives paſſage to a reflected branch of the ſecond branch of the fifth pair of nerves, which joins the portio dura of the auditory nerve, while it is in the aqueduct (*n*), ſmall branches of blood-veſſels accompanying the nerves, or paſſing through ſmaller holes near it.—The paſſage of the cutaneous vein into the lateral ſinus, or of a branch of the occipital artery, is ſeen about the middle of the large foſſa for that ſinus; and the orifice of the canal of the carotid artery is evident at the under part of the point of the petrous proceſs.

Beſides theſe proper holes of the temporal bones, which appear on their external and internal ſurfaces, there are two others in each ſide that are common to this bone, and to the occipital and ſphenoidal bones; which ſhall be mentioned afterwards in the deſcription of theſe bones.

The upper round part of the ſquamous bones is thin, but equal; while the low petrous part is thick and ſtrong, but irregular and unequal, having the diſtinction of tables and diploë confounded, with ſeveral cavities, proceſſes, and bones within its ſubſtance, which are parts of the organ of hearing. That a clear idea may be had of this beautiful, but intricate organ, anatomiſts generally chooſe to demonſtrate all its parts together. I think the method good; and

(*m*) Valſalva de aure humana, cap. 3. § 11.

(*n*) Ibid. cap. 3. § 10.

therefore, ſince it would be improper to inſert a complete treatiſe on the ear here, I ſhall omit the deſcription of the parts contained within the os petroſum of the ſkeleton.

The temporal bones are joined above to the parietal bones by the ſquamous ſutures and their poſterior additamenta: Before, to the ſphenoid bone, by the ſuture of that name; to the cheek bones, by the zygomatic ſutures: Behind, to the occipital bone by the lambdoid ſuture and its additaments; and they are articulated with the lower jaw in the manner which ſhall be deſcribed when this bone is examined.

The *purpoſes* which theſe two bones ſerve, are eaſily collected, from the general uſe of the cranium, and from what has been ſaid in the deſcription of their ſeveral parts.

In an infant, a ſmall fiſſure is to be obſerved between the thin upper part and the lower craggy part of each of theſe bones; which points out the recent union of theſe parts.——Neither maſtoid nor ſtyloid proceſſes are yet to be ſeen.——Inſtead of a bony funnel-like external meatus auditorius, there is only a ſmooth bony ring, within which the membrane of the drum is faſtened.——At the entry of the Euſtachian tube, the ſide of the tympanum is not completed.——A little more outward than the internal auditory canal, there is a deep pit, over the upper part of whoſe orifice the interior ſemicircular canal of the ear is ſtretched, and ſome way below this, the poſterior ſemicircular canal alſo manifeſtly appears.

OS OCCIPITIS.

Os occipitis (*o*), ſo called from its ſituation, is convex on the outſide, and concave internally. Its figure is an irregular ſquare, or rather rhomboid; of which the angle above

(*o*) Inion, Baſilare, prorae, memoriae, pyxidis, fibroſum, nervoſum, lambdoe.

above is generally a little rounded; the two lateral angles are more finiſhed, but obtuſe; and the lower one is ſtretched forward in form of a wedge, and thence is called by ſome the *cuneiform* proceſs.—If one would, however, be very nice in obſerving the ſeveral turns which the edges of the os occipitis make, five or ſeven ſides, and as many angles of this bone, might be deſcribed.

The external ſurface is convex, except at the cuneiform apophyſe, where it is flatted. At the baſe of this triangular proceſs, on each ſide of the great hole, but more advanced forwards than the middle of it, the large oblong protuberances named the *condyles*, appear, to ſerve for the articulation of this bone with the firſt vertebra of the neck. The ſmooth ſurface of each of theſe condyloid proceſſes is longeſt from behind forwards, where, by their oblique ſituation, they come much nearer to each other than they are at their back-part. Their inner ſides are lower than the external, by which they are prevented from ſliding to either ſide out of the cavities of the firſt vertebra (*p*). In ſome ſubjects, each of theſe plain ſmooth ſurfaces ſeems to be divided by a ſmall riſing in its middle; and the lower edge of each condyle, next the great foramen, is diſcontinued about the middle, by an intervening notch: Whence ſome (*q*) allege, that each of theſe apophyſes is made up of two protuberances. Round their root a ſmall depreſſion and ſpongy roughneſs is obſervable, where the ligaments for ſurrounding and ſecuring their articulations adhere.—Though the motion of the head is performed on the condyles, yet the centre of gravity of that globe does not fall between them, but is a good way farther forward; from which mechaniſm it is evident, that the muſcles which pull the head back muſt be in a conſtant ſtate of contraction;

(*p*) Galen. de uſu part. lib. 12. cap. 7.
(*q*) Diemerbroek, Anat. lib. 9. cap. 6.

contraction; which is ſtronger than the natural contraction of the proper flexors, elſe the head would always fall forwards, as it does when a man is aſleep, or labours under a palſy, as well as in infants, where the weight of the head far exceeds the proportional ſtrength of theſe muſcles. This ſeeming diſadvantageous ſituation of the condyles is, however, of ſingular uſe, by allowing ſufficient ſpace for the cavities of the mouth and fauces, and for lodging a ſufficient number of muſcles, which commonly ſerve for other uſes; but may at pleaſure be directed to act on the head, and then they have an advantageous lever to act with, ſo as to be able to ſuſtain a conſiderable weight, or other force applied, to pull the head back.

Somewhat more externally than the condyles, there is a ſmall riſing and ſemilunated hollow in each ſide, which make part of the holes common to the occipital and petrous bones.——Immediately behind this, on each ſide, a ſcabrous ridge is extended from the middle of the condyle towards the root of the maſtoid proceſs. Into this ridge the muſculus lateralis, commonly aſcribed to Fallopius, is inſerted.——About the middle of the external convex ſurface, a large arch runs croſs the bone; from the upper lateral parts of which the occipital muſcles have their riſe; to its middle the trapezii are attached; and half-way between this and the great hole, a leſſer arch is extended.——In the hollows between the middle of theſe arches the complexi are inſerted; and in the depreſſions more external and further forward than theſe, the ſplenii are inſerted.——Between the middle of the leſſer arch and the great hole, the little hollow marks of the recti minores appear; and on each ſide of theſe the fleſhy inſertions of the obliqui ſuperiores and recti majores make depreſſions.——Through the middle of the two arches a ſmall ſharp ſpine is placed, which ſerves as a kind of partition between the muſcles

of

of different ſides, or rather is owing to the action of the muſcles depreſſing the bone on each ſide of it, while this part is free from their compreſſion. Theſe prints of the muſcles of this bone are very ſtrong and plain in ſome ſubjects, but are not ſo diſtinct in others. All round the great foramen, the edges are unequal, for the firmer adheſion of the ſtrong circular ligament which goes thence to the firſt vertebra. One end of each lateral or moderator ligament of the head, is fixed to a rough ſurface at the fore-part of each condyle, and the perpendicular one is connected to a rough part of the edge of the great hole between the two condyles. Immediately before the condyles, two little depreſſions are made in the external ſurface of the cuneiform proceſs, for the inſertion of the recti anteriores minores muſcles, which are unjuſtly aſcribed to Cowper: And ſtill farther forward, near the ſphenoid bone, are two other ſuch depreſſions, for the reception of the recti anteriores majores. When we conſider the ſize of the prints of muſcles on the occipital bone, before and behind its condyles, and at the ſame time compare their diſtances from theſe centres of motion of the head, we muſt ſee how much ſtronger the muſcles are which pull the head backwards, than thoſe which bend it forward; and how much greater force the former acquire by the long lever they act with, than the latter which are inſerted ſo near the condyles. This great force in the extenſor muſcles is altogether neceſſary, that they might not only keep the head from falling forward in an erect poſture, but that they might ſupport it when we bow forward in the moſt neceſſary offices of ſocial life, when the weight of the head comes to act at right angles on the vertebra of the neck, and obtains a long lever to act with.

On the inner ſurface of the os occipitis we ſee two *ridges*; one ſtanding perpendicularly, the other running horizontally acroſs

acroſs the firſt. The upper part of the perpendicular limb of the croſs, to which the falx is fixed, is hollowed in the middle, or often on one ſide, for the reception of the ſuperior longitudinal ſinus; and the lower part of it has the ſmall or third proceſs of the dura mater faſtened to it, and is ſometimes hollowed by the occipital ſinus. Each ſide of the horizontal limb is made hollow by the lateral ſinuſes incloſed in the tranſverſe proceſs of the dura mater; the foſſa in the right ſide being generally a continuation of the one made by the longitudinal ſinus in the perpendicular limb, and therefore is larger than the left one (*r*).—Round the middle of the croſs there are four large depreſſions, ſeparated by its limbs; the two upper ones being formed by the back part of the brain, and the two lower ones by the cerebellum.—Farther forward than the laſt mentioned depreſſions, is the lower part of the foſſa for the lateral ſinus on each ſide.—The inner ſurface of the cuneiform apophyſe is made concave for the reception of the medulla oblongata, and of the baſilar artery.—A furrow is made on each ſide, near the edges of this proceſs, by a ſinus of the dura mater, which empties itſelf into the lateral ſinus (*s*).

The *holes* of this bone are commonly five proper, and two common to it and to the temporal bones.——The firſt of the proper holes, called *foramen magnum* (*t*) from its ſize, is immediately behind the wedge-like proceſs; and allows a paſſage to the medulla oblongata, nervi acceſſorii, to the vertebral arteries, and ſometimes to the vertebral veins.—At each ſide of this great hole, near its fore part, and immediately above the condyles, we always find a hole, ſometimes two, which ſoon unite again into one, that opens externally; through

(*r*) Morgagn. Adverſ. anat. 6. animad. 1.

(*s*) Albin. de oſſib. ſect. 65.

(*t*) Rachitidis, Medullæ ſpinalis.

through theſe the ninth pair of nerves go out of the ſkull. —The fourth and fifth holes pierce from behind the condyle of each ſide into the foſſæ of the lateral ſinuſes; they ſerve for the paſſage of the cervical veins to theſe ſinuſes. Often one of theſe holes is wanting, ſometimes both, when the veins paſs through the great foramen.—Beſides theſe five, we frequently meet with other holes near the edges of this bone, for the tranſmiſſion of veins; but their number and diameter are very uncertain. The two common foramina are the large irregular holes, one in each ſide, between the ſides of the cuneiform proceſs and the edges of the petrous bones. In a recent ſubject, a ſtrong membrane runs croſs from one ſide to the other of each of theſe holes: In ſome heads I have ſeen this membrane oſſified, or a bony partition dividing each hole: And in the greater number of adult ſkulls, a ſmall ſharp-pointed proceſs ſtands out from the os petroſum, and there is a more obtruſe riſing in the occipital bone, between which the partition is ſtretched. Behind this partition, where the largeſt ſpace is left, the lateral ſinus has its paſſage; and before it the eighth pair of nerves and acceſſorius make their exit out of the ſkull; and ſome authors ſay, an artery paſſes through this hole to be beſtowed on the dura mater.

The *occipital* bone is among the thickeſt of the cranium, though unequally ſo; for it is ſtronger above, where it has no other defence than the common teguments, than it is below; for being there preſſed by the lobes of the brain and cerebellum on one ſide, and by the action of the muſcles on the other, it is ſo very thin as to be diaphanous in many ſkulls: But then theſe muſcles ward of injuries; and the ridges and ſpines, which are frequent here, make it ſufficiently ſtrong to reſiſt ordinary forces.

The tables and diploë are tolerably diſtinct in this bone, except where it is ſo thin as to become diaphanous.

The

The occipital bone is *joined* above to the offa parietalia, and to the triquetra, when preſent, by the lambdoid ſuture; —laterally to the temporal bones, by the additamenta of the lambdoid ſuture—below to the ſphenoid bone, by the end of its cuneiform proceſs, in the ſame way that epiphyſes and their bones are joined: For in children a ligamentous cartilage is interpoſed between the occipital and ſphenoid bones, which gradually turns thinner as each of the bones advances, till their fibres at laſt run into each other; and, about ſixteen or eighteen years of age, the union of theſe two bones becomes ſo intimate, that a ſeparation cannot be made without violence.—The os occipitis is joined by a double articulation to the firſt vertebra of the neck, each condyle being received into a ſuperior oblique proceſs of that vertebra. What motion is allowed here we ſhall conſider afterwards, where the vertebrae are deſcribed.

The *uſes* of this bone appear from the preceding deſcription, and therefore need not be repeated.

An infant, born at the full time, has this bone divided, by unoſſified cartilages, into four parts.—The firſt of theſe is larger than the other three, is of a triangular ſhape, and conſtitutes all the part of the bone above the great foramen. Fiſſures generally appear in the upper part and ſides of this triangular bone, when all the cartilage is ſeparated by maceration; and ſometimes little diſtinct bones are ſeen towards the edges of it.—The ſecond and third pieces of this bone are exactly alike, and ſituated on each ſide of the great foramen, from which the whole condyles are nearly produced; and they are extended forwards almoſt to the fore part of the hole for the ninth pair of nerves.—The fourth piece is the cuneiform proceſs, which forms a ſmall ſhare of the great hole, and of thoſe for the ninth pair of nerves and of the condyles; betwixt it and the ſphenoid bone, a cartilage is interpoſed.

Of

Of the eight bones which belong to the cranium, there are only two which are not yet deſcribed, viz. the *ethmoid* and *ſphenoid*. Theſe we already mentioned, in complaiſance to the generality of writers on this ſubject, as bones common to the cranium and face, becauſe they enter into the compoſition of both; but the ſame reaſon might equally be uſed for calling the frontal bone a common one too. I ſhall, however, paſs any idle diſpute about the propriety of ranging them, and proceed to examine the ſtructure of the bones themſelves.

OS ETHMOIDES.

Os Ethmoides (*u*), or the *ſieve-like* bone, has got its name from the great number of ſmall holes with which that part of it firſt taken notice of is pierced. When this bone is entire, the figure of it is not eaſily deſcribed; but by a detail of its ſeveral parts, ſome idea may be afforded of the whole; and therefore I ſhall diſtinguiſh it into the *cribriform lamella* with its proceſs, the *naſal lamella*, *cellulæ*, and *oſſa ſpongioſa*.

The thin horizontal lamella is all (except its back-part) pierced obliquely by a great number of ſmall holes, through which the filaments of the olfactory nerves paſs. In a recent ſubject, theſe holes are ſo cloſely lined by the dura mater, that they are much leſs conſpicuous than in the ſkeleton. From the middle of the internal ſide of this plate, a thick proceſs riſes upwards; and being higheſt at the fore-part, gradually becomes lower as it is extended backwards. From ſome reſemblance which this proceſs was imagined to

(*u*) Cribriforme, σηαγγοειδης, ſpongiforme, criſtatum.

have to a cock's comb, it has been called *crista galli* (*x*). The falx is connected to its ridge, and to the unperforated part of the cribriform plate. When the crista is broke, its base is sometimes found to be hollow, with its cavity opening into the nose (*y*). Immediately before the highest part of this process, is the blind hole of the frontal bone, which, as was formerly remarked, is often in a good measure formed by a notch in the fore-part of the root of the crista.

From the middle of the outer surface of the cribriform lamella, a thin solid plate is extended downwards and forwards, having the same common base with the crista galli. Generally it is not exactly perpendicular, but is inclined to one side or other, and therefore divides the cavity of the nose unequally. Its inclination to one side, and flexure in the middle, is sometimes so great, that it fills up a large share of one of the nostrils, and has been mistaken for a polypus there. It is thin at its rise, and rather still thinner in its middle; yet afterwards, towards its lower edge, it becomes thicker, that its conjunction with the bones and middle cartilage of the nose might be firmer.

At a little distance from each side of this external process, a cellular and spongy bony substance depends from the cribriform plate. The number and figure of the cells in this irregular process of each side, are very uncertain, and not to be represented in words; only the cells open into each other, and into the cavity of the nose: The uppermost, which are below the aperture of the frontal sinuses, are formed like funnels. The outer surface of these cells is smooth and plain, where this bone assists in composing the orbit; at which place, on each side, it has got the name of os planum; on the upper edge of which a small notch or two may sometimes be observed, which go to the formation of the internal

(*x*) Verruca præduta, septum ossis spongiosi.

(*y*) Palfyn. Anat. chir. tr. 4. ch. 15.

ternal orbitar holes; as was remarked in the deſcription of the frontal bones.

Below the cells of each ſide, a thin plate is extended inwards; and then bending down, it becomes thick and of a ſpongy texture. This ſpongy part is triangular, with a ſtraight upper edge placed horizontally, an anterior one ſlanting from above, downwards and forwards, and with a pendulous convex one below. The upper and lower edges terminate in a ſharp point behind. The ſide of this pendulous ſpongy part next to the ſeptum narium is convex, and its external ſide is concave. Theſe two proceſſes of the ethmoid bone have got the name of *oſſa ſpongioſa*, or *turbinata ſuperiora*, from their ſubſtance, figure, and ſituation.

All the prominences, cavities, and meanders of this ethmoid bone, are covered with a continuation of the membrane of the noſtrils in a recent ſubject. Its horizontal cribriform is lodged between the orbitar proceſſes of the frontal bone, to which it is joined by the ethmoid future, except at the back part, where it is connected with the cuneiform bone by a future common to both theſe bones, though it is generally eſteemed part of the ſphenoidal ——Where the oſſa plana are contiguous to the frontal bone within the orbit, their conjunction is reckoned part of the tranſverſe future.——Farther forward than the oſſa plana, the cells are covered by the oſſa unguis: which are not only contiuous to theſe cells, but cannot be ſeparated from them without breaking the bony ſubſtance; and therefore, in propriety, thoſe bones ought to be demonſtrated as part of the ethmoid bone.——Below the oſſa unguis and plana, theſe cells and oſſa ſpongioſa are overlapped by the maxillary bones. The cellular part of each palate bone is contiguous to each os planum, and to the cells backwards. The lower edge of the naſal perpendicular plate is received into the furrow of the vomer.

vomer —Its poſterior edge is joined to the fore part of the proceſſus azygos of the ſphenoid bone. Its upper edge joins the naſal proceſs of the frontal and naſal bones; and its anterior one is connected to the middle cartilage of the noſe.

From all which, the *uſes* of this bone are evident, viz. to ſuſtain the anterior lobes of the brain; to give paſſage to the olfactory nerves, and attachment to the falx; to enlarge the organ of ſmelling, by allowing the membrane of the noſe a great extent; to ſtraighten the paſſage of the air through the noſe, by leaving only a narrow winding canal, on the ſenſible membranous ſides of which the ſubſtances conveyed along with the air muſt ſtrike; to form part of the orbit of the eyes and ſeptum narium; while all its parts are ſo light as not to be in hazard of ſeparating by their weight; and they are ſo thin as to form a large ſurface, without occupying much ſpace. This brittle ſubſtance, however, is ſufficiently protected from external injuries by the firm bones which cover it.

If this bone is ſeized on by any corroding matter, we may eaſily conceive what deſtruction may enſue. Hence it is, that an ozæna is difficult to cure; and that in violent ſcurvies, or in the lues venerea, the fabric of the noſe, the eyes, and life itſelf are in danger. The ſituation of the naſal plate may ſhow us how dangerous a fracture of the bones of the noſe may be, when made by a force applied to their middle fore-part, of a perſon in whom this naſal plate is perpendicular.

The ethmoid bone of ripe children is divided into two by a perpendicular cartilage, which, when oſſified, is the criſta galli and naſal plate; but its other parts are oſſified and complete.

OS SPHENOIDES.

Os SPHENOIDES (z), or *wedge-like* bone, ſo called becauſe of its ſituation in the middle of the bones of the cranium and face, is of ſuch an irregular figure, that I know not any thing to which it may be compared, unleſs, perhaps, it bear ſome faint reſemblance to a bat with its wings extended.

When we view the external ſurface of the os ſphenoides, two or three remarkable proceſſes from each ſide of it may be obſerved, which are all of them again ſubdivided. The firſt pair is the two large lateral proceſſes or wings; the upper part of each of which is called the *temporal proceſs*, becauſe they join with the temporal bones in forming the temples, and the ſeat for ſome ſhare of the crotaphite muſcles. That part of the wings which juts out towards the inſide, ſomewhat lower than the temporal apophyſes, and is ſmooth and hollowed, where it makes up part of the orbit, is thence named *orbitar proceſſes*. Behind the edge ſeparating theſe two proceſſes, there is often a ſmall groove, made by a branch of the ſuperior maxillary nerve, in its paſſage to the temporal muſcle. The loweſt and back part of each wing, which runs out ſharp to meet the oſſa petroſa, has been ſtyled the *ſpinous* proceſs: from near the point of which a ſharp-pointed proceſs is frequently produced downwards, commonly called *ſtyliform*, that affords origin to the ptery-ſtaphylinus externus muſcle. From this ſtyloid proceſs a very ſmall groove is extended along the edge of the bone to the hollow at the root of the internal plate of the following proceſſes, which

(z) Cuneiforme, πολυμορφον, multiforme, paxillum, cribratum palati, colatorii, cavilla, baſilare.

which forms part of the Euſtachian tube (*a*). The ſecond pair of external proceſſes of the cuneiform bone, is the two which ſtand out almoſt perpendicular to the baſe of the ſkull. Each of them has two plates, and a middle foſſa facing backwards; and ſhould, to carry on our compariſon, be likened to the legs of a bat; but are commonly ſaid to reſemble the wings of that creature; and therefore are named *pterygoid or aliform* (*b*) proceſſes. The external plates are broadeſt, and the internal are longeſt. From each ſide of the external plates the pterygoid muſcles take their riſe. At the root of each internal plate, a ſmall hollow may be remarked, where the muſculus ptery-ſtaphilinus internus or circumflexus palati riſes, and ſome ſhare of the cartilaginous end of the Euſtachian tube reſts: and, at the lower end of the ſame plate, is a hook like riſing or proceſs, round which the tendon of the laſt-named muſcle plays, as on a pulley. From the edge of the external plates ſome ſmall ſharp ſpikes ſtand out; but their number and bulk are uncertain.——To theſe another pair may be added, to wit, the little triangular thin proceſs, which comes from each ſide of the body of the ſphenoid bone, where the pterygoid proceſſes are riſing from it, and are extended over the lower part of the aperture of the ſinus as far as to join the ethmoid bone, while their body hangs down into the nares (*c*). Beſides theſe pair of proceſſes, there is a ſharp ridge which ſtands out from the middle of its baſe: Becauſe it wants a fellow, it may be called *proceſſus azygos*. The lower part of this proceſs, where it is received into the vomer, is thick, and often not quite perpendicular, but inclining more to one ſide than the other. The fore-part of this proceſs, where it joins the naſal plate of the os ethmoides,

(*a*) Winſlow Expoſ. anatomique du corps humain, traité des os ſecs, § 233.

(*b*) Naviculares.

(*c*) Albin. Tab. Oſſ. 5. fig. 2. 6. A. A.——Bertin. Mem. de l'acad. des ſciences, 1744.—Sue, planche viii. fig. 2, 3, 4, 5, 6.

des, is thin and ſtraight. Theſe two parts have been deſcribed as two diſtinct proceſſes by ſome writers.

The depreſſions, ſinuoſities, and foſſae, on the external ſurface of this ſphenoid bone, may be reckoned up to a great number, viz. Two on the temporal apophyſes, where the crotaphite muſcles lodge—Two on the orbitar proceſſes, to make way for the globes of the eyes—Two between the temporal and ſpinous proceſſes, for receiving the temporal bones —Two between the plates of the pterygoid proceſſes, where the muſculi pterygoidei interni and ptery-ſtaphylini interni are placed—Two between the pterygoid and orbitar proceſſes, for forming the holes common to this and to the cheek and maxillary bones—Two on the lower ends of the aliform proceſſes, which the palate bones enter into.—Two at the roots of the temporal and pterygoid proceſſes, where the largeſt ſhare of the external pterygoid muſcles have their riſe—Two at the ſides of the proceſſus azygos, for forming part of the noſe, &c.

What I deſcribed under the name of *temporal* and *ſpinous proceſſes* on the outſide of the ſkull, are likewiſe ſeen on its inſide, where they are concave, for receiving part of the brain; and commonly three apophyſes on the internal ſurface of the ſphenoid bone are only mentioned.——Two riſing broad from the fore-part of its body, become ſmaller as they are extended obliquely backwards.——The third ſtanding on a long tranſverſe baſe, near the back part of the body of this bone, riſes nearly erect, and of an equal breadth, terminating often in a little knob on each ſide. The three are called *clinoid*, from ſome reſemblance which they were thought to have to the ſupporters of a bed. Sometimes one or both the anterior clinoid proceſſes are joined to the ſides of the poſterior one, or the body of the bone itſelf. From the roots of the anterior clinoid proceſſes the bone is

extended

extended on each ſide outwards and forwards, till it ends in a ſharp point, which may have the name of the *tranſverſe* ſpinous proceſſes. Between, but a little farther back than the two anterior clinoid proceſſes, we ſee a protuberance conſiderably ſmaller than the poſterior clinoid proceſs, but of its ſhape.——Another proceſs from between the tranſverſe proceſſes, often forces itſelf forwards into the os ethmoides.

Within the ſkull, there are two *ſinuoſities* in the internal part of each wing of the ſphenoid bone, for receiving the middle part of the brain.——One between the tranſverſe ſpinous proceſſes, for lodging the part of the brain where the crura medullae oblongatae are.——Immediately before the third or middle clinoid proceſs, a ſingle pit may generally be remarked, from which a foſſa goes out on each ſide to the holes through which the optic nerves paſs. The pit is formed by the conjoined optic nerves; and in the foſſa theſe nerves are lodged, as they run divided within the ſkull. Between that third protuberance and the poſterior clinoid proceſs, the larger pit for the glandula pituitaria may be remarked. This cavity, becauſe of its reſemblance to a Turkiſh ſaddle, is always deſcribed under the name of *ſella turcica*, or *ephippium*——On the ſides of the poſterior clinoid proceſs, a foſſa may be remarked, that ſtretches upwards, then is continued forwards along the ſides of the ſella turcica, near to the anterior clinoid proceſſes, where a pit on each ſide is made. Theſe foſſae point out the courſe of the two internal carotid arteries, after they have entered the ſkull. Beſides all theſe, ſeveral other foſſae may be obſerved, leading to the ſeveral holes, and imprinted by the nerves and blood veſſels.

The *holes* on each ſide of the os ſphenoides are ſix proper, and three common.——The firſt is the round one immediately

mediately below the anterior clinoid proceſſes, for the paſſage of the optic nerve, and of the branch of the internal carotid artery that is ſent to the eye.——The ſecond is the foramen lacerum, or large ſlit between the tranſverſe ſpinous and orbitar proceſſes: The interior end of which ſlit is large; and, as it is extended outwards, it becomes narrower. The outer end of it is formed in the os frontis; and therefore this might be reckoned among the common foramina. Through it the third, fourth, the firſt branch of the fifth, and the greater ſhare of the ſixth pair of nerves, and an artery from the internal carotid, go into the orbit. Sometimes a ſmall branch of the external carotid enters near its end, to be diſtributed to the dura mater (*d*); and a vein, ſome call it the *venous duct*, or *Nuck's aqueduct*, returns through it to the cavernous ſinus.——The third hole, ſituated a little below the one juſt now deſcribed, is called *rotundum*, from its ſhape. It allows paſſage to the ſecond branch of the fifth pair of nerves, or ſuperior maxillary nerve, into the bottom of the orbit.——The fourth is the *foramen ovale*, about half an inch behind the round hole. Through it the third branch of the fifth pair, or inferior maxillary nerve, goes out; and ſometimes a vein from the dura mater paſſes out here (*e*).—— Very near the point of the ſpinous proceſs is the fifth hole of this bone; it is ſmall and round, for a paſſage to the largeſt artery of the dura mater, which is often accompanied with a vein.——The ſixth proper hole (*f*) cannot be well ſeen, till the cuneiform bone is ſeparated from all the other bones of the cranium; for one end of it is hid by a ſmall

(*d*) Winſlow, Expoſition anatomique du corps humain, traité des arteries, § 60. et de la tete, § 26.

(*e*) Ingraſſ. Commentar. in Galen. de oſſib. lib. 1. comment. 8.

(*f*) Veſal. Anat. lib. 1. cap. 12.——Euſtach. tab. 46. fig. 13. & 16.—— Vidus Vidius, Anat. lib. 2. cap. 2. explicat. tab. 5. & tab. 6. fig. 8. 9. 10. lit. O.

protuberance of the internal plate of the pterygoid proceſs, and by the point of the proceſſus petroſus of the temporal bone. Its canal is extended above the inner plate of the pterygoid proceſs; and where it opens into the cavity of the noſe, it is concealed by the thin laminous part of the palate-bone. Through it a conſiderable branch of the ſecond branch of the fifth pair of nerves is reflected.—In the middle of the ſella turcica, a ſmall hole or two often pierce as far as the cellular ſubſtance of the bone; and ſometimes at the ſides of this ſella, one or more ſmall holes penetrate into the ſphenoidal ſinuſes. Theſe obſervations afforded ſome anatomiſts (*g*) an argument of weight in their days in defence of Galen (*h*), who aſſerted the deſcent of the pituita that way into the ſinuſes below.

The firſt of the *common holes* is that unequal fiſſure at the ſide of the ſella turcica, between the extreme point of the os petroſum and the ſpinous proceſs of the cuneiform bone. This hole only appears after the bones are boiled; for in a recent ſubject its back-part is covered by a thin bony plate that lies over the internal carotid artery, and farther forward it is filled with a cartilaginous ligament, under which the cartilaginous part of the Euſtachian tube is placed: It was by this paſſage that the antients believed the ſlimy matter was conveyed from the emunctory of the brain, the glandula pituitaria, to the fauces.——The ſecond common hole is the large diſcontinuation of the external ſide of the orbit, left between the orbitar proceſſes of the cuneiform bone, the os maxillare, malæ, and palati. In this large hole the fat for lubricating the globe of the eye and temporal muſcles is lodged, and branches of the ſuperior maxillary nerve, with ſmall arteries from the carotid, and veins, paſs.——The third

(*g*) Joc. Syl. Calumniæ ſecundæ amolitio.——Laurent. Hiſt. Anat. lib. ii. queſt. 11.

(*h*) Galen. de uſu part. lib. ix. cap. 1.

third hole is formed between the baſe of this bone and the root of the orbitar proceſs of the palate-bone of each ſide. Through this a branch of the external carotid artery, and of the ſecond branch of the fifth pair of nerves, are allowed a paſſage to the noſtrils, and a returning vein accompanies them. Sometimes, however, this hole is proper to the palate-bone, being entirely formed out of its ſubſtance.

Under the ſella turcica, and ſome way farther forward, but within the ſubſtance of the ſphenoid bone, are two *ſinuſes*, ſeparated by a bony plate. Each of them is lined with a membrane, and opens into the upper and back part of each noſtril by a round hole, which is at their upper fore-part. This hole is not formed only by the os ſphenoides, which has an aperture near as large as any tranſverſe ſection of the ſinus, but alſo by the palate-bones, which are applied to the fore-part of theſe ſinuſes, and cloſe them up, that hole only excepted which was already mentioned. The two ſinuſes are frequently of unequal dimenſions, and ſometimes there is only one large cavity, with an opening into one noſtril. Theſe cavities are likewiſe ſaid (*i*) to be extended ſometimes as far back as the great foramen of the occipital bone. In other ſubjects they are not be found, when the bone is compoſed of large cells (*k*). Some (*l*) mention a cavity within the partition of the ſinuſes; but it is ſmall. The ſphenoidal ſinuſes ſerve the ſame uſes as the frontal.

As this bone is extremely ragged and unequal, ſo its ſubſtance is of very different thickneſs, being in ſome places diaphanous; in others it is a middle thickneſs, and its middle back part ſurpaſſes the greateſt ſhare of the cranium in thickneſ.

The os ſphenoides is joined by its wings to the parietal bones

(*i*) Albin. de oſſib. § 39. (*k*) Veſal. lib. 1. cap. 6. (*l*) Id. ibid.

bones above, to the os frontis and offa malarum before, to the temporal bones behind;—by the fore-part of its body and ſpinous proceſſes, to the frontal and ethmoid bones;—by its back-part, behind the two ſinuſes, to the occipital, where it looks like a bone with the epiphyſes taken off, and, as was formerly obſerved in the deſcription of the occipital bone, it cannot be ſeparated without violence in adults;——to the palate-bones, by the ends of the pterygoid proceſs, and ſtill more by the fore-part of the internal plates of the pterygoid proceſſes, and of the ſinuſes;——to the maxillary bones, by the fore-part of the external pterygoid plates;—to the vomer and naſal plate of the os ethmoides, by the proceſſus azygos. All theſe conjunctions, except the laſt, which is a ſchindyleſis, are ſaid to be by the future proper to this bone; though it is at firſt ſight evident, that ſeveral other futures, as the tranſverſe, ethmoidal, &c. are confounded with it.

We ſee now how this bone is joined to all the bones of the cranium, and to moſt of the upper jaw; and therefore obtained the name of the *wedge-like bone*.

The uſes are ſo blendid with the deſcription, as to leave nothing new to be added concerning them.

The ſphenoidal bone is almoſt complete in a foetus of nine months: only the great alae ſeparate after maceration from the body of the bone.——The proceſſus azygos is very large and hollow;——the thin triangular proceſſes are not offified;——the internal ſurface of the body is unequal and porous;——the ſinuſes do not appear.

Whoever is acquainted with each bone of the cranium, can, without difficulty, examine them as they ſtand united, ſo as to know the ſhapes, ſizes, diſtances, &c. of their ſeveral parts, and the forms, capacities, &c. of the cavities formed by them; which is of great uſe towards underſtanding the anatomy of the parts contiguous to, contained

ed within, or connected to them. Such a review is neceſſary, after conſidering each claſs of bones. Thus the orbits, noſtrils, mouth, face, head, ſpine, thorax, pelvis, trunk, extremities, and ſkeleton, ought likewiſe to be examined.

§ 2. *Bones of the* FACE.

THE face is the irregular pile of bones compoſing the fore and under part of the head, which is divided by authors into the *upper* and *lower maxillae* or jaws.

THE *ſuperior maxillae* (*m*) is the common deſignation given to the upper immoveable ſhare of the face; though, if we would follow Celſus (*n*), we ſhould apply the word *maxillae* to the lower jaw only, and the name *mala* to this upper jaw. In compliance to prevailing cuſtom, I ſhall, however, uſe the terms as they are now commonly employed. The ſhape of the ſuperior jaw cannot eaſily be expreſſed; nor is it neceſſary, provided the ſhape and ſituation of all the bones which compoſe it are deſcribed.—— It is bounded above by the tranſverſe ſuture, behind by the fore-part of the ſphenoid bone, and below by the mouth.

The upper jaw conſiſts of ſix bones on each ſide; of a thirteenth bone which has no fellow, placed in the middle; and of ſixteen teeth. The thirteen bones are, two oſſa naſi, two oſſa unguis, two oſſa malarum, two oſſa maxillaria, two oſſa palati, two oſſa ſpongioſa inferiora, and the vomer.

The *oſſa naſi* are placed at the upper part of the noſe; *oſſa unguis* are at the internal canthi of the orbits; *oſſa malarum* form the prominence of the cheeks; *oſſa maxillaria* form

(*m*) Σιαγων, γενυς, mandibula. (*n*) Lib. 8. cap. 1.

form the ſide of the noſe, with the whole lower and fore-part of the upper jaw, and the greateſt ſhare of the roof of the mouth; *oſſa palati* are ſituated at the back part of the palate, noſtrils, and orbit; *oſſa ſpongioſa* are ſeen in the lower part of the nares; and the *vomer* helps to ſeparate theſe two cavities.

The bones of the upper jaw are joined to the bones of the ſkull by the *ſchindyleſis* and ſutures already deſcribed as common to the cranium and face, and they are connected to each other by *gomphoſis* and fifteen ſutures.

The *gomphoſis* is only where the teeth are fixed in their ſockets, and the *ſchindyleſis* is only where the edges of the vomer are joined to other bones.

The *futures* are generally diſtinguiſhed by numbers, which have been differently applied; and therefore I join thoſe (*o*) who prefer the giving names to each, which may be eaſily contrived from their ſituation, or from the bones which they connect.

The firſt is the anterior naſal (*p*), which is ſtraight, and placed longitudinally in the middle fore-part of the noſe.

The ſecond and third are the lateral naſal (*q*), which are at each ſide of the noſe, and almoſt parallel to the firſt future.

Each of the two lacrymal is almoſt ſemicircular, and is placed round the lacrymal groove.

The ſixth and ſeventh are the internal orbitar: each of which is extended obliquely from the middle of the lower ſide of an orbit to the edge of its baſe.

The two external orbitars are continued, each from the end of the internal orbitar to the under and fore-part of the cheek.

The tenth is the myſtachial, which reaches only from the

(*o*) Vander Linden. Medicin. phyſiolog. cap. 13. art. 2. § 10.——Rolfinc. Anat. lib. 2. cap. 25.—Schenk. Schol. part. ſect. ult. par. 2. cap. 5.

(*p*) Naſalis recta. (*q*) Naſalis obliqua.

the lower part of the ſeptum narium to between the two middle dentes inciſores.

The longitudinal palate ſuture (*r*) ſtretches from the middle of the foremoſt teeth through the middle of all the palate.

The tranſverſe palate ſuture (*s*) runs acroſs the palate, nearer the back than the fore-part of it.

Each of the two palato-maxillary is at the back-part of the ſide of each noſtril.

The fifteenth is the ſpinous, which is in the middle of the lower part of the noſtrils. This may perhaps be rather thought a double *ſchindyleſis*.

The connection of the offa ſpongioſa to the ſide of each noſtril, is ſo much by a membrane in young ſubjects, by a ſort of hook and afterwards by concretion or union of ſubſtance in adults, that I did not know well how to rank it: But if any chuſes to call it a ſuture, the addition of two *tranſverſe naſal* ſutures may be made to thoſe above named.

The ſutures of the face (formerly called *harmoniæ*) have not ſuch conſpicuous indentations as thoſe of the ſkull; the bones here not having ſubſtance enough for forming large indentations, and there being leſs neceſſity for ſecurity againſt external injuries, or any internal protruding force, than in the cranium.—Theſe ſutures often diſappear in old people, by the bones running into each other; which can do little prejudice, becauſe the principal uſe of the bones being ſo numerous here, is to allow them to be extended into a proper form.

It is evident, from the manner of the conjunction of theſe bones, that they can have no motion, except in common with the cranium.

The purpoſes which this pile of bones ſerves, will be ſhown in the deſcription which I am going to give of each of them.

OSSA

(*r*) Laquearis, palataria recta. (*s*) Arcuata, palatina poſtica.

OSSA NASI.

Ossa nasi, ſo named from their ſituation at the root of the noſe, are each of an irregular oblong ſquare figure, being broadeſt at their lower end, narroweſt a little higher than their middle, and becoming ſomewhat larger at the top, where they are ragged and thickeſt, and have a curvature forwards, that their connection with the frontal bone might be ſtronger.—Theſe bones are convex externally, and thereby better reſiſt any violence from without; and they are concave internally, for enlarging the cavity of the noſe.

The lower edge of theſe bones is unequal, and is ſtretched outwards and backwards, to join the cartilages of the noſtrils. Their anterior ſide is thick, eſpecially above, and unequal, that their conjunction to each other might be ſtronger; and a ſmall riſing may be remarked on their inner edge, where they are ſuſtained by the ſeptum narium. —Their poſterior ſide, at its upper half, has externally a depreſſion, where it is a little overlapped by the maxillary bones, while its lower half covers theſe bones: By which contrivance, they do not yield eaſily to preſſure applied to their fore-part or ſides.

A ſmall hole is frequently to be obſerved on their external ſurface, into which two, three, or four holes, which appear internally, terminate for the tranſmiſſion of ſmall veins: Sometimes the holes go no farther than the cancelli of the bones.

The naſal bones are firm and ſolid, with very few cells of cancelli in them; the thin ſubſtance of which they conſiſt not requiring much marrow.

They are *joined* above to the frontal bone, by the middle of the tranſverſe future;——behind, to the maxillary bones, by the lateral naſal futures;——below, to the cartilages

tilages of the noſe;—before, to one another, by the anterior naſal ſuture;—internally, to the ſeptum narium.

The *uſe* of theſe bones is to cover and defend the root of the noſe.

In an infant the naſal bones are proportionally ſhorter and leſs thick at their upper part, than in an adult; but are otherwiſe complete.

OSSA UNGUIS, OR LACRYMALIA.

OSSA UNGUIS, or LACRYMALIA, are ſo named, becauſe their figure and magnitude are nearly like thoſe of a nail of one's finger, and becauſe the tears paſs upon them into the noſe.

Their external ſurface is compoſed of two ſmooth concavities and a middle ridge.——The *depreſſion* behind forms a ſmall ſhare of the orbit for the eye-ball to move on; and the one before is a deep perpendicular canal, or *foſſa*, larger above than below, containing part of the lacrymal ſac and duct. This is the part that ought to be pierced in the great operation for the fiſtula lacrymalis.———This foſſa of the bone is cribriform, or has a great number of ſmall holes through it, that the filaments for the membrane which lines it, inſinuating themſelves into theſe holes, might prevent a ſeparation of the membrane, and ſecure the bone in its natural ſituation. ——The *ridge* between theſe two cavities of the os unguis is the proper boundary of the orbit at its internal canthus; and beyond which ſurgeons ſhould not proceed backwards in performing operations here.——The internal or poſterior ſurface of this bone conſiſts of a furrow in the middle of two convexities.

The *ſubſtance* of the os unguis is as thin as paper, and very brittle; which is the reaſon that thoſe bones are often

wanting in ſkeletons, and need little force to pierce them in living ſubjects.

Each of thoſe bones is *joined*, above, to the frontal bone, by part of the tranſverſe ſuture;—behind to the os planum of the ethmoid bone, by the ſame ſuture;—before, and below, to the maxillary bone, by the lacrymal ſuture.——Internally, the offa unguis cover ſome of the ſinus ethmoidales; nay, are really continuous with the bony lamellae which make up the ſides of theſe cells; ſo that they are as much part of the ethmoid bone as the offa plana.

Theſe unguiform bones compoſe the interior internal parts of the orbits, lodge a ſhare of the lacrymal ſac and duct, and cover the ethmoid cells.——Their ſituation and tender ſubſtance make a raſh operator in danger of deſtroying a conſiderable ſhare of the organ of ſmelling, when he is performing the operation of the fiſtula lacrymalis: but when theſe bones are hurt, they caſt off without much difficulty, and conſequently the wound is ſoon cured, unleſs the patient labours under a general cacoethes, or there is a prediſpoſition in the bones to caries; in which caſe, a large train of bad ſymptoms follow, or at beſt the cure proves tedious.

The bones are fully formed in a new-born child.

OSSA MALARUM.

Ossa malarum (*t*) was the name given by Celſus, as was already remarked, to all the upper jaw; but is now appropriated to the prominent ſquare bones which form the cheek on each ſide.—Before, their ſurface is convex and ſmooth; backward, it is unequal and concave, for lodging part of the zygomatic muſcles.

The four *angles* of each of theſe bones have been reckoned proceſſes

(*t*) Jugalia vel zygomatica, hypopia, ſubocularia.

proceſſes by ſome authors.—The one at the external canthus of the orbit, called the *ſuperior orbitar* proceſs, is the longeſt and thickeſt.—The ſecond terminates near the middle of the lower edge of the orbit in a ſharp point, and is named the *inferior orbitar* proceſs.—The third, placed near the lower part of the cheek, and thence called *maxillary*, is the ſhorteſt, and neareſt to a right angle.—The fourth, which is called *zygomatic*, becauſe it is extended backwards to the *zygoma* of the temporal bone, ends in a point, and has one ſide ſtraight and the other ſloping.—Between the two orbitar angles there is a concave arch, which makes about a third of the external circumference of the orbit, from which a fifth proceſs is extended backwards within the orbit, to form near one third of that cavity; and hence it may be called the *internal orbitar* proceſs.——From the lower edge of each of the oſſa malarum, which is between the maxillary and zygomatic proceſſes, the maſſeter muſcle takes its origin; and from the exterior part of the zygomatic proceſs, the muſculus diſtortor oris riſes; in both which places the ſurface of the bone is rough.

On the external ſurface of each cheek-bone, one or more ſmall *holes* are commonly found, for the tranſmiſſion of ſmall nerves or blood-veſſels from, and ſometimes into, the orbit.—On the internal ſurface are the holes for the paſſage of the nutritious veſſels of theſe bones.—A notch on the outſide of the internal orbitar proceſs of each of theſe bones aſſiſts to form the great ſlit common to this bone and to the ſphenoid, maxillary, and palate bones.

The *ſubſtance* of theſe bones is, in proportion to their bulk, thick, hard, and ſolid, with ſome cancelli.

Each of the oſſa malarum is *joined*, by its ſuperior and internal orbitar proceſſes, to the os frontis, and to the orbitar proceſs of the ſphenoid bone, by the tranſverſe ſuture;

by

by the edge between the internal and inferior orbitar proceſſes, to the maxillary bone, by the internal orbitar future; by the ſide between the maxillary and inferior orbitar proceſs, again to the maxillary bone, by the external orbitar future; by the zygomatic proceſs, to the os temporum, by the zygomatic future.

The cheek-bones are entire, and fully oſſified in all their parts, in infants.

OSSA MAXILLARIA SUPERIORA.

Ossa Maxillaria Superiora are the largeſt bones, and conſtitute the far greater part, of the upper jaw, which has appropriated the name of *maxillaria* to them. The figure of one of them, or of the two when joined, is ſo irregular, that words can ſcarce give an idea of it.

The *proceſſes* of each os maxillare may be reckoned ſeven. The firſt is the long naſal one at its upper and fore-part, which is broad below, and turns ſmaller as it riſes upwards, to make the ſide of the noſe. At the root of this a tranſverſe ridge may be obſerved within the noſtrils, which ſupports the fore-part of the upper edge of the os ſpongioſum inferius. The ſecond is produced backwards and outwards, from the root of the naſal proceſs, to form the lower ſide of the orbit; and therefore may be called *orbitar*. The edge of this orbitar proceſs, and the ridge of the naſal one, which is continued from it, make a conſiderable portion of the external circumference of the orbit. From the proper orbitar proceſs, a very rough triangular ſurface is extended downwards and outwards, to be connected to the cheek-bone; and therefore may be called the molar proceſs, from the loweſt protuberant part of which ſome ſhare of the maſſeter muſcle takes its riſe. Behind the orbitar proceſs, a large tuberoſity or bulge of the bone

bone appears, which is eſteemed the fourth proceſs. On the internal part of this we often meet with a ridge, almoſt of the ſame height with that in the naſal proceſs, which runs tranſverſely, and is covered by a ſimilar ridge of the palate-bone, on which the back-part of the upper edge of the os ſpongioſum inferius reſts.——The convex back-part of this tuberoſity is rough for the origin of part of the external pterygoid muſcle (*u*); and more internal is ſcabrous, where the palate and ſphenoid bones are joined to it.—That ſpongy protuberance (*x*) at the lower circumference of this bone, where the ſockets for the teeth are formed, is reckoned the fifth. The ſixth is the horizontal plate, which forms the greater part of the baſe of the noſtrils, and roof of the mouth: its upper ſurface, which belongs to the noſtrils, is very ſmooth; but the other below is arched and rough, for the ſtronger adheſion of the membrane of the mouth, which is ſtretched upon it; and in chewing, ſpeaking, &c. might otherwiſe be liable to be ſeparated. The ſeventh riſes like a ſpine from the inner edge of the laſt, and forms a ſmall part of the partition of the noſtrils.

The *depreſſions* in each maxillary bone are, 1. A ſinuoſity behind the orbitar proceſs, made by the temporal muſcle. 2. A pit immediately before the ſame proceſs, where the origin of the muſculus elevator labiorum communis, and elevator labii ſuperioris, with a branch of the fifth pair of nerves, are lodged ſecurely. 3. The hollow arch of the palate. 4. The ſemicircular great notch, or entry to the lower part of the noſtrils, between the root of the naſal proceſs and ſpine of the palate plate. —— Below this, the fore part of the bone is flatted, or ſometimes hollowed, by the muſculus depreſſor labii ſuperioris. 5. Sockets for the teeth (*y*): The number of theſe ſockets is uncertain; for

(*u*) Albin. de oſſib. § 79. (*x*) φατνια.

(*y*) Βοθρια ὀλμισκων, Alveoli, foſſulae, mortariola, fraena, locelli, cavae, loculamenta.

for the ſame number of teeth is not in all people, and the four fartheſt teeth of each ſide in each jaw vary greatly in their number of roots; and when the teeth of a living perſon fall out or are taken away, the ſockets fill up with an oſſeous net work, which becomes ſolid afterwards. 6. The lacrymal foſſa in the naſal proceſs, which aſſiſts the os unguis to form a paſſage for the lacrymal duct. The part of the bone forming this foſſa is ſo firm and ſtrong, that a ſurgeon can ſcarcely perforate it with the ordinary inſtruments for the fiſtula lacrymalis; and therefore ought to avoid it in doing this operation.——Immediately on the outſide of this, there is a ſmall depreſſion, from which the inferior or leſſer oblique muſcle of the eye has its origin (z). 7. The canal on the upper part of the great tuberoſity within the orbit, which is almoſt a complete hole; in this a branch of the ſuperior maxillary nerve paſſes. Beſides theſe, the ſuperior ſurface of the great bulge is concave, to receive the under part of the eye. Immediately above the tranſverſe ridge in the naſal proceſs, a ſmall hollow is formed by the os ſpongioſum. In ſome ſubjects, the naſal proceſs has a ſmall round pit above the lacrymal duct, where the little tendon or ligament of the orbicular muſcle of the eye-lids is inſerted. It is this tendon, and not the tendon of the larger oblique muſcle of the eye, which there is ſome hazard of cutting in the operation of the fiſtula lacrymalis.

The *holes* of this bone are two proper and two common, which are always to be found; beſides ſeveral others, whoſe magnitude, number, &c. are uncertain. The firſt of the proper is the external orbitar, immediately below the orbit, by which the infra-orbitar branch of the ſecond branch of the fifth pair of nerves, and a ſmall artery, come out, after having paſſed in the canal, at the bottom of

(z) Winſlow, Expoſition anatomique des os ſecs, § 276.

of the orbit, deſcribed number 7. of the depreſſions. This hole is often double, and that when the nerve has happened to ſplit before it has eſcaped from the bone. The ſecond is the foramen inciſivum, juſt behind the fore teeth; which, at its under part, is one irregular hole common to both the maxillary bones when they are joined; but, as it aſcends, ſoon divides into two, three, or ſometimes more holes; ſome of which open into each noſtril. Through them ſmall arteries and veins, and a twig of the ſecond branch of the fifth pair of nerves, paſs, and make a communication between, or join the lining coats of the noſe and mouth. In ſome ſubjects, Steno's duct may be traced ſome length on the ſide of theſe paſſages next to the noſe, and ſmall orifices may be obſerved opening into the mouth.

The firſt common hole is that which appears at the inner ſide of the back part of the tuberoſity and of the ſockets of the teeth; and is formed by a foſſa in this bone, and a correſponding one in the os palati: Through it a nerve, which is a branch of the ſecond branch of the fifth pair, runs to the palate. The other common hole is the great ſlit in the outſide of the orbit, deſcribed already as the ſecond common hole of the ſphenoid bone.

On the naſal proceſs holes may be often obſerved for the paſſage of veſſels to the ſubſtance of the bones; and, at the back-part of each tuberoſity, ſeveral foramina are placed, for the tranſmiſſion of nerves to the cavity within: But theſe are uncertain.

All the *body* of the maxillary bone is hollow, and leaves a large *ſinus* like the frontal and ſphenoid, which is commonly, but unjuſtly, called *antrum Highmorianum* (*a*). When the os maxillare is ſingle, or ſeparated from all the other bones of a ſkeleton, its antrum appears to have a large aperture

(*a*) Genæ.

ture into the noſtrils; but, in a recent ſubject, it is ſo covered at its back-part by the palate-bone, in the middle by the os ſpongioſum inferius, before by a ſtrong membrane, that one or ſometimes two holes, ſcarcely ſo large as a crow-quill, are only left at the upper part; which, after a ſhort winding proceſs, open into the noſtrils between the two oſſa ſpongioſa.—At the bottom of this cavity, we may often obſerve ſome protuberances, in which the ſmall points of the roots of the teeth are contained (*b*).—This cavern and the ſockets of the teeth are often divided by the interpoſition only of a very thin bony plate, which is liable to be eroded by acrid matter collected in the antrum, or to be broke in drawing a tooth (*c*). The ſymptoms of a collection of matter here naturally led us to the practice of pulling out the teeth, and piercing through this plate into the antrum, to procure an evacuation of the collected matter; by which conſiderable ſervice is frequently done (*d*).

The maxillary ſinuſes have the ſame *uſes* as the frontal and ſphenoidal: and the ſituation of the ſinuſes is ſuch, that the liquor drilling from them, from the cells of the ethmoid and palate bones, and from the lacrymal ducts, may always moiſten all the parts of the membrane of the nares in the different ſituations of the head.

Though the membranes which line the frontal, ſphenoidal, and maxillary ſinuſes, are continuations of the one which covers the bones within the noſe; yet they are much thinner than it is, and have ſo much ſmaller veſſels, that the injection which makes the membrane of the noſe red all over, fills only ſome few veſſels of the maxillary ſinuſes, and is ſcarcely obſerved in the frontal and ſphenoidal.

(*b*) Highmore, Diſquiſ. Anat. lib. 3. part. 2. cap. 7.

(*c*) Highmore, ibid.

(*d*) Cowper in Drake's Anthropol. book 3. chap. 10.——Medical Eſſays and Obſerv. vol. 5. art. 30.

dal. Are not the larger veſſels intended for a more plentiful ſecretion of a viſcid liquor to defend the membrane from the effects of the perſtatus which is conſtantly through the noſe? Are not the membranes which have the ſmalleſt veſſels, *cæteris paribus*, the moſt ſenſible? Do not many phenomena of ſmelling, inflammations of theſe parts, megrim, polypi, &c. depend on this ſtructure of the membranes?

The *ſubſtance* of the oſſa maxillaria is compact and firm, except at the inferior proceſſes, in which the teeth are lodged, where it is very ſpongy.

The maxillary bones are joined above, by the upper ends of their naſal proceſſes to the os frontis, by the tranſverſe ſuture;—at the ſides of theſe proceſſes, to the oſſa unguis, by the lacrymal ſuture;—to the naſal bones, by the lateral naſal ſutures;—by their orbitar proceſſes, to the cheek-bones, by the external orbitar ſutures;—by the internal ſides of the internal orbitar proceſſes, to the oſſa plana, by part of the ethmoidal ſuture;—by the back-part of the tuberoſities, to the palate-bones, by the ſuturæ palato-maxillares;—by the poſterior edges of their palatine lamellæ, to the oſſa palati, by the tranſverſe palate ſuture;—by their naſal ſpines, to the vomer, by the ſpinous ſuture;—by their ſocket, to the teeth, by the gomphoſis;—by the internal edge of the palate-plate, to one another, by the longitudinal palate ſuture, on the upper and fore-part of which a furrow is left for receiving the cartilage which forms the partition of the noſtrils;—between the fore-part of the noſtrils and mouth, to each other, by the myſtachial ſuture:—Sometimes they are connected to the oſſa ſpongioſa inferiora, by a plain concretion or union of ſubſtance.

Theſe bones form the greater part of the noſe and of the roof of the mooth, and a conſiderable ſhare of the orbit.

They contain ſixteen teeth, give riſe to muſcles, tranſmiſſion to nerves, &c. as mentioned in the deſcription of their ſeveral parts.

In each of the maxillary bones of a new-born child, the external orbitar proceſs is hollow, with remarkable holes in it;—there are five ſockets for the teeth, of which the two poſterior are very large, and, when divided by a ſecond croſs partition, make the number of ſockets ſix (*e*). The palate-plate is cribriform about the middle. The great tuberoſity is not formed;—inſtead of the antrum, there is only an oblong depreſſion at the ſide of the noſtrils.

OSSA PALATI.

OSSA PALATI are commonly deſcribed as two ſmall ſquare bones, at the back-part of the palate or roof of the mouth, though they are of much greater extent, being continued up the back-part of the noſtrils to the orbit (*f*). Each palate-bone may therefore be divided into four parts, the palate ſquare-bone, the pterygoid proceſs, naſal lamella, and orbitar proceſs.

The *ſquare-bone* is unequally concave, for enlarging both the mouth and cavity of the noſe. The upper part of its internal edge riſes in a ſpine, after the ſame manner as the palate-plate of the maxillary bone does, to be joined with the vomer. Its anterior edge is unequally ragged, for its firmer connection with the palate-proceſs of the os maxillare. The internal edge is thicker than the reſt, and of an equal ſurface, for its conjunction with its fellow of the other ſide.

(*e*) Alb'n. Oſteogen, tab. 5. fig. 45.—Ungebav. de dentit. ſecund. jun. § 1.

(*f*) Euſtach. tab. 47. fig. 1. 3. 6. 7. 8.———Vidus Vidius, de nat. lib. 2. cap. 2. explicat. tab. 6. fig. 19.——Winſlow, Memoires de l'acad. des ſciences, 1720.

ſide. Behind, this bone is ſomewhat in form of a creſcent, and thick, for the firm connection of the velum pendulum palati; the internal point being produced backwards, to afford origin to the palato-ſtaphylinus or azygos muſcle This ſquare bone is well diſtinguiſhed from the pterygoid proceſs by a perpendicular foſſa, which applied to ſuch another in the maxillary bone, forms a paſſage for the palatine branch of the fifth pair of nerves; and by another ſmall hole behind this, through which a twig of the ſame nerve paſſes.

The *pterygoid proceſs* is ſomewhat triangular, having a broad baſe, and ending ſmaller above. The back-part of this proceſs has three foſſæ formed in it; the two lateral receive the ends of the two plates of the ſphenoid bone, that are commonly compared to a bat's wing; the middle foſſa makes up a part of what is commonly called the *foſſa pterygoidea;* the fore-ſide of this palatine pterygoid proceſs is an irregular concave, where it receives the back-part of the great tuberoſity of the maxillary bone. Frequently ſeveral ſmall holes may be obſerved in this triangular proceſs, particularly one near the middle of its baſe, which a little above communicates with the common and proper holes of this bone already mentioned.

The *naſal lamella* of this bone is extremely thin and brittle, and riſes upwards from the upper ſide of the external edge of the ſquare bone, and from the narrow extremity of the pterygoid proceſs; where it is ſo weak, and at the ſame time ſo firmly fixed to the maxillary bone, as to be very liable to be broken in ſeparating the bones. From the part where the plate riſes, it runs up broad on the inſide of the tuberoſity of the maxillary bone, to form a conſiderable ſhare of the ſides of the maxillary ſinus, and to cloſe up the ſpace between the ſphenoid and the great bulge of the maxillary bone, where there would otherwiſe be a large ſlit opening into the noſtrils.

noſtrils (*g*). From the middle internal ſide of this thin plate, a croſs ridge, placed on ſuch another of the maxillary bone, is extended; on it the back-part of the os ſpongioſum inferius reſts. Along the outſide of this plate, the perpendicular foſſa made by the palate-nerve is obſervable.

At the upper part of this naſal plate, the palate bone divides into *two proceſſes*, which I have already named *orbitar*;—between which and the body of the ſphenoid bone, that hole is formed which I mentioned as the laſt of the holes common to the ſphenoid bone. Sometimes this hole is wholly formed in the os palati, by a croſs plate going from the one orbitar proceſs to the other. A nerve, artery, and vein, belonging to the noſtrils, paſs here.—The anterior of the two orbitar proceſſes is the largeſt, and has its fore-part contiguous to the back-part of the maxillary ſinus, and its upper ſurface appears in the bottom of the orbit, behind the back-part of the os maxillare and planum. It has cells behind, reſembling thoſe of the ethmoid bone, to which it is contiguous; it is placed on the aperture of the ſinus ſphenoidalis, ſo as to leave only a round hole at its upper fore-part.—The other part of the orbitar proceſs is extended along the internal ſide of the upper back-part of the maxillary tuberoſity, the baſe of the ſphenoid bone, between the root of the proceſſus azygos and the pterygoid proceſs.

The palate ſquare part of this plate-bone, and its pterygoid proceſs, are firm and ſtrong, with ſome cancelli; but the naſal plate and orbitar proceſſes are very thin and brittle.

The palate-bones are *joined* to the maxillary, by the fore-edge of the palate ſquare bone, by the tranſverſe palate ſuture:——By their thin naſal plates, and part of their orbitar proceſſes, to the ſame bones, by the palato-maxillares ſutures:——By their pterygoid proceſſes, and back-part of the naſal plates, to the alæ veſpertilionum, by the ſphe-

noid

(*g*) Albin. de oſſib. §. 88.

noid ſuture:——By the tranſverſe ridges of the naſal palates, to the oſſa ſpongioſa inferiora, by contact; hence there frequently is an intimate union of the ſubſtance of theſe bones in old ſkulls:——By the orbitar proceſſes, to the oſſa plana and cellulae ethmoidae, by the ethmoid ſuture:——To the body of the ſphenoid bone, by the ſphenoid ſuture:—By the internal edge of the ſquare bones, to each other by the longitudinal palate ſuture;—and by their naſal ſpines, to the vomer, by the ſpinous ſuture.

The palate bones form part of the palate, noſtrils, orbits, and foſſae pterygoideae; and they cover part of the ſinus maxillares, ſphenoidales, and ethmoidei.

Theſe bones are very complete in a new-born infant, the naſal plates being then thicker and ſtronger than in adults; but the orbitar proceſſes have not the cells which appear in the bones of adults.

When we are acquainted with the hiſtory of theſe bones, the reaſon is evident, why the eyes are ſo much affected in ulcers of the palate, as to be often attended with blindneſs, which frequently happens in an ill-managed lues venerea; or why, on the other hand, the palate ſuffers from an aegylops (*b*).

OSSA TURBINATA.

Ossa Turbinata, or *ſpongioſa inferiora*, reſemble the ſuperior oſſa ſpongioſa in ſhape and ſubſtance, but have their anterior and upper edges contiguous to the tranſverſe ridges of the naſal proceſſes of the maxillary and palate-bones.——From their upper ſtraight edge, two ſmall proceſſes ſtand out: The poſterior, which is the broadeſt, deſcends to cover ſome of the antrum Highmorianum; the anterior riſes up to join the os unguis, and to make part of the lacrymal duct.

Below

(*b*) Hoffman. in Ephemerid. Germ. cent. 1. and 2. obſ. 135.

Below the ſpongy bones already mentioned, there are ſometimes two others, one in each noſtril, which ſeem to be a production of the ſides of the maxillary ſinus turned downwards (*i*). When this third ſort of ſpongy bones is found, the middle one of the three in each noſtril is the largeſt, and the loweſt is the ſmalleſt.——Beſides all theſe, there are often ſeveral other ſmall bones ſtanding out into the noſtrils, that, from their ſhape, might alſo deſerve the name of *turbinata:* but are uncertain in their bulk, ſituation, and number (*k*).

The names of theſe bones ſufficiently declare their ſpongy ſubſtance, which has no firm external plate covering it.

They are joined to the offa maxillaria, palati, and unguis, in all ſubjects, by a firm union of ſubſtance; and as this often happens in people of no great age, ſome authors (*l*) are of opinion, that they ſhould be eſteemed part of the palate-bones; others (*m*) think, that ſince their upper edge is continued by a plate to a part of the os ethmoides, they ought to be eſteemed a part of this bone.

Their uſe is, to ſtraighten the noſtrils, to afford a large ſurface for extending the organ of ſmelling, to cover part of the antra maxillaria, and to aſſiſt in forming the under part of the lacrymal ducts, the orifices of which into the noſe are concealed by theſe bones.

The offa turbinata are nearly complete in a new born infant.

VOMER.

Vomer, or bone reſembling a plough-ſhare, is the thirteenth of the upper jaw, and is without a fellow, forming the lower and back-parts of the partition of the noſe (*n*).

The

(*i*) Cowper in Drake's Anthropolog. book 3. chap. 10.

(*k*) Santorin. Obſervat. Anatomicae, cap. 5. ſect. 9.

(*l*) Id. ib. cap. 5. ſect. 7.

(*m*) Hunauld, in Memoires de l'acad. des ſciences, 1730.

(*n*) Columb. de re anat. lib. 1. cap. 8.——Fallop. Obſervat. Anatom.

The figure of this bone is an irregular rhomboid. Its ſides are flat and ſmooth. Its poſterior edge appears in an oblique direction at the back part of the noſtrils. The upper one is firmly united to the baſe of the ſphenoid bone, and to the naſal plate of the ethmoid; and, when it can be got ſeparated, is hollow for receiving the proceſſus azygos of the ſphenoid. The anterior edge has a long furrow in it, where the middle cartilage of the noſe enters. The lower edge is firmly united to the naſal ſpines of the maxillary and palate bone. Theſe edges of this bone are much thicker than its middle, which is as thin as the thinneſt paper; by which, and the firm union, or connection this bone has above and below, it can very ſeldom be ſeparated entire in adults: but in a child it is much more eaſily ſeparated entire, and its ſtructure is more diſtinctly ſeen; wherefore I ſhall examine all its parts in ſuch a ſubject.

Its ſituation is not always perpendicular, but often inclined and bended to one ſide, as well as the naſal plate of the ethmoid bone.

The vomer is convex at its upper part; and then is ſtraight, as it is extended downwards and forwards, where it is compoſed of two plates: the edges of which have a great number of ſmall proceſſes, diſpoſed ſomewhat like the teeth of a ſaw, but more irregularly, and ſeveral of them are reflected back. Between theſe plates a deep foſſa is left, which, as far as to the top of the curvature, is wide and has ſtrong ſides, for receiving the proceſſus azygos of the ſphenoid bone. Beyond the arch forwards, the foſſa is gradually narrower and ſhallower to the point of the bone, receiving for ſome way the naſal lamella ethmoidea; which, after the oſſification is complete, is ſo cloſely united to the vomer by the little proceſſes piercing its ſubſtance, as to prevent any ſeparation: on which account it has been eſteemed by ſome authors (o) a part of the ethmoid bone.

(o) Lieutaud, Eſſais anatomiques, ſect. 1. l'os ethmoide.

bone. The middle cartilage of the nose fills up what remains of the fossa at its fore-part. The posterior edge of the vomer, which appears above the back-part of the palate bones, is broader above; but as it descends forwards, becomes thinner, though it is still solid and firm. The lower edge of this bone, which rests on the nasal spine of the palate and maxillary bones, has a little furrow on each side of a small middle ridge, answering to the spines of the bones of different sides, and the interstices between them. This and the upper edge meet in the pointed fore-end of this bone.

The body of the vomer has a smooth and solid surface, but thin substance; and towards its sides, where it is thickest, some cancelli may be observed when the bone is broken.

It is joined above to the sphenoid and ethmoid bones, and to the middle cartilage of the nose by schindylesis: below, to the maxillary and palate bones by the spinous future.

The vomer divides the nostrils; enlarges the organ of smelling, by allowing place for expanding the membrane of the nose on its sides; and sustains the palate-plates of the maxillary and palate-bones, which otherwise might be in hazard of being pressed into the nostrils: while the vomer is secured from shuffling to one side or other by the double schindylesis, by which it is joined to the bones above and below.

Thus we have now described all the bones which compose the upper jaw, except the teeth, which are so much like those of the lower jaw, that I choose to make one description serve for both, in which their differences shall be remarked, after the second part of the face, (*i. e.*) the lower jaw, is examined; because the structure of the teeth

teeth cannot be well underſtood, until the caſe in which they are ſet is explained.

MAXILLA INFERIOR.

MAXILLA INFERIOR (*p*), the lower jaw, conſiſts only of one moveable bone, and ſixteen teeth incaſed in it.

This bone, which is ſomewhat of the figure of the Greek letter υ, is ſituated at the lower part of the face, ſo as its convex middle part is forward, and its legs are ſtretched back. It is commonly divided into the chin, ſides, and proceſſes.—The *chin* is the middle fore-part; the extent of which to each ſide is marked on the external ſurface by the holes obſervable there, and internally by the beginning of an oblique ridge. Beyond theſe the *ſides* appear; and are continued till the bone, by bending upwards, begins to form the *proceſſes*.

On the fore-part of the *chin*, a tranſverſe ridge appears in the middle; on each ſide of which the muſculi quadrati, or depreſſores, and the levatores labii inferioris, depreſs the bone: And below theſe prints a ſmall riſing may be obſerved, where the depreſſores commence. On the back-part of the chin, ſometimes three, always two, ſmall protuberances appear in the middle. To the uppermoſt, when it is ſeen, the frænum of the tongue is connected. From the middle one, the muſculi genio-gloſſi riſe; and from the loweſt, the genio-hyoidei have their origin. Below the laſt, we ſee two rough ſinuoſities formed by the digaſtric muſcles.

At the lower and fore-part of the external ſurface of each *ſide* of the lower jaw a ſmall eminence may be obſerved, where the depreſſor labiorum communis riſes. Near the upper edge of the ſide a ridge runs lengthwiſe, to which the

(*p*) Γενυς σιαγων, Mandibula, facies.

under-part of the muſculus buccinator is connected.—Internally, towards the upper edge of each ſide another ridge appears, from which the mylo-hyoidei have their origin, and to which the internal membrane of the gums adheres.

In the upper edge of both *chin* and *ſides*, are a great many deep pits or ſockets, for receiving the roots of the teeth. The number and magnitude of theſe ſockets are various, becauſe of the different number, as well of the teeth themſelves, as of their roots, in different people. Theſe ſockets in this lower jaw, as well as in the upper one, are leſs deep as old age comes on: When freed from the teeth by any means, they are ſome time after filled up with an oſſeous net-work, which at laſt becomes entirely ſolid, and as ſmooth as any other part of the bone; ſo that, in a great many old jaws, we cannot obſerve a veſtige of the ſockets: But then the jaw becomes leſs, and much narrower (*q*). Hence we may know why the chin and noſe of edentulous people are much nearer than before the teeth were loſt; while their lips either fall in towards their mouth, or ſtand prominent forwards.——When new teeth are protruded, new ſockets are formed (*r*). The lower edge of the chin and ſides is ſmooth and equal, and is commonly called the baſe of the lower jaw. The ends of the baſe, where the jaw turns upwards, are called its angles; the external ſurface of each of which has ſeveral inequalities upon it, where the maſſeter muſcle is inſerted; as the internal ſurface alſo has, where the pterygoideus internus is inſerted, and a ligament extended from the ſtyloid proceſs of the temporal bone is fixed.

The *proceſſes* are two on each ſide.——The anterior ſharp thin coronoid ones have the crotaphite muſcles inſerted into them. The poſterior proceſſes or condyles (*s*) terminate in an

(*q*) Veſal. Anat. lib. 1. cap. 10.

(*r*) Fallop. Obſerv. Anat.

(*s*) Articulatorii.

an oblong ſmooth head, ſupported by a cervix. The heads, whoſe greateſt length is tranſverſe, and whoſe convexity is turned forwards, are tipped with a cartilage, as the articulated parts of all other moveable bones are. The fore-part of the root and neck of theſe condyloid proceſſes are a little hollow and rough where the external pterygoid muſcles are inſerted.

The *holes* of the lower jaw are two on each ſide; one at the root of the proceſſes internally, where a large branch of the third branch of the fifth pair of nerves enters with an artery, and a vein returns. A ſmall ſharp proceſs frequently juts out backwards from the edge at the fore-part of this hole, to which a ligament extended from the temporal bone is fixed (*t*) which ſaves the nerve and veſſels from being too much preſſed by the pterygoid muſcles.—From the lower ſide of this hole, either a ſmall ſuperficial canal or furrow deſcends, where a branch of the nerve is lodged, in its way to the mylo-hyoideus muſcle and ſublingual gland (*u*). The other hole is external, at the confines of the chin, where branches of the nerve and veſſels come out. The canal between theſe two holes is formed in the middle of the ſubſtance of the bone, and is pierced by a great number of ſmall holes, by which the nerves and blood-veſſels of the cancelli and teeth paſs. This canal is continued a little farther than the external hole at the chin.—On account of the veſſels and nerves in the lower jaw, fractures of it may be attended with dangerous ſymptoms.

The *ſurface* of the lower jaw is hard and firm, except at the ſpongy ſockets, where, however, it is ſtronger than the upper jaw. Its internal *ſubſtance* is cellular, without any ſolid partition between the cancelli in its middle. At the baſe, eſpecially of the chin, where this bone is moſt expoſed

(*t*) Weitbrecht. Syndeſmolog. fig. 32. 1.

(*u*) Palfyn. anat. chirur. traite 5. chap. 6.

ed to injuries, the ſolid ſides of it are thick, compact, and hard.

The lower jaw generally receives the root of ſixteen teeth into its ſockets, by *gomphoſis;* and its condyloid proceſſes covered with a cartilage, are articulated with the temporal bones, in a manner that is not commonly right deſcribed: For, as was already mentioned in the deſcription of the temporal bones, not only the fore-part of the cavity between the zygomatic, auditory, and vaginal proceſſes, but alſo the adjoining tubercle at the root of the zygomatic proceſs of each os temporum, is covered with a ſmooth cartilage for this articulation. Here alſo an intermediate moveable cartilage is placed; which being thin in the middle, and thick at the edges, is concave on both ſides; and is connected ſo firmly by ligaments to each condyle, as to follow the motions of the condyle; and ſo looſely to the temporal bone, as readily to change its ſituation from the cavity to the tubercle, and to return again; while the common ligament of the articulation affords ſpace enough for ſuch a change of place backwards and forwards: But like other ligaments of the joints by ginglimus, is ſtrong and ſhort at the ſides, to confine the lateral motions.

When, therefore, the teeth of both jaws coincide, the condyles are lodged ſecurely in the temporal cavities; but their motions to either ſides muſt be confined both by the firmneſs of the ligaments and the riſing brims which are on each ſide of the cavities.—When the jaw is brought directly forwards, the condyle and intermediate cartilages deſcend and advance forwards upon the tubercles.—In this ſituation, the lateral motions are a little more free than in the former one, from the want of riſing brims to ſtop the condyles.—When the fore-teeth of the lower jaw are moved forwards and to a ſide, the condyle of the oppoſite ſide is either advanced from the cavity to the tubercle, while

while the condyle of the ſame ſide remains in the cavity; or, if both condyles are on the tubercles, when the jaw is moved obliquely to a ſide, the condyle of the ſide to which the motion is made ſlides back from the tubercle to the cavity. —— When the mouth is opened by the deſcent of the lower jaw, the fore-part of it, where the depreſſing muſcles are fixed, is drawn backwards, as well as downwards, while reſiſtance is made to the angles moving backwards by the maſſeter and internal pterygoid muſcles, and, at the ſame time, the external pterygoid draws the condyles and their moveable cartilages forwards; and therefore, when the mouth is opened, the condyles are carried forwards upon the tubercles, and the axis of motion of the bone is a little above its angles. But in this ſituation there is leſs reſiſtance, than in any other, to the condyles luxating forwards; a diſeaſe which ſeldom happens, except when people are gaping too wide; and therefore the common practice of nurſes, who ſupport the jaw of infants when they are yawning, is reaſonable. ——In chewing, there is a ſucceſſion of the motions above deſcribed (*x*).

Here a general remark may be made, that wherever moveable cartilages are found in joints, either the articulated bones are of ſuch a figure, or ſo joined and fixed by their ligaments, that little motion would be allowed without ſuch cartilages; or elſe ſome motions are neceſſary to the right uſe of the member, which the form of the articulation would not otherwiſe admit of. This will more fully appear after the other joints with ſuch cartilages are deſcribed.

In a child born to the full time, the lower jaw is compoſed of two bones, connected by a thin cartilage in the middle of the chin, which gradually oſſifies, and the two bones

(*x*) For a more full account of this articulation, vid. Edinburgh Medical Eſſays and Obſerv. vol. I. art. II. and vol. iii. art. 13.—Memoires de l'acad. des ſciences, 1744.

bones intimately unite.——In each of theſe bones there are five or ſix ſockets for teeth, as in the upper jaw.

After I have thus deſcribed the incaſement of the teeth; the inſertion of ſo many muſcles of the tongue, and of the os hyoides; the connection of the membrane of the tongue to the maxillary bone, and the motions of this bone; it is eaſy to ſee, that the lower jaw muſt be a principal inſtrument in manducation, deglutition, and ſpeech.

THE TEETH.

The teeth are the hard white bodies placed in the ſockets of both jaws. Their number is generally ſixteen above and as many below; though ſome people have more, others have fewer.

The broad thick part of each tooth which appears without the ſocket, is the baſe or body (*y*).——The ſmaller proceſſes ſunk into the maxillae, are the roots or fangs; which become gradually ſmaller towards the end fartheſt from the baſe, or are nearly conical; by which the ſurface of their ſides divides the preſſure made on the baſes, to prevent the ſoft parts, which are at the ſmall points of the ſockets, to be hurt by ſuch preſſure. At the place where the baſe ends and the roots begin, there is generally a ſmall circular depreſſion, which ſome call the neck or collar.

Without the gums the teeth are covered with no membrane, and they are ſaid to have no proper perioſteum within the ſockets; but that is ſupplied by the reflected membrane of the gums, which after a good injection may be evidently ſeen in a young ſubject, with the veſſels from it penetrating into the ſubſtance of the teeth; and it may be diſcovered in any tooth recently pulled, by macerating it in water (*z*). The adheſion of this membrane to theſe

(*y*) Corona. (*z*) Cowper's Anat. Explic. tab. 27. fig. 9. lit. E.

theſe roots is ſtrengthened by the ſmall furrows obſervable on them.

Each tooth is compoſed of its *cortex* or enamel, and an internal bony ſubſtance. The *cortex* has no cavity or place for marrow; and is ſo ſolid and hard, that ſaws or files can with difficulty make impreſſion on it. It is thickeſt upon the baſe, and gradually, as the roots turn ſmaller, becomes thinner, but not proportionally to the difference of the ſize of the baſe and roots.——The fibres of this enamel are all perpendicular to the internal ſubſtance; and are ſtraight on the baſe, but at the ſides are arched with a convex part towards the roots (*a*); which makes the teeth reſiſt the compreſſion of any hard body between the jaws, with leſs danger of breaking theſe fibres, than if they had been ſituated tranſverſely. The ſpongy ſockets in which the teeth are placed, likewiſe ſerve better to prevent ſuch an injury than a more ſolid baſe would have done.——Notwithſtanding the great hardneſs of this cortex, it is waſted by manducation. Hence the ſharp edges of ſome teeth are blunted and made broad, while the rough ſurfaces of others are made ſmooth and flat, as people advance in life.

The bony part of the teeth has its fibres running ſtraight, according to the length of the teeth. When it is expoſed to the air, by the breaking or falling off of the hard cortex, it ſoon corrupts. And thence carious teeth are often all hollow within, when a very ſmall hole appears only externally.

The teeth have canals formed in their middle, wherein their nerves and blood-veſſels are lodged: which they certainly need, being conſtantly waſted by the attrition they are ſubjected to in manducation; and for their further growth, not only after they firſt appear, but even in adults; as is evident when a tooth is taken out: for then the oppoſite

(*a*) Haver's Oſteolog. Nov. diſc. 1.

ſite one becomes longer, and thoſe on each ſide of the empty ſocket turn broader; ſo that when the jaws are brought together, it is ſcarcely obſervable where the tooth is wanting (*b*).

The veſſels are eaſily traced as long as they are in the large canal, but can ſcarcely be obſerved in their diſtribution from that to the ſubſtance of the teeth of adults. Ruyſch (*c*) however affirms, that after injection he could trace the arteries into the hardeſt part of the teeth: And Leeuwenhoek (*d*) ſuſpected the fibres of the cortex to be veſſels. In children I have frequently injected the veſſels of the teeth as far as their baſe; and in ſuch as are not entirely oſſified, one can with a lucky injection fill ſo many veſſels as to make both the outſide and inſide of the cortical part appear perfectly red.—This plentiful ſupply of veſſels muſt expoſe the teeth to the ſame diſorders that attack other vaſcular parts; and ſuch teeth as have the greateſt number of veſſels muſt have the moſt numerous chances of being ſeized with theſe diſeaſes.

Every root of each tooth has a diſtinct canal, with veſſels and nerves in it. Theſe canals in the teeth with more than one root, come nearer each other as they approach the baſe of the tooth; and at laſt are only ſeparated by very thin plates, which, being generally incomplete, allow a communication of all the canals; and frequently one common cavity only appears within the baſe, in which a pulpy ſubſtance compoſed of nerves and veſſels is lodged.——The condition therefore of the nerves here bears a ſtrong analogy to that of the cutaneous nerves which ſerve for the ſenſation of touching.

The entry of the canals for theſe veſſels is a ſmall hole placed

(*b*) Ingraſ. de tumor. cap. 1. p. 24. 25. 26.

(*c*) Theſaur. 10. num. 27.

(*d*) Arcan. Natur. continuat. epiſt. p. 3.

placed a little to a ſide of the extreme point of each root; ſometimes, eſpecially in old people, this hole is entirely cloſed up, and conſequently the nerves and blood-veſſels are deſtroyed (*e*).

The teeth are ſeen for a conſiderable time in form of mucus contained in a membrane; afterwards a thin cortical plate and ſome few oſſeous layers appear within the membrane, with a large cavity filled with mucus in the middle; and gradually this exterior ſhell turns thicker, the cavity decreaſes, the quantity of mucus is leſſened, and this induration proceeds till all the body is formed, from which the roots are afterwards produced.

In young ſubjects, different ſtamina or rudiments of teeth are to be obſerved. Thoſe next the gums ordinarily hinder the deeper ſeated ones from making their way out, while theſe prevent the former from ſending out roots, or from entering deep into the bony ſockets of the jaws; by which they come to be leſs fixed.

Children are ſeldom born with teeth; but at two years of age they have twenty; and their number does not increaſe till they are about ſeven years old; when the teeth that firſt made their way through the gums are thruſt out by others that have been formed deeper in the jaw, and ſome more of the teeth begin to diſcover themſelves farther back in the mouth. About fourteen years of age, ſome more of the firſt crop are ſhed, and the number is increaſed. This ſhedding of the teeth is of good uſe: for if the firſt had remained, they would have ſtood at a great diſtance one from another; becauſe the teeth are too hard in their outer cruſt to increaſe ſo faſt as the jaws do. Whereas, both the ſecond layer and the teeth that come out late, meeting, while they are ſoft, with a conſiderable reſiſtance to their growth in length, from thoſe ſituated upon them, neceſſarily come out broad,

(*e*) De la Hire, Hiſtoire de l'acad. des ſciences, 1699.

broad, and fit to make that close guard to the mouth (*f*) which they now form.

The teeth are *joined* to the sockets by *gomphosis*, and the gums contribute to fix them there; as is evident by the teeth falling out when the gums are any way destroyed or made too spongy, as in the scurvy or salivation: whence some authors (*g*) class this articulation with the *sysarcosis*.

The *uses* of the teeth are to masticate our aliment, and to assist us in the pronunciation of several letters.

Though the teeth so far agree in their structure, yet, because of some things wherein they differ, they are generally divided into three classes, viz. *incisores*, *canini*, and *molares*.

The INCISORES (*h*) are the four fore-teeth in each jaw, receiving their name from their office of cutting our aliment; for which they are excellently adapted, being each formed into a sharp-cutting edge at their base, by their fore-side turning inwards there, while they are sloped down and hollowed behind (*i*); so that they have the form of wedges, and therefore their power of acting must be considerably increased.——Seeing, in the action of the incisores, a perpendicular compression is only necessary, without any lateral motion, they are not so firmly fixed in their sockets as the other teeth are, each having only one short root; but that is broader from before backwards, than to either side, to have the greatest strength where it is exposed to the strongest force applied to it (*k*).

The incisores of the upper jaw, especially the two middle ones (*l*), are generally broader and longer than those of the upper jaw.

In

(*f*) Φραγμος. (*g*) Drake's Anthropolog. book. 4. chap. 3.

(*h*) Τελαστινοι, τομικοι, διλαστηρες, κτενες, τομεις, πρωσθιοι, οξεις, Riforii, quaterni, primi, primores, anteriores, acuti.

(*i*) Ολμισκος.

(*k*) Lettre sur l'osteologie, ascribed to Du Verney. (*l*) Duales,

In a new-born infant, only the outer ſhell of the body of theſe teeth is hardened.—Afterwards, when the ſtamina of two ſets are formed, each has its own ſocket, thoſe neareſt to the edge of the gums being placed more forward, and the others are lodged farther back within the jaw-bones.

CANINI (*m*), from the reſemblance to dog tuſks, are one on each ſide of the inciſores in each jaw.——The two in the upper jaw are called *eye-teeth*, from the communication of nerves which is ſaid to be between them and the eyes.—— The two in the lower jaw are named *angular* or *wike-teeth*, becauſe they ſupport the angles of the mouth.

The canini are broader, longer, and ſtronger, than the inciſores.——Their baſes are formed into a ſharp-edge, as the inciſores are; only that the edge riſes into a point at the middle. Each of them has generally but one long root, though ſometimes they have two (*n*). The roots are crooked towards the end. The canini of the upper jaw are larger, longer, and with more crooked roots, than thoſe of the under jaw. The form of their baſe is fit both for piercing and cutting, and the long crooked root of each makes it ſecure in the ſocket.

The canini of a child are in much the ſame condition as the inciſores are.

The DENTES MOLARES, or GRINDERS (*o*), which have got their name becauſe they grind our food, are generally five in each ſide of each jaw; in all twenty. Their baſes are broader, more ſcabrous, and with a thinner cortical ſubſtance, than the other teeth. They have alſo more roots; and as theſe roots generally divaricate from each other, the partitions of the

(*m*) Κυνοδοντες, Riſorii, fractorii, collaterales, columellares.

(*n*) Fauchard, Chirurgien dentiſte, chap. I.

(*o*) Μυλιται, γομφοι, μυλοι, πλατεις, φραξηρες, Maxillares, menſales, clavales, buccarum.

the ſockets between them bear a large ſhare of the great preſſure they ſuffer, and hinder it from acting on their points (*p*).

The baſe of the firſt grinder has an edge pointed in the middle, on its outſide, reſembling the canini; from which it ſlopes inwards till it riſes again into a point.—It has generally but one root, which ſometimes is long and crooked at its point.

The ſecond dens molaris has two points on its baſe, riſing nearly equally on its out and inſide.—It has two roots either ſeparate or run together, but ſhorter than the root of the firſt. Theſe two anterior grinders are much ſmaller than the three that are placed farther back in the mouth.

The third and fourth are very broad in their baſes, with four or five points ſtanding out; and they have three or more roots.

The fifth, commonly called *dens ſapientiæ* (*q*), from its coming through the gums later than the other grinders, has four points on its baſe, which is not ſo large as the baſe of the third and fourth, and its roots are leſs numerous.

The inciſores of the upper jaw being broader than thoſe of the lower jaw, make the ſuperior grinders to be placed ſo much farther back than the lower ones, that, when they are brought together, by ſhutting the mouth, the points of the grinders of the one jaw enter into the depreſſions of the oppoſite grinders, and they are all equally applied to each other, notwithſtanding the inequality of their ſurface.

The numerous roots of the dentes molares prevent their looſening by the lateral preſſure they ſuffer in grinding; and as the ſockets in the upper jaw are more ſpongy, and the teeth are

(*p*) Lettre ſur l'oſteologie.

(*q*) Σωφρονιστηρες, φαντηρες, οψιγονοι, Senſus, intellectus, ſerotini, ætatem complentes, genuini, moderatores.

are more liable, by their ſituation, to fall out (*r*), the grinders there have more numerous and more ſeparated roots than in the lower jaw (*s*). The number, however, of the roots of the teeth of both jaws is very uncertain; ſometimes they are more, ſometimes fewer; frequently ſeveral roots are joined together; at other times they are all diſtinct. The diſpoſition of ſuch as are diſtinct is alſo various; for in ſome the roots ſtand out ſtraight, in others they ſeparate, and in others again they are crooked inwards. When the roots are united, we can ſtill diſtinguiſh them, by remarking the number of ſmall holes at their points, which determine the number of roots each tooth ought to be reckoned to have.

At the time of birth, only two dentes molares in each jaw have begun to oſſify; and that at little more than the baſe, which has ſeveral ſharp points ſtanding out from it. The temporaneous grinders are placed more directly upon the internal ſet than the other two claſſes are: Sometimes there is a piece of the bone of the jaws between the two ſets; in other children, the two ſets have no bone interpoſed between them.

From what has been ſaid, the anſwers to the following queries may be given.

Why are children ſubject to ſalivation, fever, convulſion, vomiting, purging, &c. when their teeth are breeding or cutting the gums?

Why in children do the dentes inciſores firſt cut the gums, the canini next, and molares laſt?

Why do children ſhed their teeth?

Why have theſe temporaneous teeth generally no roots, or very ſmall ones?

Why have theſe firſt teeth ſometimes roots, and that more frequently

(*r*) Galen de oſſib. cap. 5.

(*s*) Fauchard. Chirurg. dent. chap. 3.

frequently in teeth pulled by art than in thoſe which are ſhed by nature (*t*) ?

Why do theſe roots frequently come outwards through the gums ?

Whence comes butter or buck teeth ?

How do theſe teeth ſometimes go into the natural row with the others, after pulling a rotten tooth near them ?

How have ſome people got two rows of teeth in one or both jaws (*u*) ?

Why do the teeth of old people looſen, and then drop out entire ?

Whence ariſe the new ſets of teeth which ſeveral old people obtain (*x*) ?

Why are not the gums of toothleſs old people torn by the hard ſockets in chewing ?

Why are the teeth inſenſible when ſlightly filed or raſped ?

How come they to be ſenſible of heat or cold, to be ſet on edge by acids, or to give an uneaſy ſenſation when gritty or ſandy ſubſtances are rubbed between them ?

Why does a perſon who has a pained tooth imagine it longer than any other ?

What is the reaſon of ſome perſons dying convulſed, upon raſping or filing down an overgrown tooth (*y*) ?

How do the teeth break and moulder away without any pain in ſome people, and not in others ?

What parts are affected in the toothach ?

What are the cauſes of the toothach ?

May worms be reckoned among theſe cauſes (*z*) ?

Why are the dentes molares moſt ſubject to that diſeaſe ?

In

(*t*) Fauchard, Chirurgien-dentiſte, p. 7.

(*u*) Blaſ. Comment. ad Veſling. Syntagm. cap. 1. 3.

(*x*) Hoffman. in Van. Horc. Microcoſm. p. 38.

(*y*) Bartholin. Anat. reformat. lib. 4. cap. 12.

(*z*) Jacob. in Act. Hafn. vol. 5. obſ. 107.—Pechlin. Obſerv. Medic. lib. 2. obſ. 36.—Bartholin. Hiſt. Medic. cent. 3. hiſt. 96.

In what different manners ought the ſeveral claſſes of teeth to be extracted when ſuch an operation is neceſſary?

Whence proceeds the violent obſtinate hemorrhagy which ſometimes attends the drawing of teeth (*a*)?

Why is it more difficult and dangerous to draw the eye-teeth than any other?

What makes it impoſſible frequently to draw grinders without bringing away part of the jaw-bone with them, or breaking the fangs?

Why do teeth ſoon replaced after being extracted become again fixed in the ſockets (*b*)?

According to the diviſion made of the ſkeleton, we ſhould now proceed to the deſcription of the *trunk* of the body. But we muſt firſt conſider a bone which cannot well be ſaid to belong to either the head or the trunk; nor is it immediately joined to any other, and therefore is very ſeldom preſerved with ſkeletons. However, it is generally deſcribed by authors after the bones of the face. In obedience, therefore, to the prevailing method, I ſhall next examine the ſtructure of the

OS HYOIDES.

The os hyoides (*c*), is ſituated horizontally between the root of the tongue and the larynx. It is properly enough named *hyoides*, from the reſemblance it bears to the Greek letter *υ*; and may, for a clearer demonſtration of its ſtructure, be diſtinguiſhed into its *body*, *cornua*, and *appendices*.

The

(*a*) Paré, livre 6. chap. 2.——Rolfinc, lib. 2. cap. 27. & 30.——Moebii Fundam. Medicin. cap. 9.——Ephemerid. German. dec. 1. ann. 3. obſ. 319.——Fauchard, Chirurg.-dentiſte, tom. 1. chap. 23. obſerv. 7.

(*b*) De la Motte Chirurgie, tom. 1. chap. 4. obſ. 2.—Fauchard, Chirurgien-dentiſte. tom. 1. chap. 29.

(*c*) Hypſyloides, lambdoides, παραστατη φαρυγγιτερον, Os gutturis, os linguae, os morſus Adami, aſſeſſor, os laude, bicorne.

The *body* is the middle broad part, convex before and hollow behind.——The convex fore-part is divided into two by a ridge, into the middle of which the mylo-hyoidei, and into the ſides the ſtylo-hyoidei, muſcles are inſerted.——Above the ridge, the bone is horizontal; but pitted in the middle by the inſertion of the two genio-hyodei muſcles, and a little hollowed more laterally by the baſio-gloſſi.——Below the ridge, it is convex; but a little flatted in the middle by the ſterno-hyoidei, and pitted more externally by the coraco-hyoidei.——The concavity behind faces backwards and downwards to receive the thyroid cartilage, when the larynx and the os hyoides are pulled towards each other by the action of the ſterno-hyoidei and hyo-thyroidei muſcles; and to its upper edge, the ligamentous membranes of the epiglottis, tongue, and thyroid cartilage, are fixed.

The *cornua* of the (*d*) os hyoides are ſtretched backwards from each ſide of its body, where often a ſmall furrow points out the former ſeparation; for in young ſubjects, the body and cornua are not one continued ſubſtance, as they come afterwards to be in adults. Theſe cornua are not always ſtraight, nor of an equal length; their two plain ſurfaces ſtand obliquely ſloping from above outwards and downwards.——Into the external, the cerato-gloſſis is inſerted above, and the thyro-hoideus muſcle below; and to the one behind, the ligamentous membrane of the tongue and larynx adheres.—Each of the cornua becomes gradually ſmaller as it is extended from the baſe; but ends in a round tubercle, from which a moveable cartilage ſtands out, which is connected to the upper proceſs of the cartilago thyroidea.

Where the body of the os hyoides joins on each ſide with its cornua, a ſmall ſtyliform proceſs, called *appendix* (*e*), riſes

(*d*) Crura, latera, inferiora.

(*e*) Crura ſuperiora, latera ſuperiora, oſſa graniformia.

riſes upwards and backwards, into which the muſculi ſtylo-hyoidei alteri, and part of the hyo-gloſſi muſcles, are fixed. From each of them a ligament is ſometimes extended to the ſtyloid proceſſes of the temporal bones, to keep the os hyoides from being drawn too much forwards or downwards. The part of this ligament next to theſe proceſſes ſometimes forms into ſeveral cartilages, which afterwards oſſify in old people. Ruyſch *(f)* ſays, that he has ſeen this oſſification continued as far up as the ſtyloid proceſſes, which were therefore joined to the os hyoides by anchyloſis.

The *ſubſtance* of the os hyoides is cellular; but covered with a firm external plate, which is of ſufficient ſtrength to bear the actions of ſo many muſcles as are inſerted into it.

It is not *articulated* with any bone of the body, except by means of the muſcles and ligaments already mentioned.

The *uſe* of the os hyoides, is to ſerve as a ſolid lever for the muſcles to act with, in raiſing or depreſſing the tongue and larynx, or in enlarging and diminiſhing the capacity of the fauces.

At birth, this bone is in a cartilaginous ſtate; excepting a ſmall point of bone in the middle of its body, and in each of the cornua.—The appendices frequently remain cartilaginous many years.

Sect. II. Of the TRUNK.

THE Trunk conſiſts of the *ſpine*, *pelvis*, and *thorax*.

§ 1. *The* Spine.

The ſpine (g) is the long piles of bones extended from the condyles of the occiput to the end of the rump. It ſome-

(*f*) Adverſ. Anat. dec. 3. § 9.

(g) Ραχις νωτου ακανθα, ιερα συριγξ, σωλην, Tergum hominis carina.

what reſembles two unequal pyramids joined in a common baſe. It is not, however, ſtraight; for its upper part being drawn backwards by ſtrong muſcles, it gradually advances forwards, to ſupport the oeſophagus, veſſels of the head, &c. Then it turns backwards, to make room enough for the heart and lungs. It is next bent forwards, to ſupport the viſcera of the abdomen. It afterwards turns backwards, for the enlargement of the pelvis. And, laſtly, it is reflected forwards, for ſuſtaining the loweſt great gut.

The ſpine is commonly divided into *true* and *falſe vertebrae*; the former conſtituting the long upper pyramid, which has its baſe below; while the falſe vertebrae make the ſhorter lower pyramid, whoſe baſe is above.

TRUE VERTEBRAE.

The true vertebrae (*b*) are the twenty-four upper bones of the ſpine, on which the ſeveral motions of the trunk of our bodies are performed; from which uſe they have juſtly got their name.

Each of theſe vertebrae is compoſed of its body and proceſſes.

The *body* is the thick ſpongy fore-part, which is convex before, concave backwards, horizontal and plain in moſt of them above and below.——Numerous ſmall holes, eſpecially on the fore and back part of their ſurface, give paſſage to their veſſels, and allow the ligaments to enter their ſubſtance. The edges of the body of each vertebra are covered, eſpecially at the fore-part, with a ring of bone firmer and more ſolid than the ſubſtance of the body any where elſe. Theſe rings ſeem to be joined to the vertebrae in the form

(*b*) Στρεφὶς σφόγγος, Spondyli, oſſa orbiculata, oſſa vertebrata, verticula.

form of epiphyſes, but are alleged by ſome authors (*i*) to be the ligaments oſſified. They are of great uſe in preventing the ſpongy bodies from being broken in the motions of the trunk.

Between the bodies of each two adjoining vertebrae, a ſubſtance between the nature of ligament and cartilage is interpoſed; which ſeems to conſiſt of concentrical curved fibres, when it is cut horizontally; but when it is divided perpendicularly, the fibres appear oblique and decuſſating each other (*k*).——The outer part of the invertebral ligaments is the moſt ſolid and hard, and they gradually become ſofter till they are almoſt in the form of a glairy liquor in the centre; and therefore theſe ſubſtances were not improperly called mucous ligaments by the antients (*l*). The external fibrous part of each is capable of being greatly extended, and of being compreſſed into a very ſmall ſpace, while the middle fluid part is incompreſſible, or nearly ſo; and the parts of this ligament between the circumference and centre approach in their properties to each other, in proportion to their more ſolid or more fluid texture. The middle point is therefore a fulcrum or pivot, on which the motion of a ball and ſocket may be made, with ſuch a gradual yielding of the ſubſtance of the ligament, in whatever direction our ſpines are moved, as ſaves the body from violent ſhocks, and their dangerous conſequences (*m*).——This ligamento-cartilaginous ſubſtance is firmly fixed to the horizontal ſurfaces of the bodies of the vertebrae, to connect them; in which it is aſſiſted by a ſtrong membranous ligament, which lines all their concave ſurface, and by a ſtill ſtronger ligament that covers all their anterior convex ſurface.

We

(*i*) Fallop. Obſervat. Anat.

(*k*) Blancardi Anat. reform. cap. 32.—Weitbrecht, Syndeſmolog. ſect. 4. § 15.

(*l*) Galen, De uſu part. lib. 12. cap. 16.

(*m*) Medical Eſſays and Obſerv. vol. 5. art. 28.

We may lay it down as a general rule, notwithſtanding ſome exceptions, That the bodies of the vertebrae are ſmaller, and more ſolid above; but, as they deſcend, they appear larger and more ſpongy; and that the cartilages between them are thick, and the ſurrounding ligaments ſtrong, in proportion to the ſize of the vertebrae, and to the quantity of motion they perform: by which diſpoſition, the greater weight is ſupported on the broadeſt beſt ſecured baſe, and the middle of our body is allowed a large and ſecure motion.

From each ſide of the body of each vertebra, a bony bridge is produced backwards, and to one ſide: from the poſterior end of which one ſlanting *proceſs* riſes, and another deſcends; the ſmooth, and what is generally the flatteſt, ſide of each of theſe four proceſſes, which are called the *oblique* (*n*), is covered with a ſmooth cartilage; and the two lower ones of each vertebra are fitted to, and articulated with, the two upper or aſcending oblique proceſſes of the vertebra below, having their articular ligaments fixed into the rough line round their edges.

From between the oblique proceſſes of each ſide, the vertebra is ſtretched out laterally into a proceſs that is named *tranſverſe*.

From the back-part of the roots of the two oblique and of the tranſverſe proceſs of each ſide, a broad oblique bony plate is extended backwards; where theſe meet, the ſeventh proceſs of the vertebrae takes its riſe, and ſtands out backwards: this being generally ſharp-pointed and narrow edged, it has therefore been called *ſpinal* proceſs; from which this whole chain of bones has got its name.

Beſides the common ligament which lines all the internal ſurface of the ſpinal proceſſes, as well as of the bodies, particular ligaments connect the bony bridges and proceſſes of the contiguous vertebrae together.

The

(*n*) Articulatorii, minimi.

The ſubſtance of the proceſſes is conſiderably ſtronger and firmer, and has a thicker external plate, than the bodies of the vertebrae themſelves.

The ſeven *proceſſes* form a concavity at their fore-part, which, joined to the one at the back-part of the bodies, makes a great hole; and the holes of all the vertebrae form a long large conduit (*o*), for continuing the ſpinal marrow. —In the upper and lower edge of each lateral bridge, there is a notch. Theſe are ſo adapted to each other in the contiguous vertebrae, as to form a round hole in each ſide between each two vertebrae, through which the nerves that proceed from the ſpinal marrow and its blood-veſſels paſs.

The *articulations*, then, of theſe true vertebrae are plainly double: For their bodies are joined by the intervening cartilage above deſcribed; and their oblique proceſſes, being tipped with cartilages, are ſo connected by their ligaments as to allow a ſmall degree of motion on every ſide. Hence it is evident, that their centre of motion is altered in different poſitions of the trunk: For, when we bow forwards, the upper moved part bears entirely on the bodies of the vertebrae; if we bend back, the oblique proceſſes ſupport the weight; if we recline to one ſide, we reſt upon the oblique proceſſes of that ſide and part of the bodies; if we ſtand erect, all the bodies and oblique proceſſes have their ſhare in our ſupport.

Hence it follows, 1. That becauſe the joints of which the ſpine is compoſed are ſo numerous, the ſpinal marrow, nerves, blood-veſſels, &c. are not liable to ſuch compreſſion and overſtretching in the motion of the trunk of the body as they would otherwiſe be, ſince ſeveral vertebrae muſt be concerned in every motion of the ſpine; and therefore a very ſmall curvature is made at the conjunction of any two vertebrae,

(*o*) Ιερα συριγγξ, σωλην, Canalis.

brae (*p*). 2. That an erect poſture is the ſureſt and firmeſt, becauſe the ſurface of contact of the fulcra is largeſt, and the weight is moſt perpendicular to them (*q*). 3. That the muſcles which move the ſpine act with greater force in bringing the trunk into an erect poſture than in drawing it to any other: For in bending forwards, backwards, or to a ſide, the muſcles which perform any of theſe actions are nearer the centre of motion: conſequently the lever with which they act is ſhorter than when the centre of motion is on the part of the vertebra, oppoſite to that where theſe muſcles are inſerted; which is the caſe in raiſing the trunk. This is extremely neceſſary; ſince, in the deflections of the ſpine from a perpendicular bearing, the weight of the body ſoon inclines it in the direction we chooſe; whereas, in raiſing us erect, this great weight muſt be more than counteracted. 4. In calculating the force exerted by the muſcles which move the ſpine, we ſhould always make allowance for the action of the cartilages between the vertebrae, which, in every motion from an erect poſture, muſt be ſtretched on one ſide, and compreſſed on the other, to both which they reſiſt; whereas, in raiſing the trunk, theſe cartilages aſſiſt by their ſpringy force (*r*). 5. We are hence naturally led into the reaſon of our height of ſtature increaſing in the morning, and diminiſhing at night (*s*): For the intermediate cartilages of the vertebrae being preſſed all day long by the weight of our body, become more compact and thin in the evening; but, when they are relieved from this preſſure in the night, they again expand themſelves to their former thickneſs: And ſeeing

(*p*) Galen. de uſu part. lib. 12. cap. 12.

(*q*) Paaw de oſſib. part. 2. cap. 2.

(*r*) Borelli de motu animal. pars 1. ſchol. ad propoſ. 58. Parent, Hiſtoire de l'acad. des ſciences, 1702.

(*s*) Waſſe, Philoſoph. Tranſact. numb. 383. art. 1.

ing the bulk of any part muſt vary according to the different diſtenſion or repletion of the veſſels compoſing it, we may underſtand how we become taller after a plentiful meal, and decreaſe after faſting or evacuations (*t*). 6. From the different articulations of the bodies and oblique proceſſes of the vertebrae, and the different ſtrength of the ligaments, it is plain, that they are formed ſo as to allow much larger motion forwards than backwards; this laſt being of much leſs uſe, and might be dangerous, by over-ſtretching the large blood-veſſels that are contiguous to the bodies of the vertebrae (*u*). 7. The intervertebral cartilages ſhrivelling as they become more ſolid by age, is the cauſe why old people generally bow forwards, and cannot raiſe their bodies to ſuch an erect poſture as they had in their youth.

The *uſes* of the true vertebrae are, to give us an erect poſture; to allow ſufficient and ſecure motion to the head, neck, and trunk of the body; and to ſupport and defend the bowels and other ſoft parts.

At the ordinary time of birth, each vertebra conſiſts of three bony pieces, connected by cartilages; to wit, the body, which is not fully oſſified; and a long crooked bone on each ſide, on which we ſee a ſmall ſhare of the bony bridge, the oblique proceſſes complete, the beginning tranſverſe proceſſes, and the oblique plate, but no ſpinal proceſſes: So that the teguments are in no danger of being hurt by the ſharp ends of theſe ſpinal proceſſes, while a child is in its bended poſture in the womb, nor while it is ſqueezed in the birth.

From this general mechaniſm of the ſpine, an account is eaſily deduced of all the different preternatural curvatures of which the ſpine is capable: For if one or more vertebrae, or their cartilages, are of unequal thickneſs in oppoſite ſides, the

(*t*) Abbe Fontenu. Hiſtoire de l'acad. des ſciences, 1725.

(*u*) Galen. de uſu part. lib. 1. cap. 16.

the ſpine muſt be reclined over to the thinner ſide; which now ſuſtaining the greateſt ſhare of the weight, muſt ſtill be more compreſſed, conſequently hindered from extending itſelf in proportion to the other ſide, which, being too much freed of its burden, has liberty to enjoy a luxuriant growth. The cauſes, on which ſuch an inequality of thickneſs in different ſides of the vertebrae depends, may vary. For either it may be owing to an over-diſtenſion of the veſſels of one ſide, and from thence a preternatural increaſe of the thickneſs of that part: Or, which is more commonly the caſe, it may proceed from an obſtruction of the veſſels, by which the application of proper nouriſhment to the bony ſubſtance is hindered; whether that obſtruction depends on the faulty diſpoſition of the veſſels or fluids, or if it is produced by an unequal mechanical preſſure occaſioned by a paralytic weakneſs of the muſcles and ligaments, or by a ſpaſmodic over-action of the muſc'es on any ſide of the ſpine, or by people continuing long, or putting themſelves frequently, into any poſture declining from the erect one: In all theſe caſes one common effect follows, to wit, the vertebrae, or their cartilages, or both, turn thick on that ſide where the veſſels are free, and remain thin on the other ſide where theſe veſſels are ſtraightened or obſtructed.—Whenever any morbid curvature is thus made, a ſecond turn, but in an oppoſite direction to the former, muſt be formed; both becauſe the muſcles on the convex ſide of the ſpine being ſtretched, muſt have a ſtronger natural contraction to draw the parts to which their ends are fixed, and becauſe the patient makes efforts to keep the centre of gravity of the body perpendicular to its baſe, that the muſcles may be relieved from a conſtant violent contractile ſtate, which always creates uneaſineſs and pain.

Whence once we underſtand how theſe crooked ſpines are produced, there is little difficulty in forming a juſt prognoſis;

noſis, and a proper method of cure may be eaſily contrived, which muſt vary as to the internal medicines, according to the different cauſes on which the diſeaſe depends: But one general indication muſt be purſued by ſurgeons; which is, to counteract the bending force, by increaſing the compreſſion on the convex part of the curvature, and diminiſhing it on the concave ſide. The manner of executing which in particular caſes muſt be different, and requires a very accurate examination of the circumſtances both of the diſeaſe and patient. In many ſuch caſes I have found ſome ſimple directions, as to poſtures in which the patient's body ſhould be kept, of very great advantage.

THOUGH the true vertebrae agree in the general ſtructure which I have hitherto deſcribed; yet, becauſe of ſeveral ſpecialties proper to a particular number, they are commonly divided into three claſſes, viz. *cervical*, *dorſal*, and *lumbar*.

The CERVICAL (*x*) are the ſeven uppermoſt vertebrae; which are diſtinguiſhed from the reſt by theſe marks.——Their bodies are ſmaller and more ſolid than any others; and flatted on the fore-part, to make way for the oeſophagus; or rather this flat figure is owing to the preſſure of that pipe, and to the action of the longi colli and anterior recti muſcles. They are alſo flat behind, where ſmall proceſſes riſe, to which the internal ligaments are fixed. The upper ſurface of the body of each vertebra is made hollow, by a ſlanting thin proceſs which is raiſed on each ſide:——The lower ſurface is alſo hollowed, but in a different manner, for here the poſterior edge is raiſed a little, and the anterior one is conſiderably produced.——Hence we ſee how the cartilages between thoſe bones are firmly connected, and their articulations are ſecure.

(*x*) Γραχηλια, αυχην, αυχενος, Colli.

The cartilages between theſe vertebrae are thick, eſpecially at their fore-part; which is one reaſon why the vertebrae advance forward as they deſcend, and have larger motion.

The oblique proceſſes of theſe bones of the neck more juſtly deſerve that name than thoſe of any other vertebrae. They are ſituated ſlanting; the upper ones having their ſmooth and almoſt flat ſurfaces facing obliquely backwards and upwards, while the inferior oblique proceſſes have theſe ſurfaces facing obliquely forwards and downwards.

The tranſverſe proceſſes of theſe vertebrae are framed in a different manner from thoſe of any other bones of the ſpine: For, beſides the common tranſverſe proceſs riſing from between the oblique proceſſes of each ſide, there is a ſecond one that comes out from the ſide of the body of each vertebra; and theſe two proceſſes, after leaving a circular hole for the paſſage of the cervical artery and vein, unite, and are conſiderably hollowed at their upper part, with riſing ſides, to protect the nerves that paſs in the hollow; and at laſt each ſide terminates in an obtuſe point, for the inſertion of muſcles.

The ſpinal proceſſes of theſe cervical bones ſtand nearly ſtraight backwards, are ſhorter than thoſe of any other vertebrae, and are forked or double at their ends; and hence allow a more convenient inſertion to muſcles.

The thick cartilages between the bodies of theſe cervical vertebrae, the obliquity of their oblique proceſſes, and the ſhortneſs and horizontal ſituation of their ſpinal proceſſes, all conſpire to allow them large motion.

The holes between the bony croſs bridges, for the paſſage of the nerves from the ſpinal marrow, have their largeſt ſhare formed in the loweſt of the two vertebrae, to which they are common.

So far moſt of the cervical vertebrae agree; but they have

have ſome particular differences, which oblige us to conſider them ſeparately.

The firſt, from its uſe of ſupporting the head, has the name of *atlas* (*y*); and is alſo called *epiſtrophea* from the motion it performs on the ſecond.

The atlas, contrary to all the other vertebrae of the ſpine, has no body; but, inſtead of it, there is a bony arch.——In the convex fore-part of this arch a ſmall riſing appears, where the muſculi longi colli are inſerted; and, on each ſide of this protuberance, a ſmall cavity may be obſerved, where the recti interni minores take their riſe.——The upper and lower parts of the arch are rough and unequal, where the ligaments that connect this vertebra to the os occipitis, and to the ſecond vertebra, are fixed.——The back-part of the arch is concave, ſmooth, and covered with a cartilage, in a recent ſubject, to receive the tooth like proceſs of the ſecond vertebra.——In a firſt vertebra, from which the ſecond has been ſeparated, this hollow makes the paſſage for the ſpinal marrow to ſeem much larger than it really is: On each ſide of it a ſmall rough ſinuoſity may be remarked, where the ligaments going to the ſides of the tooth-like proceſs of the following vertebra are faſtened; and on each ſide a ſmall rough protuberance and depreſſion is obſervable, where the tranſverſe ligament, which ſecures the tooth-like proceſs in the ſinuoſity, is fixed, and hinders that proceſs from injuring the medulla ſpinalis in the flections of the head.

The atlas has as little ſpinal proceſs as body; but inſtead of it, there is a large bony arch, that the muſcles which paſs over this vertebra at that place might not be hurt in extending the head. On the back and upper part of this arch there are two depreſſions, where the recti poſtici minores take their riſe; and at the lower part are two other ſinuoſities,

(*y*) Ατλας.

ſinuoſities, into which the ligaments that connect this bone to the following one are fixed.

The ſuperior oblique proceſs of this atlas are large, oblong, hollow, and more horizontal than in any other vertebra.——They riſe more in their external than internal brim; by which their articulations with the condyloid proceſſes of the os occipitis are firmer.——Under the external edge of each of theſe oblique proceſſes is the foſſa, or deep open channel, in which the vertebral arteries make the circular turn, as they are about to enter the great foramen of the occipital bone, and where the tenth pair of nerves go out.——In ſeveral bodies I have ſeen this foſſa covered with bone.——The inferior oblique proceſſes, extending from within outwards and downwards, are large, concave, and circular. So that this vertebra, contrary to the other ſix, receives the bones with which it is articulated, both above and below.

The tranſverſe proceſſes here are not much hollowed or forked; but are longer and larger than thoſe of any other vertebra of the neck, from the origin and inſertion of ſeveral muſcles; of which thoſe that ſerve to move this vertebra on the ſecond have a conſiderable lever to act with, becauſe of the diſtance of their inſertion from the axis of revolution.

The hole for the ſpinal marrow is larger in this than in any other vertebra, not only on account of the marrow being largeſt here, but alſo to prevent its being hurt by the motions of this vertebra on the ſecond one. This large hole, and the long tranſverſe proceſſes, make this the broadeſt vertebra of the neck.

The condyles of the os occipitis move forwards and backwards in the ſuperior oblique proceſſes of this vertebra; but from the figure of the bones forming theſe joints, it appears,

appears, that very little motion can here be allowed to either ſide; and there muſt be ſtill leſs circular motion.

In new born children this vertebra has only the two lateral pieces offified; the arch, which it has at its fore-part inſtead of a body, being cartilaginous.

The ſecond vertebra colli is called *dentata*, from the tooth-like proceſs on the upper part of its body. Some authors call it *epiſtrophea*; but improperly, ſince this deſignation is only applicable to the firſt, which moves on this as on an axis.

The body of this vertebra is ſomewhat of a pyramidal figure, being large, and produced downwards, eſpecially at its fore-ſide, to enter into a hollow of the vertebra below; while the upper part has a ſquare proceſs, with a ſmall point ſtanding out from it. This it is that is imagined to reſemble a tooth (z), and has given name to the vertebra.—The ſide of this proceſs, on which the hollow of the anterior arch of the firſt vertebra plays, is convex, ſmooth, and covered with a cartilage; and it is of the ſame form behind, for the ligament, which is extended tranſverſely from one rough protuberance of the firſt vertebra to the other, and is cartilaginous in the middle, to move on it.——A ligament likewiſe goes out in an oblique tranſverſe direction, from each ſide of the proceſſus dentatus, to be fixed at its other end to the firſt vertebra, and to the occipital bone; and another ligament riſes up from near the point of the proceſs to the os occipitis.

The ſuperior oblique proceſſes of this vertebra dentata are large, circular, very nearly in an horizontal poſition, and ſlightly convex, to be adapted to the inferior oblique proceſſes of the firſt vertebra.—A moveable cartilage is ſaid by ſome authors to be interpoſed between theſe oblique proceſſes of the firſt and ſecond vertebra; but I could never find

(z) Conoides, pyrenoides, odontoides.

find it.—The inferior oblique proceſſes of this vertebra dentata anſwer exactly to the deſcription given of thoſe common to all the cervical vertebrae.

The tranſverſe proceſſes of the vertebra dentata are ſhort, very little hollowed at their upper part, and not forked at their ends; and the canals through which the cervical arteries paſs are reflected outwards about the middle ſubſtance of each proceſs; ſo that the courſe of theſe veſſels may be directed towards the tranſverſe proceſſes of the firſt vertebra. Had this curvature of the arteries been made in a part ſo moveable as the neck is, while they were not defended by a bone, and fixed to that bone, ſcarce a motion could have been performed without the utmoſt hazard of compreſſion, and a ſtop put to the courſe of the liquids, with all its train of bad conſequences. Hence we obſerve this ſame mechaniſm ſeveral times uſed, when there is any occaſion for a ſudden curvature of a large artery. This is the third remarkable inſtance we have ſeen of it. The firſt was the paſſage of the carotids through the temporal bones; and the ſecond was that lately deſcribed in the vertebral arteries, turning round the oblique proceſſes of the firſt vertebra, to come at the great hole of the occipital bone.

The ſpinal proceſs of this vertebra dentata is thick, ſtrong, and ſhort, to give ſufficient origin to the muſculi recti majores and obliqui inferiores, and to prevent the contuſion of theſe and other muſcles in pulling the head back.

This ſecond vertebra conſiſts, at the birth, of four bony pieces: For, beſides the three which I have already mentioned as common to all the vertebrae, the tooth-like proceſs of this bone is begun at this time to be oſſified in its middle, and is joined as an appendix to the body of the bone.——Leſt this appendix be bent or diſplaced, nurſes ought to keep the heads of new born children from falling

top

too far backwards, by ſtay-bands, or ſome ſuch means, till the muſcles attain ſtrength ſufficient to prevent that dangerous motion.

When we are acquainted with the ſtructure and articulations of the firſt and ſecond vertebrae, and know exactly the ſtrength and connection of their ligaments, there is no difficulty in underſtanding the motions that are performed upon, or by, the firſt; though this ſubject was formerly matter of hot diſpute among ſome of the greateſt anatomiſts (*a*). It is not my purpoſe at preſent to enter into a detail of the reaſons advanced by either party; but to explain the fact, as any one may ſee it, who will remove the muſcles, which, in a recent ſubject, hinder the view of theſe two joints, and then will turn the head into all the different poſitions of which it is capable. The head may then be ſeen to move forwards and backwards on the firſt vertebra, as was already ſaid, while the atlas performs the circumgyratio upon the ſecond vertebra: The inferior oblique proceſſes of the firſt vertebra ſhuffling eaſily in a circular motion on the ſuperior oblique proceſſes of the ſecond, and its body or anterior arch having a rotation on the tooth-like proceſs, by which the perpendicular ligament that is ſent from the point of the tooth-like proceſs to the occipital bone is twiſted, while the lateral ligaments that fix the proceſſus dentatus to the ſides of the firſt vertebra, and to the os occipitis, are very differently affected; for the one upon the ſide towards which the face is turned by the circumgyratio is much ſhortened and lax, while the oppoſite one is ſtretched and made tenſe, and yielding at laſt no more, prevents the head from turning any farther round on the axis. So that theſe lateral ligaments are the proper moderators of the circumgyratio of the head here; which muſt be larger or ſmaller, as theſe ligaments are weaker

(*a*) See Euſtach. de motu capitis.

weaker or ſtronger, longer or ſhorter, and more or leſs capable of being ſtretched. Beſides the revolution on this axis, the firſt vertebra can move a ſmall way to either ſide; but is prevented from moving backwards or forwards by its anterior arch, and by the croſs ligament, which are both cloſely applied to the tooth-like proceſs. Motion forwards here would have been of very bad conſequence, as it would have brought the beginning of the ſpinal marrow upon the point of the tooth-like proceſs.

The rotatory motion of the head is of great uſe to us on many accounts, by allowing us to apply quickly the organs of the ſenſes to objects: And the axis of rotation was altogether proper to be at this place; for, if it had been at a greater diſtance from the head, the weight of the head, if it had at any time been removed from a perpendicular bearing to the ſmall very moveable joint, and thereby had acquired a long lever, would have broken the ligaments at every turn inconſiderately performed, or theſe ligaments muſt have been formed much ſtronger, and conſequently could not have been connected to ſuch ſmall bones. Neither could this circular motion be performed on the firſt vertebra without danger, becauſe the immoveable part of the medulla oblongata is ſo near, as, at each large turn, the beginning the ſpinal marrow would have been in danger of being twiſted, and of ſuffering by the compreſſion this would have made on its tender fibrils.

It is neceſſary to obſerve, that the lateral or moderator ligaments confined ſo much the motion of the firſt vertebra upon the ſecond, that though this joint may ſerve us on ſeveral occaſions, yet we are often obliged to turn our faces farther round than could be done by this joint alone, without the greateſt danger of twiſting the ſpinal marrow too much, and alſo of luxating the oblique proceſſes: Therefore, in large turns of this kind, the rotation is aſ-

ſiſted

ſiſted by all the vertebrae of the neck and loins; and if this is not ſufficient, we employ moſt of the joints of the lower extremities.——This combination of a great many joints towards the performance of one motion, is alſo to be obſerved in ſeveral other parts of the body; notwithſtanding ſuch motion being generally ſaid to be performed by ſome ſingle joint alone.

The third vertebra of the neck is by ſome called *axis*; but this name is applied to it with much leſs reaſon than to the ſecond.——This third, and the three below, have nothing particular in their ſtructure, but all their parts come under the general deſcription formerly given, each of them being larger as they deſcend.

The ſeventh (*b*) vertebra of the neck is near to the form of thoſe of the back, having the upper and lower ſurfaces of its body leſs hollow than the others: The oblique proceſſes are more perpendicular; neither ſpinal nor tranſverſe proceſſes are forked. This ſeventh and the ſixth vertebra of the neck have the hole in each of their tranſverſe proceſſes more frequently divided by a ſmall croſs bridge, that goes between the cervical vein and artery, than any of the other vertebrae.

The twelve DORSAL (*c*) may be diſtinguiſhed from the other vertebrae of the ſpine by the following marks.

Their bodies are of a middle ſize, between thoſe of the neck and loins:—they are more convex before than either of the other two ſorts; and are flatted laterally by the preſſure of the ribs, which are inſerted into ſmall cavities formed in their ſides. This flatneſs of their ſides, which makes the figure of theſe vertebrae almoſt an half oval, is of good uſe; as it affords a firm articulation to the ribs, allows the trachea arteria to divide at a ſmall angle, and

(*b*) Atlas quibuſdam, maxima, magna vertebra, prominens.

(*c*) Θωρακος, μεταφρενα, νωτα, υποτραχηλια, antiſterni, pectoris, tergi.

the other large veſſels to run ſecure from the action of the vital organs.——Their bodies are more concave behind than any of the other two claſſes.——Their upper and lower ſurfaces are horizontal.

The cartilages interpoſed between the bodies of theſe vertebrae are thinner than in any other of the true vertebrae; and contribute to the concavity of the ſpine in the thorax, by being thinneſt at their fore-part.

The oblique proceſſes are placed almoſt perpendicularly; the upper ones ſlanting but a little forwards, and the lower ones ſlanting as much backwards.——Their convexity or concavity is not ſo remarkable as to require particular notice. Between the oblique proceſſes of oppoſite ſides, ſeveral ſharp proceſſes ſtand out from the upper and lower parts of the plates which join to form the ſpinal proceſſes; into theſe ſharp proceſſes, ſtrong ligaments are fixed for connecting the vertebrae.

The tranſverſe proceſſes of the dorſal vertebrae are long, thicker at their ends than in the middle, and turned obliquely backwards; which may be owing to the preſſure of the ribs, the tubercles of which are inſerted into a depreſſion near the end of theſe proceſſes.

The ſpinal proceſſes are long, ſmall-pointed, and ſloping downwards and backwards: from their upper and back-part a ridge riſes, which is received by a ſmall channel in the fore-part of the ſpinal proceſs immediately above, which is here connected to it by a ligament.

The conduit of the ſpinal marrow is here more circular, but, correſponding to the ſize of that cord, is ſmaller than in any of the other vertebrae; and a larger ſhare of the holes in the bony bridges, for the tranſmiſſion of the nerves, is formed in the vertebra above than in the one below.

The

The connection of the dorſal vertebrae to the ribs, the thinneſs of their cartilages, the erect ſituation of the oblique proceſſes, the length, ſloping, and connection of the ſpinal proceſſes, all contribute to reſtrain theſe vertebrae from much motion, which might diſturb the actions of the heart and lungs; and in conſequence of the little motion allowed here, the intervertebral cartilages ſooner ſhrivel, by becoming more ſolid; and therefore the firſt remarkable curvature of the ſpine obſerved, as people advance to old age, is in the leaſt ſtretched vertebrae of the back; or old people firſt become round ſhouldered.

The bodies of the four uppermoſt dorſal vertebrae deviate from the rule of the vertebrae becoming larger as they deſcend: for the firſt of the four is the largeſt, and the other three below gradually become ſmaller, to allow the trachea and large veſſels to divide at ſmaller angles.

The two uppermoſt vertebrae of the back, inſtead of being very prominent forwards, are flatted by the action of the muſculi longi colli and recti majores.

The proportional ſize of the two little depreſſions in the body of each vertebra for receiving the heads of the ribs, ſeems to vary in the following manner; the depreſſion on the upper edge of each vertebra decreaſes as far down as the fourth, and after that increaſes.

The tranſverſe proceſſes are longer in each lower vertebra to the ſeventh or eighth, with their ſmooth ſurfaces, for the tubercles of the ribs, facing gradually more downwards; but afterwards, as they deſcend, they become ſhorter, and the ſmooth ſurfaces are directed more upwards.

The ſpinous proceſſes of the vertebrae of the back become gradually longer and more ſlanting, from the firſt as far down as the eighth or ninth vertebra; from which they manifeſtly turn ſhorter and more erect.

The

The firſt (*d*) vertebra, beſides an oblong hollow in its lower edge, that aſſiſts in forming the cavity wherein the ſecond rib is received, has the whole cavity for the head of the firſt rib formed in it.

The ſecond has the name of axillary (*e*), without any thing particular in its ſtructure.

The eleventh (*f*) often has the whole cavity for the eleventh rib in its body, and wants the ſmooth ſurface on each tranſverſe proceſs.

The twelfth (*g*) always receives the whole head of the laſt rib, and has no ſmooth ſurfaces of its tranſverſe proceſſes, which are very ſhort.——The ſmooth ſurfaces of its inferior oblique proceſſes face outwards as the lumbar do.———And we may ſay in general, that the upper vertebrae of the back loſe gradually their reſemblance to thoſe of the neck, and the lower ones come nearer to the figure of the lumbar.

The articulation of the vertebrae of the back with the ribs, ſhall be more particularly conſidered after the ribs are deſcribed. Only it may be proper now to remark, that the ligaments which ſerve that articulation aſſiſt in connecting the vertebrae.

The loweſt order of the true vertebrae is the *lumbar* (*h*), which are five bones, that may be diſtinguiſhed from any others by theſe marks: 1. Their bodies, though of a circular form at their fore-part, are ſomewhat oblong from one ſide to the other; which may be occaſioned by the preſſure of the large veſſels, the aorta and cava, and of the viſcera. The epiphyſes on their edges are larger; and therefore the upper and lower ſurfaces of their bodies are more concave than in the vertebrae of the back. 2. The cartilages

(*d*) Λοφια, Gutturalis. (*e*) Μασχαλιστηρ.

(*f*) Αρρεπης, in neutram partem inclinans.

(*g*) Διαζωστηρ, Praecingens.

(*h*) Οσφυος, ἰξυος, ψοιων, Renum, lumborum.

cartilages between these vertebrae are much the thickest of any, and render the spine convex within the abdomen, by their greatest thickness being at their fore-part. 3. The oblique processes are strong and deep; those in opposite sides being almost placed in parallel planes; the superior, which are concave, facing inwards, and the convex inferior ones facing outwards: And therefore each of these vertebrae receives the one above it, and is received by the one below; which is not so evident in the other two classes already described. 4. Their transverse processes are small, long, and almost erect, for allowing large motion to each bone, and sufficient insertion to muscles, and for supporting and defending the internal parts. 5. Between the roots of the superior oblique and transverse processes a small protuberance may be observed, where some of the muscles that raise the trunk of the body are inserted. 6. Their spinal processes are strong, straight, and horizontal, with broad flat sides, and a narrow edge above and below; this last being depressed on each side by muscles. And at the root of these edges, we see rough surfaces for fixing the ligaments. 7. The canal for the numerous cords called *cauda equina*, into which the spinal marrow divides, is rather larger in these bones than what contains that marrow in the vertebrae of the back. 8. The holes for the passage of the nerves are more equally formed out of both the contiguous vertebrae than in the other classes; the upper one furnishes, however, the larger share of each hole.

The thick cartilages between these lumbar vertebrae, their deep oblique processes, and their erect spinal processes, are all fit for allowing large motion: Though it is not so great as what is performed in the neck; which appears from comparing the arches which the head describes when moving on the neck or the loins only.

The lumbar vertebrae, as they descend, have their oblique

processes

proceſſes at a greater diſtance from each other, and facing more backwards and forwards.

Both tranſverſe and ſpinal proceſſes of the middlemoſt vertebrae of the loins are longeſt and thickeſt; in the vertebrae above and below, they are leſs: So that theſe proceſſes of the firſt (*i*) and fifth (*k*) are the leaſt, to prevent their ſtriking on the ribs or oſſa ilium, or their bruiſing the muſcles in the motions of the ſpine.

The epiphyſes round the edges of the bodies of the lumbar vertebrae are moſt raiſed in the two loweſt; which conſequently make them appear hollower in the middle than the others are.

The body of the fifth vertebra is rather thinner than that of the fourth.—The ſpinal proceſs of this fifth is ſmaller, and the oblique proceſſes face more backwards and forwards than in any other lumbar vertebra.

After conſidering the ſtructure of the particular vertebrae, and their mutual connection, we may obſerve a ſolicitous care taken that they ſhall not be disjoined without great difficulty. For, beſides being connected by ſtrong ligaments proportioned to the forces which are to be reſiſted, their bodies either enter ſo into each other as to prevent their being diſplaced any way, as in the vertebrae of the neck; or they are propped on all ſides, as thoſe of the back are by the ribs; or their ſurfaces of contact are ſo broad, as to render the ſeparation almoſt impracticable, as in the loins: While the depth and articulation of the oblique proceſſes are exactly proportioned to the quantity of motion which the other parts of the bones allow, or the muſcles can perform. Yet as theſe oblique proceſſes are ſmall, and therefore not capable of ſo ſecure a conjunction as the larger bodies, they may ſooner yield

(*i*) Νεφριτης, Renalis. (*k*) Ασχολιτης, Fulciens.

yield to a disjoining force; but then their diſlocation is not of near ſo bad conſequence as the ſeparation of the bodies would be: For by the oblique proceſſes being diſlocated, the muſcles, ligaments, and ſpinal marrow, are indeed ſtretched; but this marrow muſt be compreſſed, or entirely deſtroyed, when the body of the vertebra is removed out of its place.

THE FALSE VERTEBRAE.

The False Vertebrae compoſe the under pyramid of the ſpine. They are diſtinguiſhed from the bones already deſcribed juſtly enough by this epithet of falſe; becauſe, though each bone into which they can be divided in young people reſembles the true vertebrae in figure, yet none of them contribute to the motion of the trunk of the body; they being intimately united to each other in adults, except at their lower part, where they are moveable; whence they are commonly divided into two bones, *os ſacrum*, and *os coccygis*.

Os Sacrum (*l*), is ſo called, from being offered in ſacrifice by the antients, or rather becauſe of its largeneſs in reſpect of the other vertebrae.—This bone is of an irregular triangular ſhape, broad above, narrow below; convex behind, for the advantageous origin of the muſcles that move the ſpine and thigh backwards; and concave before, for enlarging the capacity of the pelvis.—Four tranſverſe lines, of a colour different from the reſt of the bone which are ſeen on its fore-part, are the marks of diviſion of the five different bones of which it conſiſts in young perſons.

The fore-part of the os ſacrum, analogous to the bodies of the true vertebrae, is ſmooth and flat, to allow a large ſpace

(*l*) Ιερον, σπονδυλος μιλας, Hippocrat. υπερσπονδυλον, Oribaſ. Πλατυ, Latum, os clunium, clavium.

ſpace for the contained bowels, without any danger of hurting them; or this flat figure may be owing to the equal preſſure of theſe bowels, particularly of the laſt gut.—The back-part of it is almoſt ſtraight, without ſo large a cavity as the vertebrae have; becauſe the ſpinal marrow, now ſeparated into the cauda equina, is ſmall.——The bridges between the bodies and proceſſes of this bone are much thicker, and in proportion ſhorter, than in the former claſs of bones.—The ſtrength of theſe croſs bridges is very remarkable in the three upper bones, and is well proportioned to the incumbent weight of the trunk of the body, which theſe bridges ſuſtain in a tranſverſe, conſequently an unfavourable, ſituation, when the body is erect.

There are only two oblique proceſſes of the os ſacrum; one ſtanding out on each ſide from the upper part of the firſt bone.—Their plain erect ſurfaces face backwards, and are articulated with the inferior oblique proceſſes of the laſt vertebra of the loins, to which each of theſe proceſſes is connected by a ſtrong ligament, which riſes from a ſcabrous cavity round their roots, where mucilaginous glands are alſo lodged.—Inſtead of the other oblique proceſſes of this bone, four rough tubercles are to be ſeen on each ſide of its ſurface behind, from which the muſculus ſacer has its origin.

The tranſverſe proceſſes here are all grown together into one large ſtrong oblong proceſs on each ſide; which, ſo far as it anſwers to the firſt three bones, is very thick, and divided into two irregular cavities by a long perpendicular ridge——The foremoſt of the two cavities has commonly a thin cartilaginous ſkin covering it in the recent ſubject, and is adapted to the unequal protuberance of the os ilium; and a ſtrong ligament connects the circumference of theſe ſurfaces of the two bones.——The cavity behind is divided by a tranſverſe ridge into two, where ſtrong

ligamentous

ligamentous ſtrings that go from this bone to the os ilium, with a cellular ſubſtance containing mucus, are lodged.

The tranſverſe proceſſes of the two laſt bones of the os ſacrum are much ſmaller than the former.—At their back-part near their edge, a knob and oblong flat ſurface give riſe to two ſtrong ligaments which are extended to the os iſchium; and are therefore called *ſacroſciatic*.

The ſpinal proceſſes of the three uppermoſt bones of the os ſacrum appear ſhort, ſharp, and almoſt erect, while the two lower ones are open behind; and ſometimes a little knob is to be ſeen on the fourth, though generally it is bifurcated, without the two legs meeting into a ſpine; in which condition alſo the firſt is often to be ſeen; and ſometimes none of them meet, but leave a ſinus, or rather foſſa, inſtead of a canal (*m*).—The muſculus latiſſimus and longiſſimus dorſi, ſacro-lumbalis, and glutaeus maximus, have part of their origins from theſe ſpinal proceſſes.

The canal between the bodies and proceſſes of this bone, for the cauda equina, is triangular; and becomes ſmaller, as the cauda alſo does, as it deſcends.—Below the third bone, this paſſage is no more a complete bony canal, but is open behind; and is only there defended by a ſtrong ligamentous membrane ſtretched over it, which, with the muſcles that cover it, and that are very prominent on each ſide, is a ſufficient defence for the bundle of nerves within.

At the root of each oblique proceſs of this bone, the notch is conſpicuous; by which, and another ſimilar one in the laſt vertebra of the loins, a paſſage is left for the twenty-fourth ſpinal nerve; and in viewing the os ſacrum, either before or behind, four large holes appear in each ſide, in much the ſame height as where the marks of the union of its ſeveral bones remain. Some of the largeſt nerves of the body paſs through the anterior holes; and ſuperficial

(*m*) Verheyen, Anat. tract. 5. cap. 9.—Sue Trad. d'oſteol. p. 127.

grooves, running outwards from them in different directions, ſhew the courſe of theſe nerves.——From the intervals of theſe grooves, the pyriformis muſcle chiefly riſes.——The holes in the back-part of the bone are covered by membranes which allow ſmall nerves to paſs through them.——The two uppermoſt of theſe holes, eſpecially on the fore-ſide, are the largeſt; and as the bone deſcends, the holes turn ſmaller. Sometimes a notch is only formed at the lower part in each ſide of this bone; and in other ſubjects there is a hole common to it and the os coccygis, through which the twenty-ninth pair of ſpinal nerves paſſes; and frequently a bony bridge is formed on the back-part of each ſide by a proceſs ſent up from the back-part of the os coccygis, and joined to the little knobs which the laſt bone of the os ſacrum has inſtead of a ſpinal proceſs. Under this bridge or jugum, the twenty-ninth pair of ſpinal nerves runs in its courſe to the common holes juſt now deſcribed.

The upper part of the body of the firſt bone reſembles the vertebrae of the loins; but the ſmall fifth bone is oblong tranſverſely, and hollow in the middle of its lower ſurface.

The ſubſtance of the os ſacrum is very ſpongy, without any conſiderable ſolid external plates, and is lighter proportionally to its bulk than any other bone in the body; but it is ſecured from injuries by the thick muſcles that cover it behind, and by the ſtrong ligamentous membranes that cloſely adhere to it.—As this is one of the moſt remarkable inſtances of this ſort of defence afforded a ſoft weak bone, we may make the general obſervation: That wherever we meet with ſuch a bone, one or other, or both theſe defences, are uſed; the firſt to ward off injuries; and the ſecond to keep the ſubſtance of the bone from yielding too eaſily.

This

This bone is *articulated* above to the laſt vertebra of the loins, in the manner that the lumbar vertebrae are joined; and therefore the ſame motions may be performed here.—The articulation of the lower-part of the os ſacrum to the os coccygis, ſeems well enough adapted for allowing conſiderable motion to this laſt bone, were it not much confined by ligaments. Laterally, the os ſacrum is joined to the oſſa ilium by an immoveable ſynchondroſis, or what almoſt deſerves the name of a ſuture: For the cartilaginous cruſt on the ſurface of the bones is very thin; and both their ſurfaces are ſo ſcabrous and unequal, as to be indented into each other; which makes ſuch a ſtrong connection, that great force is required to ſeparate them, after all the muſcles and ligaments are cut.—Frequently the two bones grow together in old ſubjects.

The *uſes* of the os ſacrum are, to ſerve as the common baſe and ſupport of the trunk of the body, to guard the nerves proceeding from the end of the ſpinal marrow, to defend the back-part of the pelvis, and to afford ſufficient origin to the muſcles which move the trunk and thigh.

The bones that compoſe the os ſacrum of infants, have their bodies ſeparated from each other by a thick cartilage; and, in the ſame manner as the true vertebrae, each of them conſiſts of a body and two lateral plates, connected together by cartilages; the ends of the plates ſeldom being contiguous behind.

Os Coccygis (*n*), or *rump-bone*, is that triangular chain of bones depending from the os ſacrum; each bone becoming ſmaller as they deſcend, till the laſt ends almoſt in a point. The os coccygis is convex behind, and concave before; from which crooked pyramidal figure, which was thought to reſemble a cuckow's beak, it has got its name.

This

(*n*) Οξυκοκκυγιον, οσεος, Caudae os, ſpondylium os cuculi.

This bone conſiſts of four pieces in people of middle age: —In children, almoſt the whole of it is cartilage: In old ſubjects, all the bones are united, and become frequently one continued bone with the os ſacrum.

The higheſt of the four bones is the largeſt, with ſhoulders extended farther to each ſide than the end of the os ſacrum; which enlargement ſhould, in my opinion, ſerve as a diſtinguiſhing mark to fix the limits of either bone; and therefore ſhould take away all diſpute about reckoning the number of bones, of which one or other of theſe two parts of the falſe vertebrae is compoſed; which diſpute muſt ſtill be kept up, as long as the numbering five or ſix bones in the os ſacrum depends upon the uncertain accident of this broad ſhouldered little bone being united to or ſeparated from it.—The upper ſurface of this bone is a little hollow.—From the back of that bulbous part, called its ſhoulders, a proceſs often riſes up on each ſide, to join with the bifurcated ſpine of the fourth and fifth bones of the os ſacrum, to form the bony bridge mentioned in the deſcription of the os ſacrum.—— Sometimes theſe ſhoulders are joined to the ſides of the fifth bone of the os ſacrum, to form the hole in each ſide common to theſe two bones, for the paſſage of the twenty-ninth pair of ſpinal nerves.—Immediately below the ſhoulders of the os coccygis, a notch may be remarked on each ſide, where the thirtieth pair of the ſpinal nerves paſſes.——The lower end of this bone is formed into a ſmall head, which very often is hollow in the middle.

The three lower bones gradually become ſmaller, and are ſpongy; but are ſtrengthened by a ſtrong ligament which covers and connects them.—Their ends, by which they are articulated, are formed in the ſame manner as thoſe of the firſt bone are.

Between each of theſe four bones of young ſubjects a cartilage

cartilage is interposed; therefore their articulation is analogous to that of the bodies of the vertebrae of the neck: For, as has been above remarked, the lower end of the os sacrum, and of each of the three superior bones of the os coccygis, has a small depression in the middle; and the upper part of all the bones of the os coccygis is a little concave, and consequently the interposed cartilages are thickest in the middle, to fill up both cavities; by which they connect the bones more firmly.—When the cartilages ossify, the upper end of each bone is formed into a cavity, exactly adapted to the protuberant lower end of the bone immediately above.—From this sort of articulation, it is evident, that, unless when these bones grow together, all of them are capable of motion; of which the first and second, especially this last, enjoys the largest share.

The lower end of the fourth bone terminates in a rough point, to which a cartilage is appended.

To the sides of these bones of the os coccygis, the coccygaei muscles (*o*), and part of the levatores ani, and of the glutaei maximi, are fixed.

The substance of these bones is very spongy, and in children cartilaginous; there being only a part of the first bone ossified in a new born infant.——Since therefore the intestinum rectum of children is not so firmly supported as it is in adults, this may be one reason why they are more subject to a procidentia ani than old people (*p*).

From the description of this bone, we see how little it resembles the vertebrae; since it seldom has processes, never has any cavity for the spinal marrow, nor holes for the passage of nerves. Its connection hinders it from being moved to either side: and its motion backwards and forwards is much

(*o*) Douglas, Myograph. chap. 40.—Eustach. tab. 36. N°. 45. 20.

(*p*) Spigel. de humani corp. fabric. lib. 2. cap. 22.—Paaw de ossib. par. 2. cap. 3.

much confined: Yet, as its ligaments can be ſtretched by a conſiderable force, it is of great advantage in the excretion of the faeces alvinae, and much more in child-bearing, that this bone ſhould remain moveable; and the right management of it, in delivering women, may be of great benefit to them (*q*). The mobility of the os coccygis diminiſhing as people advance in age, eſpecially when its ligaments and cartilages have not been kept flexible by being ſtretched, is probably one reaſon why the women, who are old maids before they marry, have generally difficult parturition.

The os coccygis ſerves to ſuſtain the inteſtinum rectum; and, in order to perform this office more effectually, it is made to turn with a curve forwards; by which alſo the bone itſelf, as well as the muſcles and teguments, is preſerved from any injury when we ſit with our body reclined back.

§ 2. *Of the* PELVIS.

THE ſecond part of the trunk of the ſkeleton, viz. the pelvis, is the cylindrical cavity at the lower part of the abdomen formed by the os ſacrum, os coccygis, and oſſa innominata; which laſt therefore fall now in courſe to be examined.

OSSA INNOMINATA.

THOUGH the name of *oſſa innominata* (*r*) contributes nothing to the knowledge of their ſituation, ſtructure, or office; yet they have been ſo long univerſally known by it, that there is no occaſion for changing it. They are two large broad bones, which form the fore-part and ſides of the

(*q*) Paaw, ibid.—Deventer, Operat. chirurg. cap. 27.

(*r*) Σκελιων προσφυσεις, Sacro conjuncta.

the pelvis, and the lower part of the ſides of the abdomen.—In children, each of theſe bones is evidently divided into three; which are afterwards ſo intimately united, that ſcarce the leaſt mark of their former ſeparation remains: They are nevertheleſs deſcribed as conſiſting each of three bones, to wit, the os ilium, iſchium, and pubis; which I ſhall firſt deſcribe ſeparately, and then ſhall conſider what is common to any two of them, or to all the three.

Os ILIUM (*s*), or *haunch-bone*, is ſituated higheſt of the three, and reaches as far down as one third of the great cavity into which the head of the thigh-bone is received.

The external ſide of this bone is unequally convex, and is called its *dorſum*;—the internal concave ſurface is by ſome authors (but improperly) named its *coſta*.——The ſemicircular edge at the higheſt part of this bone, which is tipped with a cartilage in the recent ſubject, is named the *ſpine*, into which the external or deſcending oblique muſcle of the abdomen is inſerted; and from it the internal aſcending oblique, and the tranſverſe muſcles of the belly, with the glutaeus maximus, quadratus lumborum, and latiſſimus dorſi, have their origin. Some writers (*t*) are of opinion, that it is only the tendinous cruſt of all theſe muſcles, and not a cartilage, as is commonly alleged, that covers this bony edge.——The ends of the ſpine being more prominent than the ſurface of the bone below them, are therefore reckoned proceſſes.———From the anterior ſpinal proceſs, the ſartorius and faſcialis muſcles have their riſe, and the outer end of the doubled tendon of the external oblique muſcle of the abdomen, commonly called *Fallopius's* or *Paupart's* ligament, is fixed to it. The inſide of the poſterior ſpinal proceſs, and of part of the ſpine forward

(*s*) Λαγονων κεινεων, Scaphium, lumbare, clunium, clavium, anchas.

(*t*) Winſlow, Expoſition anatomique du corps humain, traité des os frais, § 96.

ward from that, is made flat and rough where the ſacro-lumbalis and longiſſimus dorſi riſe; and to its outſide are fixed ligaments extended to the os ſacrum and tranſverſe proceſſes of the fifth and fourth vertebrae of the loins (*u*).——Below the anterior ſpinal proceſs another protuberance ſtands out, which by its ſituation may be diſtinguiſhed from the former, by adding the epithet of *inferior*, where the muſculus rectus tibiae has its origin (*x*).——Between theſe two anterior proceſſes the bone is hollowed where the beginning of the ſartorius muſcle is lodged.——Below the poſterior ſpinal proceſs, a ſecond protuberance of the edge of this bone is in like manner obſervable, which is cloſely applied to the os ſacrum.——Under this laſt proceſs a conſiderable large niche is obſervable in the os ilium; between the ſides of which and the ſtrong ligament that is ſtretched over from the os ſacrum to the ſharp pointed proceſs of the os iſchium of the recent ſubject, a large hole is formed, through which the muſculus pyriformis, the great ſciatic nerve, and the poſterior crural veſſels, paſs, and are protected from compreſſion.

The external broad ſide, or dorſum of the os ilium, is a little hollow towards the fore-part; farther back, it is as much raiſed; then is conſiderably concave; and, laſtly, it is convex.——Theſe inequalities are occaſioned by the actions of the muſcles that are ſituated on this ſurface.——From behind the uppermoſt of the two anterior ſpinal proceſſes, in ſuch bones as are ſtrongly marked by the muſcles, a ſemicircular ridge is extended to the hollow paſſage of the ſciatic nerve. Between the ſpine and this ridge, the gluteus medius takes its riſe. Immediately from above the loweſt of the anterior ſpinal proceſſes, a ſecond ridge is ſtretched to the niche. Between this and the former ridge,

(*u*) Weitbrecht, Syndeſmolog. ſect. 4. § 39. 40. 46. 47.
(*x*) Baker, Curſ. Oſteolog. demonſtr. 3.

ridge, the glutaeus minimus has its origin.——On the outside of the posterior spinal processes, the dorsum of the os ilium is flat and rough, where part of the musculus glutaeus maximus and pyriformis rises. The lowest part of this bone is the thickest, and is formed into a large cavity with high brims, to assist in composing the great acetabulum; which shall be considered, after all the three bones that constitute the os innominatum are described.

The internal surface of the os ilium is concave in its broadest fore-part, where the internal iliac muscle has its origin, and where some share of the intestinum ilium and colon is lodged. From this large hollow, a small sinuosity is continued obliquely forwards, at the inside of the anterior inferior spinal process, where part of the psoas and iliacus muscles, with the crural vessels and nerves, pass.—The large concavity is bounded below by a sharp ridge, which runs from behind forwards; and, being continued with such another ridge of the os pubis, forms a line of partition between the abdomen and pelvis.—Into this ridge the broad tendon of the psoas parvus is inserted.

All the internal surface of the os ilium, behind this ridge, is very unequal: For the upper part is flat, but spongy, where the sacro-lumbalis and longissimus dorsi rise.——Lower down, there is a transverse ridge from which ligaments go out to the os sacrum.——Immediately below this ridge, the rough unequal cavities and prominences are placed, which are exactly adapted to those described on the side of the os sacrum. In the same manner, the upper part of this rough surface is porous, for the firmer adhesion of the ligamentous cellular substance; while the lower part is more solid, and covered with a thin cartilaginous skin, for its immoveable articulation with the os sacrum.——From all the circumference of this large unequal surface, ligaments are extended to the os

ſacrum, to ſecure more firmly the conjunction of theſe bones.

The paſſages of the medullary veſſels are very conſpicuous, both in the dorſum and coſta of many oſſa ilium; but in others they are inconſiderable.

The poſterior and lower parts of theſe bones are thick; but they are generally exceedingly thin and compact at their middle, where they are expoſed to the actions of the muſculi glutæi and iliacus internus, and to the preſſure of the bowels contained in the belly.—The ſubſtance of the oſſa ilium is moſtly cellular, except a thin external table.

In a ripe child, the ſpine of the os ilium is cartilaginous; and is afterwards joined to the bone, in form of an epiphyſe. ——The large lower end of this bone is not completely offified.

Os Ischium (*y*), or hip-bone, is of a middle bulk between the two other parts of the os innominatum, is ſituated loweſt of the three, and is of a very irregular figure. Its extent might be marked by a horizontal line drawn near through the middle of the acetabulum; for the upper bulbous part of this bone forms rather leſs than the lower half of that great cavity, and the ſmall leg of it riſes to much the ſame height on the other ſide of the great hole common to this bone and the os pubis.

From the upper thick part of the os iſchium, a ſharp proceſs, called by ſome authors ſpinous, ſtands out backwards, from which chiefly the muſculus coccygæus and ſuperior gemellus, and part of the levator ani, riſe; and the anterior or internal ſacroſciatic ligament is fixed to it. Between the upper part of this ligament and the bones, it was formerly obſerved that the pyriform muſcle, the poſterior crural veſſels, and the ſciatic nerve, paſs out of the pelvis. —Im-

(*y*) Coxae, coxendicis, pyxis.

—Immediately below this proceſs, a ſinuoſity is formed for the tendon of the muſculus obturator internus.———In a recent ſubject this part of the bone, which ſerves as a pulley on which the obturator muſcle plays, is covered with a ligamentous cartilage, that, by two or three ſmall ridges, points out the interſtices of the fibres in the tendon of this muſcle. ——The outer ſurface of the bone at the root of this ſpinous proceſs is made hollow by the pyriformis or iliacus externus muſcle.

Below the ſinuoſity for the obturator muſcle, is the great knob or tuberoſity, covered with cartilage or tendon (z). ——The upper part of the tuberoſity gives riſe to the inferior gemellus muſcle.——To a ridge at the inſide of this the external or poſterior ſacroſciatic ligament is ſo fixed, that between it, the internal ligament, and the ſinuoſity of the os iſchium, a paſſage is left for the internal obturator muſcle. ——The upper thick ſmooth part of the *tuber*, called by ſome its *dorſum*, has two oblique impreſſions on it. The inner one gives origin to the long head of the biceps flexor tibiae, and ſeminervoſus muſcles; and the ſemimembranoſus riſes from the exterior one, which reaches higher and nearer the acetabulum than the other.——The lower, thinner, more ſcabrous part of the knob which bends forwards, is alſo marked with two flat ſurfaces; whereof the internal is what we lean upon in ſitting, and the external gives riſe to the largeſt head of the triceps adductor femoris. Between the external margin of the tuberoſity and the great hole of the os innominatum, there is frequently an obtuſe ridge extended down from the acetabulum, which gives origin to the quadratus femoris.——As the tuber advances forwards, it becomes ſmaller, and is rough, for the origin of the muſculus tranſverſalis and erector penis.—The ſmall leg of it, which mounts upwards to join the os pubis, is rough and prominent at

(z) Winſlow, Expoſit. Anat. des os frais, § 96.

at its edge, where the two lower heads of the triceps or quadriceps adductor femoris take their rise.

The upper and back-part of the os ischium is broad and thick; but its lower and fore-part is narrower and thinner. ——Its substance is of the structure common to broad bones.

The os ilium and pubis of the same side are the only bones which are contiguous to the os ischium.

The part of the os ischium, which forms the acetabulum, the spinous process, the great tuber, and the recurved leg, are all cartilaginous at birth.—The tuber, with part of the leg or process above it, becomes an epiphyse before this bone is fully formed.

The Os Pubis (*a*), *share-bone*, is the least of the three parts of the os innominatum, and is placed at the upper fore-part of it.———The thick largest part of this bone is employed in forming the acetabulum; from which becoming much smaller, it is stretched inwards to its fellow of the other side, where it again grows larger, and sends a small branch downwards to join the end of the small leg of the os ischium.———The upper fore part of each os pubis is tuberous and rough where the musculus rectus and pyramidalis are inserted.—From this a ridge is extended along the upper edge of the bone, in a continued line with such another of the os ilium, which divides the abdomen and pelvis. The ligament of Fallopius is fixed to the internal end of this ridge, and the smooth hollow below it is made by the psoas and iliacus internus muscles passing with the anterior crural vessels and nerves behind the ligament.—— Some way below the former ridge, another is extended from the tuberous part of the os pubis downwards and outwards

(*a*) ἦβης, Pectinis, penis, pudibundum, fenestratum.

outwards towards the acetabulum; between these two ridges the bone is hollow and smooth, for lodging the head of the pectineus muscle.—Immediately below, where the lower ridge is to take the turn downwards, a winding niche is made, which is comprehended in the great foramen of a skeleton; but is formed into a hole by a subtended ligament in the recent subject, for the passage of the posterior crural nerve, an artery, and a vein.——The internal end of the os pubis is rough and unequal, for the firmer adhesion of the thick ligamentous cartilage that connects it to its fellow of the other side:——The process which goes down from that to the os ischium is broad and rough before, where the gracilis and upper heads of the triceps, or rather *quadriceps*, adductor femoris have their origin.

The substance of the os pubis is the same as of other broad bones.

Only a part of the large end of this bone is ossified, and the whole leg is cartilaginous, in a child born at the full time.

Between the os ischium and pubis a very large irregular hole is left, which, from its resemblance to a door or shield, has been called *thyroides*. This hole is all, except the niche for the posterior crural nerve, filled up, in a recent subject, with a strong ligamentous membrane, that adheres very firmly to its circumference. From this membrane chiefly the two obturator muscles, external and internal, take their rise.—The great design of this hole, besides rendering the bone lighter, is to allow a strong enough origin to the obturator muscles, and sufficient space for lodging their bellies, that there may be no danger of disturbing the functions of the contained viscera of the pelvis by the actions of the internal, nor of the external being bruised by the thigh bone, especially by its lesser trochanter, in the motions of the thigh inwards:

wards: Both which inconveniencies muſt have happened, had the offa innominata been complete here, and of ſufficient thickneſs and ſtrength to ſerve as the fixed point of theſe muſcles.—The bowels ſometimes make their way through the niche for the veſſels, at the upper part of this thyroid hole; and this cauſes a hernia in this place (*b*).

In the external ſurface of the offa innominata, near the outſide of the great hole, a large deep cavity is formed by all the three bones conjunctly: For the os pubis conſtitutes about one fifth, the os ilium makes ſomething leſs than two fifths, and the os iſchium as much more than two fifths. The brims of this cavity are very high, and are ſtill much more enlarged by the ligamentous cartilage, with which they are tipped in a recent ſubject. From this form of the cavity it has been called *acetabulum*; and for a diſtinguiſhing character the name of the bone that conſtitutes the largeſt ſhare of it is added; therefore *acetabulum oſſis iſchii* (*c*) is the name this cavity commonly bears. Round the baſe of the ſupercilia the bone is rough and unequal, where the capſular ligament of the articulation is fixed.——The brims at the upper and back part of the acetabulum are much larger and higher than any where elſe; which is very neceſſary to prevent the head of the femur from ſlipping out of its cavity at this place, where the whole weight of the body bears upon it, and conſequently would otherwiſe be in perpetual danger of thruſting it out. As theſe brims are extended downwards and forwards, they become leſs; and at their internal lower part a breach is made in them; from the one ſide of which to the other, a ligament is placed in the recent ſubject; under which a large hole is left, which contains a fatty cellular ſubſtance and veſſels. The reaſon of which appearance has

(*b*) Memoires de l'acad. de chirurgie, tom. 1. p. 709. &c.

(*c*) Coxæ, coxendicis.

has afforded matter of debate. To me it ſeems evidently contrived for allowing a large motion to the thigh inwards: For if the bony brims had been here continued, the neck of the thigh-bone muſt have ſtruck upon them when the thighs were brought acroſs each other; which, in a large ſtrong motion this way, would have endangered the neck of the one bone, or brim of the other. Then the veſſels which are diſtributed to the joint may ſafely enter at the ſinuoſity in the bottom of the breach; which being, however, larger than is neceſſary for that purpoſe, allows the larger mucilaginous gland of the joint to eſcape below the ligament, when the head of the thigh-bone is in hazard of preſſing too much upon it in the motions of the thigh outwards (*d*). Beſides this difference in the height of the brims, the acetabulum is otherwiſe unequal; for the lower internal part of it is depreſſed below the cartilaginous ſurface of the upper part, and is not covered with cartilage; into the upper part of this particular depreſſion, where it is deepeſt and of a ſemilunar form, the ligament of the thigh-bone, commonly, though improperly, called the *round* one is inſerted; while, in its more ſuperficial lower part, the large mucilaginous gland of this joint is lodged. The greateſt part of this ſeparate depreſſion is formed in the os iſchium.

From what has been ſaid of the condition of the three bones compoſing this acetabulum in new-born children, it muſt be evident that a conſiderable part of this cavity is cartilaginous in them.

THE *oſſa innominata* are *joined* at their back-part to each ſide of the os ſacrum by a ſort of ſuture, with a very thin intervening cartilage, which ſerves as ſo much glue to cement thoſe bones together; and ſtrong ligaments go from the

(*d*) Petit. Memoires de l'acad. des ſciences, 1722.

the circumference of this unequal surface, to connect them more firmly. The ossa innominata are connected together at their fore-part by the ligamentous cartilage interposed between the two ossa pubis. These bones can therefore have no motion in a natural state, except what is common to the trunk of the body, or to the os sacrum. But it has been disputed, whether or not they loosen so much from each other, and from the os sacrum, in child-birth, by the flow of mucus to the pelvis, and by the throes of labour, as that the ossa pubis recede from each other, and thereby allow the passage between the bones to be enlarged. Several observations (*e*) shew that this relaxation sometimes happens: But those who had frequently opportunities of dissecting the bodies of women who died immediately after being delivered of children, teach us to beware of regarding this as the common effect of child-birth; for they found such a relaxation in very few of the bodies which they examined (*f*).

Considering the great weight that is supported in our erect posture, by the articulation of the ossa innominata with the os sacrum, there is great reason to think, that if the conglutinated surfaces of these bones were once separated (without which the ossa pubis cannot shuffle on each other), the ligaments would be violently stretched, if not torn: From whence many disorders would arise (*g*).

Each os innominatum affords a socket (the acetabulum) for the thigh-bones to move in; and the trunk of the body rolls here so much on the heads of the thigh-bones, as to allow the most conspicuous motions of the trunk, which are commonly thought to be performed by the bones of the

(*e*) Bauchin. Theat. anat. lib. 1. cap. 49.—Spigel. Anat. lib. 2. c. 24.—Riolan. Anthropog. lib. 6. cap. 12.—Diemerbroek, Anat. lib 9 cap. 16.

(*f*) Hildan. epist. cent. obs. 46.—Dionis Sixieme demonstrat. des os.—Morgagn. Adverss. 3. animad. 15.

(*g*) Ludov. in Ephem. German. dec. 1. ann. 3. obs. 255.

the ſpine. This articulation is to be more fully deſcribed after the oſſa femoris are examined.

The pelvis, then, has a large open above where it is continued with the abdomen; is ſtrongly fenced by bones on the ſides, back and fore-part; and appears with a wide opening below, in the ſkeleton; but, in a recent ſubject, a conſiderable part of the opening is filled by the ſacroſciatic ligaments, pyriform, internal obturator, levatores ani, gemini, and coccygaei muſcles, which ſupport and protect the contained parts better than bones could have done; ſo that ſpace is only left at the loweſt part of it, for the large excretories, the veſica urinaria, inteſtinum rectum, and in females, the uterus, to diſcharge themſelves.

§ 3. *Bones of the* THORAX.

THE THORAX, (*b*), or *cheſt*, which is the only part of the trunk of the body that we have not yet deſcribed, reaches from below the neck to the belly; and, by means of the bones that guard it, is formed into a large cavity: The figure of it is ſomewhat conoidal: but its upper ſmaller end is not finiſhed, being left open for the paſſage of the wind-pipe, gullet, and large blood-veſſels; and its lower part or baſe, has no bones, and is ſhorter before than behind; ſo that, to carry on our compariſon, it appears like an oblique ſection of the conoid. Beſides which, we ought alſo to remark, that the lower part of this cavity is narrower than ſome way above (*i*); and that the middle of its back-part is conſiderably diminiſhed by the bones ſtanding forwards into it.

The bones which form the thorax are the twelve dorſal vertebrae behind, the ribs on the ſides, and the ſternum before.

(*b*) Pectus caſſum. (*i*) Albin. de oſſib. § 169.

The vertebrae have already been deſcribed as part of the ſpine; and therefore are now to be paſſed.

THE RIBS.

The Ribs, or *coſtae* (*k*), (as if they were *cuſtodes*, or guards, to thoſe principal organs of the animal machine, the heart and lungs), are the long crooked bones placed at the ſide of the cheſt, in an oblique direction downwards in reſpect of the back-bone.——Their number is generally twelve on each ſide; though frequently eleven or thirteen have been found (*l*).——Sometimes the ribs are found preternaturally conjoined or divided (*m*).

The ribs are all concave internally; where they are alſo made ſmooth by the action of the contained parts, which, on this account, are in no danger of being hurt by them; and they are convex externally, that they might reſiſt that part of the preſſure of the atmoſphere which is not balanced by the air within the lungs during *inſpiration*.——The ends of the ribs next the vertebrae are rounder than they are after theſe bones have advanced forwards, when they become flatter and broader, and have an upper and lower edge; each of which is made rough by the action of the intercoſtal muſcles inſerted into them. Theſe muſcles, being all of nearly equal force, and equally ſtretched in the interſtices of the ribs, prevent the broken ends of theſe bones, in a fracture, from being removed far out of their natural place, to interrupt the motion of the vital organs.——The upper edge of the ribs is more obtuſe and rounder than the lower, which is depreſſed on its internal ſide by

a

(*k*) Πλευραι, περιτερνα, σπαδικι.

(*l*) Riolan. Comment. de oſſibus, cap. 19.——Marchetti, cap. 9. Cowper. Explicat. tab. 93. and 94.——Morgagn. Adverſ. Anat.

(*m*) Sue, Trad. d'oſteolog. p. 141.

a long foſſa, for lodging the intercoſtal veſſels and nerves; on each ſide of which there is a ridge, to which the intercoſtal muſcles are fixed. The foſſa is not obſervable, however, at either end of the ribs; for, at the poſterior or root, the veſſels have not yet reached the ribs; and, at the fore-end, they are ſplit away into branches, to ſerve the parts between the ribs: Which plainly teaches ſurgeons one reaſon of the greater ſafety of performing the operation of the empyema towards the ſides of the thorax, than either near the back or the breaſt.

At the poſterior end (*n*) of each rib, a little head is formed, which is divided by a middle ridge into two plain or hollow ſurfaces; the loweſt of which is the broadeſt and deepeſt in moſt of them. The two plains are joined to the bodies of two different vertebrae, and the ridge forces itſelf into the intervening cartilage.——A little way from this head, we find on the external ſurface, a ſmall cavity, where mucilaginous glands are lodged; and round the head, the bone appears ſpongy, where the capſular ligament of the articulation is fixed.——Immediately beyond this a flatted tubercle riſes, with a ſmall cavity at, and roughneſs about, its root, for the articulation of the rib with the tranſverſe proceſs of the loweſt of the two vertebrae, with the bodies of which the head of the rib is joined.——Advancing farther on this external ſurface, we obſerve in moſt of the ribs another ſmaller tubercle, into which ligaments connecting the ribs to each other, and to the tranſverſe proceſſes of the vertebrae and portions of the longiſſimus dorſi, are inſerted.——Beyond this the ribs are made flat by the ſacro-lumbalis muſcle, which is inſerted into the part of this flat ſurface fartheſt from the ſpine, where each rib makes a conſiderable curve, called by ſome its angle.——Then the rib begins

(*n*) Κυρτον, Remulus.

begins to turn broad, and continues ſo to its anterior end (*o*), which is hollow and ſpongy, for the reception of, and firm coalition with, the cartilage that runs thence to be inſerted into the ſternum, or to be joined with ſome other cartilage. ——In adults, the cavity at this end of the ribs is generally ſmooth and poliſhed on its ſurface; by which the articulation of the cartilage with it has the appearance of being deſigned for motion, but it has none.

The *ſubſtance* of the ribs is ſpongy, cellular, and only covered with a very thin external lamellated ſurface, which increaſes in thickneſs and ſtrength as it approaches the vertebrae.

To the fore-end of each rib a long, broad, and ſtrong cartilage is fixed, and reaches thence to the ſternum, or is joined to the cartilage of the next rib. This courſe, however, is not in a ſtraight line with the rib: for the cartilages generally make a conſiderable curve, the concave part of which is upwards; therefore, at their inſertion into the ſternum, they make an obtuſe angle above, and an acute one below.——Theſe cartilages are of ſuch a length as never to allow the ribs to come to a right angle with the ſpine; but they keep them ſituated ſo obliquely as to make the angle very conſiderably obtuſe, above, till a force exceeding the elaſticity of the cartilage is applied.——Theſe cartilages, as all others, are firmer and harder internally than they are on their external ſurface; and ſometimes in old people, all their middle ſubſtance becomes bony, while a thin cartilaginous lamella appears externally (*p*). The oſſification, however, begins frequently at the external ſurface.——The greateſt alternate motions of the cartilages being made at their great curvature, that part remains frequently cartilaginous after all the reſt is oſſified (*q*).

The

(*o*) Πλατη, Palmul.

(*p*) Veſal. lib. 2. cap. 19.

(*q*) Havers, Oſteolog. Nov. diſc. 5. p. 289.

The ribs then are *articulated* at each end, of which the one behind is doubly joined to the vertebrae; for the head is received into the cavities of two bodies of the vertebrae, and the larger tubercle is received into the depreſſion in the tranſverſe proceſs of the lower vertebra.—When we examine the double articulation, we muſt immediately ſee that no other motion can be allowed here than upwards and downwards; ſince the tranſverſe proceſs hinders the rib to be thruſt back; the reſiſtance on the other ſide of the ſternum prevents the ribs coming forward; and each of the two joints, with the other parts attached, oppoſe its turning round. But then it is likewiſe as evident, that even the motion upwards and downwards can be but ſmall in any one rib at the articulation itſelf. But as the ribs advance forwards, the diſtance from their centre of motion increaſing, the motion muſt be larger; and it would be very conſpicuous at their anterior ends, were they not reſiſted there by the cartilages, which yield ſo little, that the principal motion is performed by the middle part of the ribs, which turns outwards and upwards, and occaſions the twiſt remarkable in the long ribs at the place near their fore-end where they are moſt reſiſted (*r*).

Hitherto I have laid down the ſtructure and connection which moſt of the ribs enjoy, as belonging to all of them; but muſt now conſider the ſpecialities wherein any of them differ from the general deſcription given, or from each other.

In viewing the ribs from above downwards, their figure is ſtill ſtraighter; the uppermoſt being the moſt crooked of any.——Their obliquity in reſpect of the ſpine increaſes as they deſcend; ſo that though their diſtances from each other is very little different at their back-part, yet at their fore-ends the diſtances between the lower ones muſt increaſe.—

In

(*r*) Winſlow, Memoires de l'acad. des ſciences, 1720.

In conſequence, too, of this increaſed obliquity of the lower ribs, each of their cartilages makes a greater curve in its progreſs from the rib towards the ſternum; and the tubercles, that are articulated to the tranſverſe proceſſes of the vertebrae, have their ſmooth ſurfaces gradually facing more upwards.—The ribs becoming thus more oblique, while the ſternum advances forwards in its deſcent, makes the diſtance between the ſternum and the anterior end of the lower ribs greater than between the ſternum and the ribs above; conſequently the cartilages of thoſe ribs that are joined to the breaſt-bone are longer in the lower than in the higher ones. —Theſe cartilages are placed nearer to each other as the ribs deſcend, which occaſions the curvature of the cartilages to be greater.

The length of the ribs increaſes from the firſt and uppermoſt rib, as far down as the ſeventh; and from that to the twelfth, it gradually diminiſhes.—The ſuperior of the two plain, or rather hollow ſurfaces, by which the ribs are articulated to the bodies of the vertebrae, gradually increaſe from the firſt to the fourth rib, and is diminiſhed after that in each lower rib.—The diſtance of their angles from the heads always increaſes as they deſcend to the ninth, becauſe of the greater breadth of the ſacrolumbalis muſcle (*s*).

The ribs are commonly divided into *true* and *falſe*.

The *true* (*t*) coſtae are the ſeven upper ones of each ſide, whoſe cartilages are all gradually longer as the ribs deſcend, and are joined to the breaſt-bone: So that, being preſſed conſtantly between two bones, they are flatted at both ends; and are thicker, harder, and more liable to oſſify, than the other cartilages that are not ſubject to ſo much preſſure. Theſe ribs include the heart and lungs: and therefore are the proper or true *cuſtodes* of life.

The

(*s*) Winſlow, Expoſition anatomique des os ſecs, § 643.

(*t*) Γνησιαι, Germanæ, legitimæ.

The five inferior ribs of each ſide are the *falſe* or *baſtard* (*u*), whoſe cartilages do not reach to the ſternum; and therefore, wanting the reſiſtance at their fore-part, they are there pointed; and on this account, having leſs preſſure, their ſubſtance is ſofter.—The cartilages of theſe falſe ribs are ſhorter as the ribs deſcend.—To all theſe five ribs the circular edge of the diaphragm is connected: And its fibres, inſtead of being ſtretched immediately tranſverſely, and ſo running perpendicular to the ribs, are preſſed ſo as to be often, eſpecially in exſpiration, parallel to the plane in which the ribs lie: Nay, one may judge by the attachments which theſe fibres have ſo frequently to the ſides of the thorax a conſiderable way above where their extremities are inſerted into the ribs, and by the ſituation of the viſcera always to be obſerved in a dead ſubject laid ſupine, that there is conſtantly a large concavity formed on each ſide by the diaphragm within theſe baſtard ribs, in which the ſtomach, liver, ſpleen, &c. are contained; which being only reckoned among the viſcera naturalia, have occaſioned the name of *baſtard cuſtodes* to theſe bones.

Hence, in ſimple fractures of the falſe-ribs, without fever, the ſtomach ought to be kept moderately filled with food, leſt the pendulous ribs falling inwards, ſhould thereby increaſe the pain, cough, &c. (*x*).——Hence likewiſe we may learn how to judge better of the ſeat of ſeveral diſeaſes, and to do the operation of the empyema, and ſome others, with more ſafety than we can do if we follow the common directions.

The eight upper ribs were formerly (*y*) claſſed into pairs, with particular names to each two; to wit, the *crooked*, the *ſolid*,

(*u*) Μαλθακαι, χονδροιδεις, ακανθαι, κλινες, ῥοαι, Adultrinae, ſpuriae, illegitimae.

(*x*) Hippocrat. de articulo, § 51.—Paré, lib. 15. cap. 11.

(*y*) Laurent. Hiſt. Anat. lib. 2. cap. 29.—Paaw, de oſſibus, pars 3. cap 2.

ſolid, the *pectoral*, the *twiſted*: But theſe names are of ſo little uſe, that they are now generally neglected.

The firſt rib of each ſide is ſo ſituated, that the flat ſides are above and below, while one edge is placed inwards and the other outwards, or nearly ſo: Therefore ſufficient ſpace is left above it for the ſubclavian veſſels and muſcles; and the broad concave ſurface is oppoſed to the lungs. But then, in conſequence of this ſituation, the channel for the intercoſtal veſſels is not to be found; and the edges are differently formed from all the other, except the ſecond; the lower one being rounded, and the other ſharp.—The head of this rib is not divided into two plain ſurfaces by a middle ridge, becauſe it is only articulated with the firſt vertebra of the thorax.—Its cartilage is oſſified in adults, and is united to the ſternum at right angles.—This firſt rib frequently has a ridge riſing near the middle of its poſterior edge, where one of the heads of the ſcalenus muſcles riſes.—Farther forward it is flatted, or ſometimes depreſſed by the clavicle.

The fifth, ſixth, and ſeventh, or rather the ſixth, ſeventh, eighth, and ſometimes the fifth, ſixth, ſeventh, eighth, and ninth ribs, have their cartilages at leaſt contiguous: And they are frequently joined to each other by croſs cartilages; and moſt commonly the cartilages of the eighth, ninth, and tenth, are connected to the former and to each other by firm ligaments.

The eleventh, and ſometimes the tenth rib, has no tubercle for its articulation with the tranſverſe proceſs of the vertebra, to which it is only looſely fixed by ligaments.—— The foſſa in its lower edge is not ſo deep as in the upper ribs, becauſe the veſſels run more towards the interſtice between the ribs.—Its fore-end is ſmaller than its body, and its ſhort ſmall cartilage is but looſely connected to the cartilage of the rib above.

The

The twelfth rib is the ſhorteſt and ſtraighteſt.—Its head is only articulated with the laſt vertebra of the thorax; therefore is not divided into two ſurfaces.—This rib is not joined to the tranſverſe proceſs of the vertebra; and therefore has no tubercle, being often pulled neceſſarily inwards by the diaphragm, which an articulation with the tranſverſe proceſs would not have allowed.—The foſſa is not found at its under edge, becauſe the veſſels run below it.—The fore-part of this rib is ſmaller than its middle, and has only a very ſmall pointed cartilage fixed to it.—To its whole internal ſide the diaphragm is connected.

The motions and uſes of the ribs ſhall be more particularly treated of after the deſcription of the ſternum.

The heads and tubercles of the ribs of a new-born child have cartilages on them; part of which becomes afterwards thin epiphyſes.——The bodies of the ribs encroach gradually after birth upon the cartilages; ſo that the latter are proportionally ſhorter, when compared to the ribs, in adults than in children.

Here I cannot help remarking the wiſe Providence of our Creator, in preſerving us from periſhing as ſoon as we come into the world. The end of the bones of the limbs remain in a cartilaginous ſtate after birth, and are many years before they are entirely united to the main body of their ſeveral bones; whereas the condyles of the occipital bone, and of the lower jaw, are true original proceſſes, and oſſified before birth; and the heads and tubercles of the ribs are nearly in the ſame condition: And therefore the weight of the large head is firmly ſupported; the actions of ſucking, ſwallowing, reſpiration, &c. which are indiſpenſably neceſſary for us as ſoon as we come into the world, are performed without danger of ſeparating the parts of the bones that are moſt preſſed on in theſe motions: Whereas, had theſe proceſſes of the head, jaw, and ribs, been epiphyſes at birth, children muſt

have been expoſed to danger of dying by ſuch a ſeparation; the immediate conſequences of which would be the compreſſion of the beginning of the ſpinal marrow, or want of food, or a ſtop put to reſpiration.

THE STERNUM.

The ſternum (z), or breaſt-bone, is the broad flat bone, or pile of bones, at the fore-part of the thorax. The number of bones into which this ſhould be divided, has occaſioned debates among anatomiſts, who have conſidered it in ſubjects of different ages.—In adults of a middle age, it is compoſed of three bones, which eaſily ſeparate after the cartilages connecting them are deſtroyed. The two lower bones are frequently found intimately united; and very often, in old people, the ſternum is a continued bony ſubſtance from one end to the other; though we ſtill obſerve two, ſometimes three, tranſverſe lines on its ſurface; which are marks of the former diviſions.

When we conſider the ſternum as one bone, we find it broadeſt and thickeſt above, and becoming ſmaller as it deſcends. The internal ſurface of this bone is ſomewhat hollowed for enlarging the thorax: But the convexity on the external ſurface is not ſo conſpicuous, becauſe the ſides are preſſed outwards by the true ribs; the round heads of whoſe cartilages are received into ſeven ſmooth pits formed in each ſide of the ſternum, and are kept firm there by ſtrong ligaments, which, on the external ſurface, have a particular radiated texture (a). The cartilaginous fibres frequently thruſt themſelves into the bony ſubſtance of the ſternum, and are joined by a ſort of ſuture.—The pits at the upper-part of the ſternum are at the greateſt diſtance one from another, and,

as

(z) Στηθος, Os pectoris, enſiforme, ſcutum cordis.

(a) Ruyſch, Catalog. Rar. fig. 9.

as they descend, are nearer; so that the two lowest are contiguous.

The *substance* of the breast-bone is cellular, with a very thin external plate, especially on its internal surface, where we may frequently observe a cartilaginous crust spread over it (*b*). —On both surfaces, however, a strong ligamentous membrane is closely braced; and the cells of this bone are so small, that a considerable quantity of osseous fibres must be employed in the composition of it. Whence, with the defence which the muscles give it, and the moveable support it has from the cartilages, it is sufficiently secured from being broken: For it is strong by its quantity of bone; its parts are kept together by ligaments; and it yields enough to elude considerably any violence offered (*c*).

So far may be said of this bone in *general*; but the *three* bones, of which, according to the common account, it is composed in adults, are each to be examined.

The *first*, all agree, is somewhat of the figure of a heart, as it is commonly painted; only it does not terminate in a sharp point.—This is the uppermost thickest part of the sternum.

The upper middle part of this first bone, where it is thickest, is hollowed, to make place for the trachea arteria, though this cavity (*d*) is principally formed by the bone being raised on each side of it, partly by the clavicles thrusting it inwards, and partly by the sterno-mastoidei muscles pulling it upwards.—On the outside of each tubercle there is an oblong cavity, that, in viewing it transversely from before backwards, appears a little convex. Into these glenae, the ends of the clavicle are received.—Immediately below these, the sides of this bone begin to turn thinner; and in each a superficial cavity

(*b*) Jac. Sylv. in Galen de ossibus, cap. 12.

(*c*) Senac, in Memoires de l'acad. des sciences, 1724.

(*d*) Σφαγη, Jugulum, furcula superior.

cavity or a rough ſurface is to be ſeen, where the firſt ribs are received or joined to the ſternum.—In the ſide of the under end of this firſt bone, the half of the pit for the ſecond rib on each ſide is formed.——The upper part of the ſurface behind is covered with a ſtrong ligament, which ſecures the clavicles; and is afterwards to be more particularly taken notice of.

The *ſecond* or middle diviſion of this bone is much longer, narrower, and thinner, than the firſt; but excepting that it is a little narrower above than below, it is nearly equal all over in its dimenſions of breadth or thickneſs.—In the ſides of it are complete pits for the third, fourth, fifth, and ſixth ribs, and an half of the pits for the ſecond and ſeventh; the lines, which are marks of the former diviſion of this bone, being extended from the middle of the pits of one ſide to the middle of the correſponding pits of the other ſide.—Near its middle an unoſſified part of the bone is ſometimes found; which, freed of the ligamentous membrane or cartilage that fills it, is deſcribed as a hole: And in this place, for the moſt part, we may obſerve a tranſverſe line, which has made authors divide this bone into two.—When the cartilage between this and the firſt bone is not offified, a manifeſt motion of this upon the firſt may be obſerved in reſpiration, or in raiſing the ſternum, by pulling the ribs upwards, or diſtending the lungs with air in a recent ſubject.

The *third* bone is much leſs than the other two, and has only one half of the pit for the ſeventh rib formed in it; wherefore it might be reckoned only an appendix of the ſternum.——In young ſubjects it is always cartilaginous, and is better known by the name of *cartilago xiphoides* or *enſiformis* (*e*) than any other; though the antients often called

(*e*) Clypealis, gladialis, mucronata, malum granatum, ſcutum ſtomachi, epiglottalis, cultralis, medium furculae inferioris, ſcutiformis, enſiculata.

called the whole ſternum *enſiforme*; comparing the two firſt bones to the handle, and this appendix to the blade of a ſword.———This third bone is ſeldom of the ſame figure, magnitude, or ſituation, in any two ſubjects; for ſometimes it is a plain triangular bone, with one of the angles below, and perpendicular to the middle of the upper ſide, by which it is connected to the ſecond bone.———In other people the point is turned to one ſide, or obliquely forwards or backwards. Frequently it is all nearly of an equal breadth, and in ſeveral ſubjects it is bifurcated; whence ſome writers give it the name of *furcilla or furcula inferior*; or elſe it is unoſſified in the middle.———In the greateſt number of adults it is oſſified, and tipped with a cartilage; in ſome, one half of it is cartilaginous; and in others, it is all in a cartilaginous ſtate.———Generally ſeveral oblique ligaments, fixed at one end to the cartilages of the ribs, and by the other to the outer ſurface of the xiphoid bone, connect it firmly to thoſe cartilages *(f)*.

So many different ways this ſmall bone may be formed without any inconvenience: But then ſome of theſe poſitions may be ſo directed, as to bring on a great train of ill conſequences; particularly when the lower end is oſſified, and is too much turned outwards or inwards *(g)*, or when the conjunction of this *appendix* with the ſecond bone is too weak *(h)*.

The ſternum is *joined* by cartilages to the ſeven upper ribs unleſs when the firſt coaleſces with it in an intimate union of ſubſtance; and its unequal cavity on each ſide of its upper end is fitted for the ends of the clavicles.

The

(f) Weitbrecht, Syndeſmolog. p. 121.

(g) Rolfinc. Differt. Anat. lib. 2. cap. 41.——Paaw de oſſib. pars 1. cap. 3. & pars 3. cap. 3.———Codronchi de prolapſu cartilagin. mucronat.

(h) Paaw de oſſib. pars 1. cap. 3. & pars 3. cap. 3.—Bartich. Act. Haſn. vol. 5. ob. 79.---Donet. Sepulchret. Anat. tom. 2. lib. 3. § 5. Append. ad. obſ. 8. et ibid. § 7. obſ. 19.

The ſternum moſt frequently has four ſmall round bones, ſurrounded with cartilage, in children born at the full time; the uppermoſt of theſe, which is the firſt bone, being the largeſt.———Two or three other very ſmall bony points are likewiſe to be ſeen in ſeveral children.———The number of bones increaſes for ſome years, and then diminiſhes, but uncertainly, till they are at laſt united into thoſe above deſcribed of an adult.

The *uſes* of this bone are, to afford origin and inſertion to ſeveral muſcles; to ſuſtain the mediaſtinum; to defend the vital organs, the heart and lungs, at the fore part; and, laſtly, by ſerving as a moveable fulcrum of the ribs, to aſſiſt conſiderably in reſpiration: which action, ſo far as it depends on the motion of the bones, we are now at liberty to explain.

When the ribs that are connected by their cartilages to the ſternum, or to the cartilages of the true ribs, are acted upon by the intercoſtal muſcles, they muſt all be pulled from the oblique poſition which their cartilages kept them in, nearer to right angles with the vertebrae and ſternum, becauſe the firſt or uppermoſt rib is by much the moſt fixed of any; and the cartilages making a greater reſiſtance to raiſing the anterior ends of the ribs, their large arched middle parts turn outwards as well as upwards.———The ſternum, preſſed ſtrongly on both ſides by the cartilages of the ribs, is puſhed forwards, and that at its ſeveral parts, in proportion to the length and motion of its ſupporters the ribs; that is, moſt at its lower end.———The ſternum and the cartilages, thus raiſed forwards, muſt draw the diaphragm connected to them: conſequently ſo far ſtretch it, and bring it nearer to a plane. The power that raiſes this bone and the cartilages, fixes them ſufficiently to make them reſiſt the action of the diaphragm, whoſe fibres contract at the ſame time, and thruſt the viſcera of the abdo-

men

men downwards.——The arched part of the ribs being thus moved outwards, their anterior ends and the ſternum being advanced forwards, and the diaphragm being brought nearer to a plain ſurface, inſtead of being greatly convex on each ſide within each cavity of the thorax, it is evident how conſiderably the cavity, of which the nine or ten upper ribs are the ſides, muſt be widened, and made deeper and longer. While this is doing in the upper ribs, the lower ones, whoſe cartilages are not joined to the ſternum or to other cartilages, move very differently, though they conſpire to the ſame intention, the enlargement of the thorax: for having no fixed point to which their anterior ends are faſtened, and the diaphragm being inſerted into them at the place where it runs nearly ſtraight upwards from its origin at the vertebrae, theſe ribs are drawn downwards by this ſtrong muſcle, and by the muſcles of the abdomen, which at this time reſiſt the ſtretching force of the bowels; while the intercoſtal muſcles are pulling them in the contrary direction, to wit, upwards. The effect, therefore, of either of theſe powers, which are antagoniſts to each other, is very little, as to moving the ribs either up or down; but the muſcles of the abdomen, puſhed at this time outwards by the viſcera, carry theſe ribs along with them.——Thus the thorax is not only not allowed to be ſhortened, but is really widened at its lower part, to aſſiſt in making ſufficient ſpace for the due diſtention of the lungs.

As ſoon as the action of theſe ſeveral muſcles ceaſes, the elaſtic cartilages, extending themſelves to their natural ſituations, depreſs the upper ribs, and the ſternum ſubſides; the diaphragm is thruſt up by the viſcera abdominalia, and the oblique and tranſverſe muſcles of the belly ſerve to draw the inferior ribs inwards at the ſame time.——By theſe cauſes, the cavity of the breaſt is diminiſhed in all its dimenſions.

Though

Though the motions above described of the ribs and sternum, especially of the latter bone, are so small in the mild respiration of a healthy person, that we can scarce observe them; yet they are manifest whenever we designedly increase our respiration, or are obliged to do it after exercise, and in several diseases.

SECT. III.

OF THE SUPERIOR EXTREMITIES.

AUTHORS are much divided in their opinions about the number of bones of which each superior extremity (*i*) should be said to consist; some describing the clavicle and scapula as part of it, others classing these two bones with those of the thorax: But since most quadrupeds have no clavicles, and the human thorax can perform its functions right when the scapula is taken away (*k*), while it is impossible for us to have the right use of our arms without these bones, I must think that they belong to the superior extremities. Each of the superior extremities may be divided into the shoulder, arm, fore-arm, and hand.

§ 1. *Bones of the* SHOULDER.

The SHOULDER consists of the *clavicle* and *scapula*.

CLAVICULA.

CLAVICULA, or *collar bone* (*l*), is the long crooked bone, in figure like an Italic *f*, placed almost horizontally between

(*i*) Κωλα, γυια, εκφυαδες, Enata, adnata, explanata membra, artus.

(*k*) Philosoph. Transact. numb. 449. § 5.

(*l*) Os jugulare, jugulum, furcula, ligula, clavis, humerus quibusdam.

between the upper lateral part of the ſternum and what is commonly called the top of the ſhoulder; which, as a clavis or beam, it bears off from the trunk of the body.

The clavicle, as well as other long round bones, is larger at its two ends than in the middle. The end next to the ſternum (*m*) is triangular: The angle behind is conſiderably protruded, to form a ſharp ridge, to which the tranſverſe ligament, extended from one clavicle to the other, is fixed (*n*). The ſide oppoſite to this is ſomewhat rounded. The middle of this protuberant end is as irregularly hollowed as the cavity in the ſternum for receiving it is raiſed: But, in a recent ſubject, the irregular concavities of both are ſupplied by a moveable cartilage; which is not only much more cloſely connected every where by ligaments to the circumference of the articulation than thoſe of the lower jaw are, but it grows to the two bones at both its internal and external end; its ſubſtance at the external end being ſoft, but very ſtrong, and reſembling the intervertebral cartilages (*o*).

From this internal end, the clavicle, for about two fifths of its length, is bended obliquely forwards and downwards. On the upper and fore-part of this curvature a ſmall ridge is ſeen, with a plain rough ſurface before it; whence the muſculus ſterno-hyoideus and ſterno maſtoideus have in part their origin. Near the lower angle a ſmall plain ſurface is often to be remarked, where the firſt rib and this bone are contiguous (*p*), and are connected by a firm ligament (*q*). From this a rough plain ſurface is extended outwards, where

(*m*) Παραφαγις.

(*n*) Riolan. Encheirid. anat. lib. 6. cap. 13.——Winſlow, Expoſ. Anat. des os frais, § 248.——Weitbrecht. Act. Petropolit. tom. 4. p. 255. et Syndeſmolog. ſect. 2. I. §. 3.

(*o*) Weitbrecht. Syndeſmolog. ſect. 2. I. § 6.

(*p*) Dionis, Sixieme demonſt. des os.

(*q*) Weitbrecht. Syndeſmolog. ſect. 2. I. § 7.

the pectoral muſcle has part of its origin. Behind, the bone is made flat and rough by the inſertion of the larger ſhare of the ſubclavian muſcle. After the clavicle begins to be bent backwards, it is round: But it ſoon after becomes broad and thin; which ſhape it retains to its external end. Along the external concavity, a rough inequality runs, from which ſome part of the deltoid muſcle takes its riſe; oppoſite to this, on the convex edge, a ſcabrous ridge gives inſertion to a ſhare of the cucullaris muſcle. The upper ſurface of the clavicle is here flat; but the lower is hollow, for lodging the beginning of the muſculus ſubclavius; and towards its back part a tubercle riſes; to which, and to a roughneſs near it, the ſtrong ſhort thick ligament, connecting this bone to the coracoid proceſs of the ſcapula, is fixed.

The external end (*r*) of this bone is horizontally oblong, ſmooth, ſloping at the poſterior ſide, and tipped in a recent ſubject with a cartilage, for its articulation with the acromion ſcapulae. Round this the bone is ſpongy, for the firmer connection of the ligaments.

The medullary arteries having their direction obliquely outwards, enter the clavicles by one or more ſmall paſſages in the middle of their back-part.

The *ſubſtance* of this bone is the ſame as of the other round long bones.

The triangular unequal interior end of each clavicle has the cartilage above deſcribed interpoſed betwen it and the irregular cavity of the ſternum —The ligaments which ſurround this articulation to ſecure it, are ſo ſhort and ſtrong, that little motion can be allowed any way; and the ſtrong ligament that is ſtretched acroſs the upper furcula of the ſternum, from the poſterior prominent angle of the one clavicle to

(*r*) Ἔσχατος.

to the ſame place of the other clavicle, ſerves to keep each of theſe bones more firmly in their place.——By the aſſiſtance, however, of the moveable intervening cartilage, the clavicle can, at this joint, be raiſed or depreſſed, and moved backwards and forwards ſo much as that the external end, which is at a great diſtance from that axis, enjoys very conſpicuous motions. The articulation of the exterior end of the clavicle ſhall be conſidered after the deſcription of the ſcapula.

The clavicles of infants are not deficient in any of their parts; nor have they any epiphyſes at their extremities joined afterwards to their bodies, as moſt other ſuch long bones have, which preſerve them from being bent too much, and from the danger of any unoſſified parts being ſeparated by the force which pulls the arms forwards.

The *uſes* of the clavicles are, to keep the ſcapulae, and conſequently all the ſuperior extremities, from falling in and forward upon the thorax; by which, as in moſt quadrupeds, the motions of the arms would be much confined, and the breaſt made too narrow. The clavicles likewiſe afford origin to ſeveral muſcles, and a defence to large veſſels.

From the ſituation, figure, and uſe of the clavicles, it is evident that they are much expoſed to fractures; and their broken parts muſt generally paſs each other; and that they are difficultly kept in their place afterwards.

SCAPULA.

Scapula, or *ſhoulder-blade* (*s*), is the triangular bone ſituated on the outſide of the ribs: with its longeſt ſide, called its *baſe*, towards the ſpinal proceſſes of the vertebrae; and with

(*s*) Ωμοπλατης, πτερυγιον. Latitudo humeri, ſceptulum vel ſcutulum opertum, ſpatula, ala, humerus, clypeus, ſcutum thoracis.

with the angle at the upper part of this ſide about three inches, and the lower angle at a greater diſtance, from theſe proceſſes. The back-part of the ſcapula has nothing but the thin ends of the ſerratus anticus major and ſubſcapularis muſcles between it and the ribs: But as this bone advances forwards, its diſtance from the ribs increaſes. The upper or ſhorter ſide, called the *ſuperior coſta*, of the ſcapula, is nearly horizontal, and parallel with the ſecond rib. The lower ſide, which is named the *inferior coſta*, is extended obliquely from the third to the eighth rib. The ſituation of this bone, here deſcribed, is when people are ſitting or ſtanding in a ſtate of inactivity, and allowing the members to remain in the moſt natural eaſy poſture. The inferior angle of the ſcapula is very acute; the upper one is near to a right angle; and what is called the anterior does not deſerve the name, for the two ſides do not meet to form an angle. The body of this bone is concave towards the ribs, and convex behind, where it has the name of *dorſum* (*t*). Three proceſſes are generally reckoned to proceed from the ſcapula. The firſt is the large ſpine that riſes from its convex ſurface behind, and divides it unequally. The ſecond proceſs ſtands out from the fore part of the upper ſide: And, from its imaginary reſemblance to a crow's beak, is named *coracoides* (*u*). The third proceſs is the whole thick bulbous fore-part of the bone.

After thus naming the ſeveral conſtituent parts of the ſcapula, the particular deſcription will be more eaſily underſtood.

The baſe, which is tipped with cartilage in a young ſubject, is not all ſtraight: For above the ſpine it runs obliquely forwards to the ſuperior angle, that here it might not be too protuberant backwards, and ſo bruiſe the muſcles

(*t*) Χελωνιον.

(*u*) Anchoroides, ſigmoides, digitalis, anciſtroides.

cles and teguments. Into the oblique ſpace the muſculus patientiae is inſerted. At the root of the ſpine, on the back-part of the baſe, a triangular plain ſurface is formed by the preſſure of the lower fibres of the trapezius. Below this the edge of the ſcapula is ſcabrous and rough, for the inſertion of the ſerratus major anticus and rhomboid muſcles.

The back-part of the inferior angle is made ſmooth by the latiſſimus dorſi paſſing over it. This muſcle alſo alters the direction of the inferior coſta ſome way forwards from this angle: and ſo far it is flatted behind by the origin of the teres major. As the inferior coſta advances forward, it is of conſiderable thickneſs, it is ſlightly hollowed and made ſmooth behind by the teres minor, while it has a foſſa formed into it below by part of the ſubſcapularis; and between the two a ridge with a ſmall depreſſion appears, where the longus extenſor cubiti has its origin.

The ſuperior coſta is very thin: and near its fore-part there is a ſemilunar niche, from one end of which to the other a ligament is ſtretched; and ſometimes the bone is continued to form one, or ſometimes two, holes for the paſſage of the ſcapular blood-veſſels and nerves. Immediately behind this ſemilunar cavity the coraco-hyoid muſcle has its riſe. From the niche to the termination of the foſſa for the teres minor, the ſcapula is narrower than any where elſe, and ſupports the third proceſs. This part has the name of *cervix*.

The whole dorſum of the ſcapula is always ſaid to be convex; but, by reaſon of the raiſed edges that ſurround it, it is divided into two cavities by the ſpine, which is ſtretched from behind forwards, much nearer to the ſuperior than to the inferior coſta. The cavity above the ſpine is really concave where the ſupra-ſpinatus muſcle is lodged; while the ſurface of this bone below the ſpine, on which the infra-ſpinatus

natus muſcle is placed, is convex, except a foſſa that runs at the ſide of the inferior coſta.

The internal or anterior ſurface of this bone is hollow, except in the part above the ſpine, which is convex.——The ſubſcapularis muſcle is extended over this ſurface, where it forms ſeveral ridges and intermediate depreſſions, commonly miſtaken for prints of the ribs; they point out the interſtices of the bundles of fibres of which the ſubſcapularis muſcle is compoſed (*x*).

The ſpine (*y*) riſes ſmall at the baſe of the ſcapula, and becomes higher and broader as it advances forwards.——On the ſides it is unequally hollowed and crooked, by the actions of the adjacent muſcles.——Its ridge (*z*) is divided into two rough flat ſurfaces: Into the upper one, the trapezius muſcle is inſerted; and the lower one has part of the deltoid fixed to it.——The end of the ſpine, called *acromion* (*a*), or top of the ſhoulder, is broad and flat, and is ſometimes only joined to the ſpine by a cartilage (*b*).——The anterior edge of the acromion is flat, ſmooth, and covered with a cartilage, for its articulation with the external end of the clavicle; and it is hollowed below, to allow a paſſage to the infra and ſupra ſpinati muſcles, and free motion to the os humeri.

The coracoid (*c*) proceſs is crooked, with its point inclining forwards; ſo that a hollow is left at the lower ſide of its root, for the paſſage of the infra ſcapularis muſcle.——The end of this proceſs is marked with three plain ſurfaces. Into the internal, the ſerratus minor anticus is inſerted;

(*x*) Winſlow, in Memoires de l'acad. des ſciences, 1722.

(*y*) Ραχις, υπερoχη ωμοπλατων, Eminentia ſcapularum.

(*z*) Ptergium, criſta.

(*a*) Εσωμις, ακρωμιον, κορακοειδης, κατακλεις, Acromii os, ſummus armus, roſtrum porcinum, proceſ. digitalis.

(*b*) Sue, Trad. d'oſteol. p. 160.

(*c*) Ακρωμιοειδης, σιγμοειδης, Roſtriformis.

inſerted: from the external, one head of the biceps flexor cubiti riſes; and from the lower one, the coraco-brachialis has its origin. At the upper part of the root of this proceſs, immediately before the ſemilunar cavity, a ſmooth tubercle appears, where a ligament from the clavicle is fixed. From all the external ſide of this coracoid apophyſe, a broad ligament goes out, which becomes narrower where it is fixed to the acromion. The ſharp pain, violent inflammation, and tedious cure of contuſions in this part, are probably owing to theſe tendons and ligaments being hurt.

From the cervix ſcapulae the third proceſs is produced. The fore-part of this is formed into a glenoid cavity (*d*), which is of the ſhape of the longitudinal ſection of an egg, being broad below and narrow above. Between the brims of this hollow and the fore-part of the root of the ſpine, a large ſinuoſity is left for the tranſmiſſion of the ſupra and infra ſpinati muſcles; and on the upper part of theſe brims we may remark a ſmooth ſurface, where the ſecond head of the biceps flexor cubiti has its origin. The root of the ſupercilia is rough all round, for the firmer adhe ion of the capſular ligament of the articulation, and of the cartilage which is placed on theſe brims, where it is thick, but becomes very thin as it is continued towards the middle of the cavity, which it lines all over.

The medullary veſſels enter the ſcapula near the baſe of the ſpine.

The *ſubſtance* of the ſcapula, as in all other broad flat bones, is cellular, but of an unequal thickneſs: for the neck and third proceſs are thick and ſtrong; the inferior coſta, and ſpine, and coracoid proceſs, are of a middle thickneſs; and the body is ſo preſſed by the muſcles, as to become thin and diaphanous.

(*d*) Ωμοχοτυλης.

The ſcapula and clavicle are *joined* by plain ſurfaces, tipped with cartilage (*e*); by which neither bone is allowed any conſiderable motion, being tightly tied down by the common capſular ligament, and by a very ſtrong one which proceeds from the coracoid proceſs; but divides into two before it is fixed into the clavicle, with ſuch a direction, as can either allow this bone to have a ſmall rotation, in which its poſterior edge turns more backwards, while the anterior one riſes farther forwards; or it can yield to the fore-part of the ſcapula moving downwards, while the back-part of it is drawn upwards; in both which caſes, the oblong ſmooth articulated ſurfaces of the clavicle and ſcapula are not in the ſame plane, but ſtand a little tranſverſely, or acroſs each other, and thereby preſerve this joint from luxations, to which it would be ſubject if either of the bones was to move on the other perpendicularly up and down without any rotation.——Sometimes a moveable ligamentous cartilage is found in this joint; and ſometimes ſuch a cartilage is only interpoſed at the anterior half of it: and in ſome old ſubjects I have found a ſeſamoid bone here (*f*). The ſcapula is connected to the head, os hyoides, vertebrae, ribs, and arm-bone, by muſcles, that have one end faſtened to theſe bones, and the other to the ſcapula, which can move it upwards, downwards, backwards, or forwards; by the quick ſucceſſion of theſe motions, its whole body is carried in a circle. But being alſo often moved as upon an axis perpendicular to its plane, its circumference turns in a circle whoſe centre this axis is (*g*). Whichever of theſe motions it performs, it always carries the outer end of the clavicle and the arm along with it.——The glenoid cavity of this bone receives the os humeri,

(*e*) Acromion, *κατακλεις*, Clauſurae.

(*f*) Jac. Sylv. Iſagog. Anat. lib. 1. cap. 2.

(*g*) See Winſlow, Memoires de l'acad. des ſciences, 1726.

humeri, which plays in it as a ball in a ſocket, as will be more fully explained hereafter.

The *uſe* of the ſcapula is, to ſerve as a fulcrum to the arm; and, by altering its poſition on different occaſions, to allow always the head of the os humeri a right ſituated ſocket to move in: And thereby to aſſiſt and to enlarge greatly the motions of the ſuperior extremity, and to afford the muſcles which riſe from it more advantageous actions, by altering their direction to the bone which they are to move.——This bone alſo ſerves to defend the back-part of the thorax, and is often employed to ſuſtain weights, or to reſiſt forces, too great for the arm to bear.

The baſe, acromion, coracoid proceſs, and head of the ſcapula, are all in a cartilaginous ſtate at birth; and the three firſt are joined as epiphyſes; while the head, with the glenoid cavity, is not formed into a diſtinct ſeparate bone, but is gradually produced by the oſſification of the body of this bone being continued forwards.

§ 2. *The* ARM.

The *Arm* has only one bone, beſt known by the Latin name of *os humeri* (*h*); which is long, round, and nearly ſtraight.

The upper end of this bone (*i*) is formed in a large round ſmooth head, whoſe middle point is not in a ſtraight line with the axis of the bone, but ſtands obliquely backwards from it.—The extent of the head is diſtinguiſhed by a circular foſſa ſurrounding its baſe, where the head is united to the bone, and the capſular ligament of the joint is fixed.—Below the fore-part of its baſe, two tubercles ſtand out: The

(*h*) Ακρώλια, ωλενη, Os brachii, armi, adjutorium, parvum, brachium, canna brachii.

(*i*) Acrocolium.

ſmalleſt one, which is ſituated moſt to the inſide, has the tendon of the ſubſcapularis muſcle inſerted into it.—The larger more external protuberance is divided, at its upper part, into three ſmooth plain ſurfaces; into the anterior of which, the muſculus ſupra-ſpinatus; into the middle or largeſt, the infra-ſpinatus; and into the one behind, the teres minor, is inſerted.—Between theſe two tubercles, exactly in the fore-part of the bone, a deep long foſſa is formed, for lodging the tendinous head of the biceps flexor cubiti; which, after paſſing, in a manner peculiar to itſelf, through the cavity of the articulation, is tied down by a tendinous ſheath extended acroſs the foſſa; in which, and in the neighbouring tubercles, are ſeveral remarkable holes, which are penetrated by the tendinous and ligamentous fibres, and by veſſels —— On each ſide of this foſſa, as it deſcends in the os humeri, a rough ridge, gently flatted in the middle, runs from the roots of the tubercles. The tendon of the pectoral muſcle is fixed into the anterior of theſe ridges, and the latiſſimus dorſi and teres major are inſerted into the internal one. A little behind the lower end of this laſt, another rough ridge may be obſerved, where the coraco-brachialis is inſerted. From the back-part of the root of the largeſt tubercle a ridge alſo is continued, from which the brevis extenſor cubiti riſes. This bone is flatted on the inſide, about its middle, by the belly of the biceps flexor cubiti. In the middle of this plain ſurface, the entry of the medullary artery is ſeen ſlanting obliquely downwards. At the fore-ſide of this plane the bone riſes in a ſort of a ridge, which is rough, and often has a great many ſmall holes in it, where the tendon of the ſtrong deltoid muſcle is inſerted; on each ſide of which the bone is ſmooth and flat, where the brachius internus riſes. The exterior of theſe two flat ſurfaces is the largeſt; behind it a ſuperficial ſpiral channel, formed by the muſcular nerve and the veſſels that accompany it, runs from behind

behind forwards and downwards. The body of the os humeri is flatted behind by the extenſors of the fore-arm.

Near the lower end of this bone, a large ſharp ridge is extended on its outſide, from which the muſculus ſpinator radii longus, and the longeſt head of the extenſor carpi radialis, riſe. Oppoſite to this, there is another ſmall ridge to which the aponeurotic tendon, that gives origin to the fibres of the internal and external brachial muſcles, is fixed; and from a little depreſſion on the fore-ſide of it, the pronator radii teres riſes.

The body of the os humeri becomes gradually broader towards the lower end, where it has ſeveral proceſſes; at the roots of which there is a cavity before and another behind (*k*). The anterior is divided by a ridge into two; the external, which is the leaſt, receives the end of the radius; and the internal receives the coronoid proceſs of the ulna in the flexions of the fore arm, while the poſterior deep triangular cavity lodges the olecranon in the extenſions of that limb. The bone between theſe two cavities is preſſed ſo thin by the proceſſes of the ulna, as to appear diaphanous in ſeveral ſubjects. The ſides of the poſterior cavity are ſtretched out into two proceſſes, one on each ſide: Theſe are called *condyles*; from each of which a ſtrong ligament goes out to the bones of the fore-arm. The external condyle, which has an oblique direction alſo forwards in reſpect of the internal, when the arm is in the moſt natural poſture (*l*), is equally broad, and has an obtuſe ſmooth head riſing from it forwards. From the rough part of the condyle, the inferior heads of the bicornis, the extenſor digitorum communis, extenſor carpi ulnaris, anconaeus, and ſome part of the ſpinator radii brevis, take their riſe; and on the ſmooth head the upper end of the radius plays. Immediately on the out-ſide of this, a

ſinuoſity

(*k*) Βαθμιδες. (*l*) Winſlow, Memoires de l'acad. des ſciences, 1722.

ſinuoſity is made by the ſhorter head of the bicornis muſcle, upon which the muſcular nerve is placed. The internal condyle is more pointed and protuberant than the external, to give origin to ſome part of the flexor carpi radialis, pronator radii teres, palmaris longus, flexor digitorum ſublimis, and flexor carpi ulnaris. Between the two condyles, is the trochlea or pulley; which conſiſts of two lateral protuberances, and a middle cavity, that are ſmooth and covered with cartilage. When the fore-arm is extended, the tendon of the internal brachiaeus muſcle is lodged in the fore part of the cavity of this pulley. The external protuberance, which is leſs than the other, has a ſharp edge behind; but forwards, this ridge is obtuſe, and only ſeparated from the little head, already deſcribed, by a ſmall foſſa, in which the joined edges of the ulna and radius move. The internal protuberance of the pulley is largeſt and higheſt; and therefore, in the motions of the ulna upon it, that bone would be inclined outwards, were it not ſupported by the radius on that ſide.——Between this internal protuberance and condyle, a ſinuoſity may be remarked, where the ulnar nerve paſſes.

The *ſubſtance* and the internal ſtructure of the os humeri is the ſame, and diſpoſed in the ſame way, as in other long bones.

The round head at the upper end of this bone is *articulated* with the glenoid cavity of the ſcapula; which being ſuperficial, and having long ligaments, allows the arm a free and extenſive motion.———Theſe ligaments are, however, conſiderably ſtrong. For beſides the common capſular one, the tendons of the muſcles perform the office, and have been deſcribed under the name of *ligaments*.——— Then the acromion and coracoid proceſs, with the ſtrong broad ligaments ſtretched between them, ſecure the articulation

lation above, where the greateſt and moſt frequent force is applied to thruſt the head of the bone out of its place. It is true, that there is not near ſo ſtrong a defence at the lower part of the articulation; but, in the ordinary poſtures of the arm, that is, ſo long as it is at an acute angle with the trunk of the body, there cannot be any force applied at this place to occaſion a luxation, ſince the joint is protected ſo well above.

The motions which the arm enjoys by this articulation are to every ſide; and by the ſucceſſion of theſe different motions, a circle may be deſcribed. Beſides which, the bone performs a ſmall rotation round its own axis. But though this can be performed with the round head in all poſitions; yet as theſe vary, the effects upon the body of the bone are very different: For if the middle of the head is the centre of rotation, as it is when the arm hangs down by the ſide, the body of the bone is only moved forwards and backwards; becauſe the axis of motion of the head it nearly at right angles with the length of the bone (*m*); whereas, when the arm is raiſed to right angles with the trunk of the body, the centre of motion, and the axis of the bone, come to be of the ſame ſtraight line: and therefore the body of the os humeri performs the ſame motion with its head. Though the motions of the arm ſeem to be extenſive, yet the larger ſhare of them depends on the motions of the ſcapula. The lower end of the os humeri is articulated with the bone of the fore-arm, and carries them with it in all its motions, but ſerves as a baſe on which they perform the motions peculiar to themſelves; as ſhall be deſcribed afterwards.

Both the ends of this bone are cartilaginous in a new-born infant; and the large head with the two tubercles, and

(*m*) Hippocrat. de articul. § 1.

and the trochlea with the two condyles, become epiphyſes before they are united to the body of the bone.

§ 3. *The* Fore-Arm.

The fore-arm (*n*) conſiſts of two long bones, the *ulna* and *radius*; whoſe ſituation, in reſpect of each other, is oblique in the leaſt ſtraining or moſt natural poſture; that is, the ulna is not directly behind, nor on the outſide of the radius, but in a middle ſituation between theſe two, and the radius croſſes it. The ſituation, however, of theſe bones, and of all the other bones of the ſuperior extremity that are not yet deſcribed, is frequently altered: and therefore, to ſhun repetitions, I deſire it may now be remarked, that, in the remaining account of the ſuperior extremity, I underſtand by the term of *poſterior*, that part which is in the ſame direction with the back of the hand: by *anterior*, that anſwering to the palm; by *internal*, that on the ſame ſide with the thumb; by *external*, the ſide neareſt to the little finger; ſuppoſing the hand always to be in a middle poſition between the pronation and ſupination.

ULNA.

Ulna (*o*), ſo named from its being uſed as a meaſure, is the longeſt of the two bones of the fore arm, and ſituated on the outſide of the radius.

At the upper end of the ulna are two proceſſes.——The poſterior is the largeſt, and formed like a hook, whoſe concave ſurface moves upon the pulley of the os humeri, and is called *olecranon* (*p*), or top of the cubit. The convex

(*n*) Cubitus, πηχυς. ωλενη, πυγων, Ulna, lacertus.

(*o*) Cubitus, πηχυς, προπηχιον, Focile majus, canna, vel arundo major, et inferior brachii

(*p*) Αγκων, Gibber cubitus, additamentum necatum.

vex back part of it is rough and ſcabrous, where the longus, brevis, and brachiaeus externus, are inſerted. The olecranon makes it unneceſſary that the tendons of the extenſor muſcles ſhould paſs over the end of the os humeri; which would have been of ill conſequence in the great flexions of this joint, or when any conſiderable force is applied to this part (*q*). The anterior proceſs is not ſo large, nor does it reach ſo high, as the one behind; but is ſharper at its end, and therefore is named *coronoid*.——Between theſe two proceſſes, a large ſemicircular or ſigmoid concavity is left; the ſurface of which on each ſide of a middle riſing, is ſlanting, and exactly adapted to the pulley of the bone of the arm.——Acroſs the middle of it, there is a ſmall ſinuoſity for lodging mucilaginous glands; where, as well as in a ſmall hollow on the internal ſide of it, the cartilage that lines the reſt of its ſurface is wanting.——Round the brims of this concavity the bone is rough, where the capſular ligament of the joint is implanted.——Immediately below the olecranon, on the back-part of the ulna, a flat triangular ſpongy ſurface appears, on which we commonly lean.——At the internal ſide of this, there is a larger hollow ſurface, where the muſculus anconaeus is lodged; and the ridge at the inſide of this gives riſe to the muſculus ſupinator radii brevis. Between the top of the ridge and the coronoid proceſs is the ſemilunated ſmooth cavity, lined with cartilage; in which, and in a ligament extended from the one to the other end of this cavity, the round head of the radius plays.——Immediately below it, a rough hollow gives lodging to mucilaginous glands.——Below the root of the coronoid proceſs, this bone is ſcabrous and unequal, where the brachiaeus internus is inſerted.——On the outſide of that, we obſerve a ſmooth concavity, where

(*q*) Winſlow, Expoſition anatomique du corps humain, traité des os ſecs, § 979.

where the beginning of the flexor digitorum profundus ſprouts out.

The body of the ulna is triangular.——The internal angle is very ſharp where the ligament that connects the two bones is fixed: The ſides which make this angle are flat and rough, by the action and adheſion of the many muſcles which are ſituated here. At the diſtance of one-third of the length of the ulna from the top, in its fore-part, the paſſage of the medullary veſſels may be ſeen ſlanting upwards. The external ſide of this bone is ſmooth, ſomewhat convex, and the angles at each edge of it are blunted by the preſſure of the muſcles equally diſpoſed about them.

As this bone deſcends, it becomes gradually ſmaller; ſo that its lower end terminates in a little head, ſtanding on a ſmall neck. Towards the fore but outer part of which laſt, an oblique ridge runs, that gives riſe to the pronator radii quadratus. The head is round, ſmooth, and covered with a cartilage on its internal ſide, to be received into the ſemilunar cavity of the radius; while a ſtyloid proceſs (*r*) riſes from its outſide, to which is fixed a ſtrong ligament that is extended to the os cuneiforme and piſiforme of the wriſt. Between the back-part of that internal ſmooth ſide and this proceſs, a ſinuoſity is left for the tendon of the extenſor carpi ulnaris. On the fore-part of the root of the proceſs, ſuch another depreſſion may be remarked for the paſſage of the ulnar artery and nerve. The end of the bone is ſmooth, and covered with a cartilage. Between it and the bones of the wriſt, a doubly concave moveable cartilage is interpoſed; which is a continuation of the cartilage that covers the lower end of the radius, and is connected looſely to the root of the ſtyloid proceſs, and to the rough cavity there; in which mucilaginous glands are lodged.

The

(*r*) Περονοειδη, Malleolus externus.

The ulna is articulated above with the lower end of the os humeri, where theſe bones have depreſſions and protuberances correſponding to each other, ſo as to allow an eaſy and ſecure extenſion of the fore-arm to almoſt a ſtraight line with the arm, and flexion to a very acute angle; but, by the ſlanting poſition of the pulley, the lower part of the fore-arm is turned outwards in the extenſion, and inwards in the flexion (*s*); and a very ſmall kind of rotation is likewiſe allowed in all poſitions, eſpecially when the ligaments are moſt relaxed by the fore-arm being in a middle degree of flexion. The ulna is alſo articulated with the radius and carpus, in a manner to be related afterwards.

RADIUS.

Radius (*t*), ſo called from its imagined reſemblance to a ſpoke of a wheel or to a weaver's beam, is the bone placed at the inſide of the fore-arm. Its upper end is formed into a circular little head, which is hollowed for an articulation with the tubercle at the ſide of the pulley of the os humeri; and the half of the round circumference of the head next to the ulna is ſmooth, and covered with a cartilage, in order to be received into the ſemilunated cavity of that bone.——Below the head, the radius is much ſmaller; therefore this part, which is made round by the action of the ſupinator radii brevis, is named its *cervix*.——At the external root of this neck, a tuberous proceſs riſes; into the outer part of which the biceps flexor cubiti is inſerted.——From this a ridge runs downwards and inwards, where the ſupinator radii brevis is inſerted; and a little below, and behind this ridge, there is a rough ſcabrous ſurface, where the pronator radii teres is fixed.

The body of the radius is not ſtraight, but convex on its

(*s*) Winſlow, Memoires de l'acad. des ſciences, 1722.

(*t*) Κερκις, παρατηχιον, Focile minus, canna minor, arundo minor.

its internal and poſterior ſurfaces; where it is alſo made round by the equal preſſure of the circumjacent muſcles, particularly of the extenſors of the thumb; but the ſurfaces next to the ulna are flatted and rough, for the origin of the muſcles of the hand; and both terminate in a common ſharp ſpine, to which the ſtrong ligament extended between the two bones of the fore-arm is fixed. A little below the beginning of the plain ſurface, on its fore-part, where the flexor muſcle of the laſt joint of the thumb takes its origin, the paſſage of the medullary veſſels is ſeen ſlanting upwards. The radius becomes broader and flatter towards the lower end, eſpecially on its fore-part, where its pronator quadratus muſcle is ſituated.

The lower end of the radius is larger than the ſuperior; though not in ſuch a diſproportion as the upper end of the ulna is larger than its lower end.——Its back-part has a flat ſtrong ridge in the middle, and foſſae on each ſide.——In a ſmall groove, immediately on the outſide of the ridge, the tendon of the extenſor tertii internodii pollicis plays. In a large one behind this, the tendons of the indicator and of the common extenſor muſcles of the fingers paſs.——Contiguous to the ulna there is a ſmall depreſſion made by the extenſor minimi digiti.——On the inſide of the ridge there is a broad depreſſion, which ſeems again ſubdivided, where the two tendons of the bicornis, or extenſor carpi radialis, are lodged.—the internal ſide of this end of the radius is alſo hollowed by the extenſors of the firſt and ſecond joint of the thumb; immediately above which a little rough ſurface ſhews where the ſupinator radii longus is inſerted.——The ridges at the ſides of the grooves, in which the tendons play, have an angular ligament fixed to them, by which the ſeveral ſheaths for the tendons are formed. The fore-part of this end of the radius is alſo depreſſed, where the flexors of the fingers and flexor carpi radialis paſs.——The external ſide is formed

into

into a ſemilunated ſmooth cavity, lined with a cartilage, for receiving the lower end of the ulna.——The loweſt part of the radius is formed into an oblong cavity: in the middle of which is a ſmall tranſverſe riſing, gently hollowed, for lodging mucilaginous glands; while the riſing itſelf is inſinuated into the conjunction of the two bones of the wriſt that are received into the cavity.——The internal ſide of this articulation is fenced by a remarkable proceſs (*u*) of the radius, from which a ligament going out to the wriſt, as the ſtyloid proceſs of the ulna with its ligament, guard it on the outſide.

The ends of both the bones of the fore-arm being thicker than the middle, there is a conſiderable diſtance between the bodies of theſe bones; in the larger part of which a ſtrong tendinous, but thin ligament, is extended, to give a large enough ſurface for the origin of the numerous fibres of the muſcles ſituated here, that are ſo much ſunk between the bones as to be protected from injuries, to which they would otherwiſe be expoſed. But this ligament is wanting near the upper end of the fore-arm, where the ſupinator radii brevis, and flexor digitorum profundus, are immediately connected (*x*).

Both ends of the bones of the fore-arm are firſt cartilages, and then epiphyſes, in children.

As the head of the radius receives the tubercle of the os humeri, it is not only bended and extended along with the ulna, but may be moved round its axis in any poſition; and, that this motion round its axis may be ſufficiently large, the ligament of the articulation is extended, further down than ordinary, on the neck of this bone, before it is connected to it: and it is very thin at its upper and lower part,

(*u*) Malleolus internus, proceſſus ſtyloides.

(*x*) Weitbrecht, Syndeſmolog. fig. 10, 11.

part, but makes a firm ring in the middle. This bone is alſo joined to the ulna by a double articulation: for above, a tubercle of the radius plays in a ſocket of the ulna; whilſt below, the radius gives the ſocket, and the ulna the tubercle. But then the motion performed in theſe two is very different: for, at the upper end, the radius does no more than turn round its axis; while, at the lower end, it moves in a ſort of cycloid upon the round part of the ulna; and as the hand is articulated and firmly connected here with the radius, they muſt move together.——When the palm is turned uppermoſt, the radius is ſaid to perform the *ſupination*: when the back of the hand is above, it is ſaid to be *prone*. But then the quickneſs and large extent of theſe two motions are aſſiſted by the ulna, which, as was before obſerved, can move with a kind of ſmall rotation on the ſloping ſides of the pulley. This lateral motion, though very inconſiderable in the joint itſelf, is conſpicuous at the lower end of ſuch a long bone; and the ſtrong ligament connecting this lower end to the carpus, makes the hand more readily obey theſe motions. When we deſign a large circular turn of our hand, we increaſe it by the rotation of the os humeri, and ſometimes employ the ſpine and inferior extremities to make theſe motions of pronation or ſupination of the hand large enough.

§ 4. *The* HAND.

THE hand *(y)* comprehends all from the joint of the wriſt to the points of the fingers. Its back-part is convex, for greater firmneſs and ſtrength; and it is concave before, for containing more ſurely and conveniently ſuch bodies as we take hold of. One half of the hand has an obſcure motion

(y) Ακρωτηριον, Summa manus.

motion in compariſon of what the other has, and ſerves as a baſe to the moveable half; which can be extended back very little farther than to a ſtraight line with the fore-arm, but can be conſiderably bent forwards.

As the bones that compoſe the hand are of different ſhapes and uſes, while ſeveral of them that are contiguous agree in ſome general characters; the hand is, on this account, commonly divided into carpus, metacarpus, and fingers; among which laſt the thumb is reckoned.

CARPUS.

THE carpus (z) is compoſed of eight ſmall ſpongy bones, ſituated at the upper part of the hand. I ſhall deſcribe each of theſe bones, under a proper name taken from their figure (a); becauſe the method of ranging them by numbers leaves anatomiſts too much at liberty to debate very idly, which ought to be preferred to the firſt number; or, what is worſe, ſeveral, without explaining the order they obſerve, apply the ſame numbers differently, and ſo confound their readers. But, that the deſcription of theſe bones may be in the ſame order as they are found in the generality of anatomical books, I ſhall begin with the range of bones that are concerned in the moveable joint of the wriſt, or are connected to the fore-arm, and ſhall afterwards conſider the four that ſupport the thumb and oſſa metacarpi of the fingers.

The eight bones of the CARPUS are, Os ſcaphoides, lunare, cuneiforme, piſiforme, trapezium, trapezoides, magnum, unciforme.

The *ſcaphoides* is ſituated moſt internally of thoſe that are articulated with the fore-arm.——The *lunare* immediately on

(z) Καρπὸς, Brachiale, prima palmae pars, raſetta.

(a) Lyſer, Cult. Anat. lib. 5. cap. 2.

on the outside of the former.——The *cuneiforme* is placed still more externally, but does not reach so high as the other two.——The *pisiforme* stands forwards in the palm from the *cuneiforme*.——The *trapezium* is the first of the second row, and is situated between the scaphoides and first joint of the thumb.——The *trapezoides* is immediately on the outside of the trapezium.——The *os magnum* is still more external.——The *unciforme* is farther to the side of the little finger.

Os scaphoides (*b*) is the largest of the eight, excepting one. It is convex above, concave and oblong below: From which small resemblance to a boat, it has got its name. Its smooth convex surface is divided by a rough middle fossa, which runs obliquely cross it. The upper largest division is articulated with the radius. The common ligament of the joint of the wrist is fixed into the fossa; and the lower division is joined to the trapezium and trapezoides. The concavity receives more than an half of the round head of the os magnum. The external side of this hollow is formed into a semilunar plane, to be articulated with the following bone.———The internal, posterior, and anterior edges are rough, for fixing the ligaments that connect it to the surrounding bones.

Os lunare (*c*) has a smooth convex upper surface, by which it is articulated with the radius. The internal side, which gives name to the bone, is in the form of a crescent, and is joined with the scaphoid;———the lower surface is hollow, for receiving part of the head of the os magnum. On the outside of this cavity is another smooth, but narrow, oblong sinuosity, for receiving the upper end of the os unciforme:———On the outside of which a small convexity is found, for its connection with the os cuneiforme. Between the great convexity above, and the first deep inferior cavity, there

(*b*) Κοτυλοειδες. Naviculare. (*c*) Lunatum.

there is a rough foſſa, in which the circular ligament of the joint of the wriſt is fixed.

Os cuneiforme (*d*) is broader above, and towards the back of the hand, than it is below and forwards; which gives it the reſemblance of a wedge. The ſuperior ſlightly convex ſurface is included in the joint of the wriſt, being oppoſed to the lower end of the ulna.—Below this the cuneiform bone has a rough foſſa, wherein the ligament of the articulation of the wriſt is fixed. On the internal ſide of this bone, where it is contiguous to the os lunare, it is ſmooth and ſlightly concave. Its lower ſurface, where it is contiguous to the os unciforme, is oblong, ſomewhat ſpiral, and concave. Near the middle of its anterior ſurface a circular plane appears, where the os piſiforme is ſuſtained.

Os piſiforme (*e*) is almoſt ſpherical, except one circular plane, or ſlightly hollow ſurface, which is covered with cartilage for its motion on the cuneiform bone, from which its whole rough body is prominent forwards into the palm; having the tendon of the flexor carpi ulnaris, and a ligament from the ſtyloid proceſs of the ulna, fixed to its upper part; the tranſverſe ligament of the wriſt is connected to its internal ſide; ligaments extended to the unciform bone, and to the os metacarpi of the little finger, are attached to its lower part; the abductor minimi digiti has its origin from its forepart; and, at the internal ſide of it, a ſmall depreſſion is formed, for the paſſage of the ulnar nerve.

Trapezium (*f*) has four unequal ſides and angles in its back-part, from which it has got its name.———Above, its ſurface is ſmooth, ſlightly hollowed, and ſemicircular, for its conjunction with the os ſcaphoides.——Its external ſide is

(*d*) Triquetrum.

(*e*) Cartilaginoſum, ſubrotundum, rectum.

(*f*) Os cubiforme, trapezoides, multangulum, majus.

is an oblong concave ſquare for receiving the following bone. The inferior ſurface is formed into a pulley; the two protuberant ſides of which are external and internal. On this pulley the firſt bone of the thumb is moved.—At the external ſide of the external protuberance, a ſmall oblong ſmooth ſurface is formed by the os metacarpi indicis. The fore-part of the trapezium is prominent in the palm, and near to the external ſide has a ſinuoſity in it, where the tendon of the flexor carpi radialis is lodged; on the ligamentous ſheath of which the tendon of the flexor tertii internodii pollicis plays: And ſtill more externally the bone is ſcabrous, where the tranſverſe ligament of the wriſt is connected, the abductor and flexor primi internodii pollicis have their origin, and ligaments go out to the firſt bone of the thumb.

Os trapezoides (*g*), ſo called from the irregular quadrangular figure of its back-part, is the ſmalleſt bone of the wriſt except the piſiforme. The figure of it is an irregular cube. It has a ſmall hollow ſurface above, by which it joins the ſcaphoides; a long convex one internally, where it is contiguous to the trepezium; a ſmall external one, for its conjunction with the os magnum; and an inferior convex ſurface, the edges of which are, however, ſo raiſed before and behind, that a ſort of pulley is formed, where it ſuſtains the os metacarpi indicis.

Os magnum (*h*), ſo called becauſe it is the largeſt bone of the carpus, is oblong, having four quadrangular ſides, with a round upper end, and a triangular plain one below. The round head is divided by a ſmall riſing, oppoſite to the connection of the os ſcaphoides and lunare, which together form the cavity for receiving it. On the inſide a ſhort plain ſurface joins the os magnum to the trapezoides. On the outſide is a long narrow concave ſurface, where it is

(*g*) Trapezium, multangulum minus.
(*h*) Maximum, capitatum.

is contiguous to the os unciforme. The lower end, which ſuſtains the metacarpal bone of the middle finger, is triangular, ſlightly hollowed, and farther advanced on the internal ſide than on the external, having a conſiderable oblong depreſſion made on the advanced inſide by the metacarpal bone of the fore-finger; and generally there is a ſmall mark of the os metacarpi digiti annularis on its external ſide.

Os unciforme (i) has got its name from a thin broad proceſs that ſtands out from it forwards into the palm, and is hollow on its inſide, for affording paſſage to the tendons of the flexors of the fingers. To this proceſs alſo the tranſverſe ligament is fixed that binds down and defends theſe tendons; and the flexor and abductor muſcles of the little finger have part of their origin from it. The upper plain ſurface is ſmall, convex, and joined with the os lunare: The internal ſide is long, and ſlightly convex, adapted to the contiguous os magnum. The external ſurface is oblique, and irregularly convex, to be articulated with the cuneiform bone. The lower end is divided into two concave ſurfaces; the external is joined with the metacarpal bone of the little finger; and the internal one is fitted to the metacarpal bone of the ring-finger.

In the deſcription of the preceding eight bones, I have only mentioned thoſe plain ſurfaces covered with cartilage, by which they are articulated to each other, or to ſome other bones, except in ſome few caſes, where ſomething extraordinary was to be obſerved; and I have deſignedly omitted the other rough ſurfaces, leſt, by crowding too many words in the deſcription of ſuch ſmall bones, the whole ſhould be unintelligible. But theſe ſcabrous parts of the bones may eaſily be underſtood after mentioning their figure, if it is obſerved, that they are generally found only towards the back or palm

(i) Cuneiforme.

of the hand; that they are all plain, larger behind than before; and that they receive the different ligaments, by which they are either connected to neighbouring bones or to one another; for these ligaments cover all the bones, and are so accurately applied to them, that at first view the whole carpus of a recent subject, appears one smooth bone (*k*).

As the surfaces of these bones are largest behind, the figure of the whole conjoined must be convex there, and concave before; which concavity is still more increased by the os pisiforme, and process of the os unciforme, standing forwards on one side, as the trapezium does on the other: And the bones are securely kept in this form by the broad strong transverse ligament connected to those parts of them that stand prominent into the palm of the hand. The convexity behind renders the whole fabric stronger, where it is most exposed to injuries; and the large anterior hollow is necessary for a safe passage to the numerous vessels, nerves, and tendons of the fingers.

The *substance* of these bones is spongy and cellular, but strong in respect of their bulk.

The three first bones of the carpus make an oblong head, by which they are *articulated* with the cavity at the lower ends of the bones of the fore-arm, so as to allow motion on all sides; and by a quick succession of these motions, they may be moved in a circle. But as the joint is oblong, and therefore the two dimensions are unequal, no motion is allowed to the carpus round its axis, except what it has in the pronation and supination along with the radius.—— The articulation of the first three bones of the superior row, with the bones of the inferior, is such as allows of motion, especially backwards and forwards; to the security and easiness of

(*k*) Galen. De usu part. lib. 2. cap. 8. For a particular description of these ligaments, see Weitbrecht. Syndesmolog. p. 5.——68.

of which, the reception of the os magnum into the cavity formed by the ſcaphoides and lunare conſiderably contributes: And the greateſt number of the muſcles that ſerve for the motion of the wriſt on the radius, being inſerted beyond the conjunction of the firſt row of bones with the ſecond, act equally on this articulation as they do on the former; but the joint formed with the radius being the moſt eaſily moved, the firſt effect of theſe muſcles is on it; and the ſecond row of the carpus is only moved afterwards. By this means a larger motion of the wriſt is allowed than otherwiſe it could have had ſafely: for if as large motion had been given to one joint, the angle of flexion would have been very acute, and the ligaments muſt have been longer than was conſiſtent with the firmneſs and ſecurity of the joint. The other articulations of the bones here being by nearly plain ſurfaces, ſcarcely allow of any more motion, becauſe of the ſtrong connecting ligaments, than to yield a little, and ſo elude the force of any external power; and to render the back of the wriſt a little more flat, or the palm more hollow, on proper occaſions. The articulations of the thumb and metacarpal bones ſhall be examined afterwards.

The *uſes* of the carpus are to ſerve as a baſe to the hand, to protect its tendons, and to afford it a free large motion.

All the bones of the carpus are in a cartilaginous ſtate at the time of birth.

On account of the many tendons that paſs upon the lower end of the fore-arm and the carpus, and of the numerous ligaments of theſe tendons and of the bones, which have lubricating liquors ſupplied to them, the pain of ſprains here is acute, the parts take a long time to recover their tone, and their ſwellings are very obſtinate.

ME-

METACARPUS.

METACARPUS (*l*) consists of four bones which sustain the fingers. Each bone is long and round, with its ends larger than its body.——The upper end, which some call the base, is flat and oblong, without any considerable head or cavity; but it is, however, somewhat hollowed, for the articulation with the carpus: It is made flat and smooth on the sides where these bones are contiguous to each other. Their bodies are flatted on their back-part by the tendons of the extensors of the fingers. The anterior surface of these bodies is a little concave, especially in their middle; along which a sharp ridge stands out, separating the musculi interossei placed on each side of these bones, which are there made flat and plain by these muscles.

Their lower ends are raised into large oblong smooth heads, whose greatest extent is forwards from the axis of the bone. At the fore-part of each side of the root of each of these heads, one or two tubercles stand out, for fixing the ligaments that go from one metacarpal bone to another, to preserve them from being drawn asunder: Round the heads a rough ring may be remarked, for the capsular ligaments of the first joints of the fingers to be fixed to; and both sides of these heads are flat, by pressing on each other.

The *substance* of the metacarpal bones is the same with that of all long bones.

At the time of birth, these bones are cartilaginous at both ends, which afterwards become epiphyses.

The metacarpal bones are *joined* above to the ossa carpi and to each other by nearly plain surfaces. These connections

(*l*) Κτεις, προπαχτιον, στηθος, ανδηρον, κτενιον, Postbrachiale, pectus, palma, pecten.

tions are not fit for large motions. The articulation of their round heads at the lower ends with the cavities of the firſt bones of the fingers, is to be taken notice of hereafter.

The concavity on the fore-part of theſe metacarpal bones, and the placing their baſes on the arched carpus, cauſe them to form a hollow in the palm of the hand, which is often uſeful to us. The ſpaces between them lodge muſcles, and their ſmall motion makes them fit ſupporters for the fingers to play on.

Though the oſſa metacarpi ſo far agree, yet they may be diſtinguiſhed from each other by the following marks.

The *os metacarpi indicis* is generally the longeſt.—Its baſe, which is articulated with the os trapezoides, is hollow in the middle. The ſmall ridge on the internal ſide of this oblong cavity is ſmaller than the one oppoſite to it, and is made flat on the ſide by the trapezium. The exterior ridge is alſo ſmooth, and flat on its outſide, for its conjunction with the os magnum; immediately below which a ſemicircular ſmooth flat ſurface ſhews the articulation of this to the ſecond metacarpal bone. The back part of this baſe is flatted where the long head of the extenſor carpi radialis is inſerted, and its fore-part is prominent where the tendon of the flexor carpi radialis is fixed. The external ſide of the body of this bone is more hollowed by the action of muſcles than the internal. The tubercle at the internal root of its head is larger than the external. Its baſe is ſo firmly fixed to the bone it is connected with, that it has no motion.

Os metacarpi medii digiti is generally the ſecond in length but often it is as long as the former; ſometimes it is longer; and it frequently appears only to equal the firſt by the os magnum being farther advanced downwards than any other bone of the wriſt. Its baſe is a broad ſuperficial cavity,

vity, ſlanting outwards: The internal poſterior angle of which is ſo prominent, as to have the appearance of a proceſs. The internal ſide of this baſe is made plain in the ſame way as the external ſide of the former bone, while its external ſide has two hollow circular ſurfaces, for joining the third metacarpal bone; and between theſe ſurfaces there is a rough foſſa, for the adheſion of a ligament, and lodging mucilaginous glands. The ſhorter head of the bicornis is inſerted into the back part of this baſe. The two ſides of this bone are almoſt equally flatted; only the ridge on the fore-part of the body inclines outwards. The tubercles at the fore-part of the root of the head are equal. The motion of this bone is very little more than that of the former; and therefore theſe two firmly reſiſt bodies preſſed againſt them by the thumb or fingers, or both.

Os metacarpi digiti annularis is ſhorter than the ſecond metacarpal bone. Its baſe is ſemicircular and convex, for its conjunction with the os unciforme. On its internal ſide are two ſmooth convexities, and a middle foſſa, adapted to the ſecond metacarpal bone. The external ſide has a triangular ſmooth concave ſurface to join it with the fourth one. The anterior ridge of its body is ſituated more to the out than to the inſide.—The tubercles near the head are equal.—The motion of this third metacarpal bone is greater than the motion of the ſecond.

Os metacarpi minimi digiti is the ſmalleſt and ſharpeſt. Its baſe is irregularly convex, and riſes ſlanting outwards.——Its internal ſide is exactly adapted to the third metacarpal bone. ——The external has no ſmooth ſurface, becauſe it is not contiguous to any other bone; but it is prominent where the extenſor carpi ulnaris is inſerted.——As this metacarpal bone is furniſhed with a proper moving muſcle, has the plaineſt articulation, is moſt looſely connected and leaſt confined, it

not

not only enjoys a much larger motion than any of the reſt, but draws the third bone with it, when the palm of the hand is to be made hollow by its advancement forwards, and by the prominence of the thumb oppoſite to it.

THUMB AND FINGERS.

THE thumb and four fingers are each compoſed of three long bones.

The THUMB (*m*) is ſituated obliquely in reſpect of the fingers, neither oppoſite directly to them, nor in the ſame plane with them.——All its bones are much thicker and ſtronger in proportion to their length, than the bone of the fingers are: Which are extremely neceſſary, ſince the thumb counteracts all the fingers.

The firſt bone of the thumb has its baſe adapted to the double pulley of the trapezium: For, in viewing it from one ſide to the other, it appears convex in the middle; but when conſidered from behind forwards, it is concave there. ——The edge at the fore-part of this baſe is produced farther than any other part; and round the back-part of the baſe a rough foſſa may be ſeen, for the connection of the ligaments of this joint. The body and head of this bone are of the ſame ſhape as the oſſa metacarpi; only that the body is ſhorter, the head flatter, and the tubercles at the fore-part of its root larger.

The articulation of the upper end of this bone is uncommon: For, though it has protuberances and depreſſions adapted to the double pulley of the trapezium; yet it enjoys a circular motion, as the joints do where a round head of the one

(*m*) Αντιχειρ, δικονδυλος, Magnus digitus, promanus.

one plays in the orbicular ſocket of another; only it is ſomewhat more confined, and leſs expeditious, but ſtronger and more ſecure than ſuch joints generally are.

This bone in children is in the ſame ſtate with the metacarpal bones.

The ſecond bone of the thumb has a large baſe formed into an oblong cavity, whoſe greateſt length is from one ſide to the other.———Round it ſeveral tubercles may be remarked, for the inſertion of ligaments.———Its body is convex, or a half round behind; but flat before, for lodging the tendon of the long flexor of the thumb, which is tied down by ligamentous ſheaths that are fixed on each ſide to the angle at the edge of this flat ſurface.———The lower end of this ſecond bone has two lateral round protuberances, and a middle cavity, whoſe greateſt extent of ſmooth ſurface is forwards.

The articulation and motion of the upper end of this ſecond bone is as ſingular as that of the former.———For its cavity being joined to the round head of the firſt bone, it would ſeem to enjoy motion in all directions; yet, becauſe of the ſtrength of its lateral ligaments, oblong figure of the joint itſelf, and mobility of the firſt joint, it only allows flexion and extenſion; and theſe are generally much confined.

The third bone of the thumb is the ſmalleſt, with a large baſe, whoſe greateſt extent is from one ſide to the other.———This baſe is formed into two cavities and a middle protuberance, to be adapted to the pulley of the former bone.—Its body is rounded behind; but is flatter than in the former bone for ſuſtaining the nail.—It is flat and rough before, by the inſertion of the flexor tertii internodii.—This bone becomes gradually ſmaller, till near the lower end, where it is a little enlarged, and has an oval ſcabrous edge.

The

The motion of this third bone is confined to flexion and extenſion.

THE orderly diſpoſition of the bones of the FINGERS into three rows, has made them generally obtain the name of three *phalanges* (*n*).—All of them have half-round convex ſurfaces, covered with an aponeuroſis, formed by the tendons of the extenſors, lumbricales, and interoſſei, and placed directly backwards, for their greater ſtrength; and their flat concave part is forwards, for taking hold more ſurely, and for lodging the tendons of the flexor muſcles.—The ligaments for keeping down theſe tendons are fixed to the angles that are between the convex and concave ſides.

The bones of the firſt phalanx (*o*) of the fingers anſwer to the deſcription of the ſecond bone of the thumb; only that the cavity in their baſe is not ſo oblong; nor is their motion on the metacarpal bones ſo much confined: For they can be moved laterally or circularly; but have no rotation, or a very ſmall degree of it, round their axis.

Both the ends of this firſt phalanx are in a cartilaginous ſtate at the birth; and the upper one is afterwards affixed in form of an epiphyſe.

The ſecond bone (*p*) of the fingers has its baſe formed into two lateral cavities, and a middle protuberance; while the lower end has two lateral protuberances and a middle cavity; therefore it is joined at both ends in the ſame manner, which none of the bones of the thumb are.

This bone is in the ſame condition with the former in children.

The third bone (*q*) differs nothing from the deſcription of the third bone of the thumb, excepting in the general

(*n*) Scytalidae, internodia, ſcuticula, agmina, acies, condyli articuli.

(*o*) Πρωτοσφόνδυλοι. (*p*) Κονδύλοι. (*q*) Μεταποσφόνδυλοι, ραχώνυχια.

diſtinguiſhing marks; and therefore the ſecond and third phalanx of the fingers enjoy only flexion and extenſion.

The upper end of this third phalanx is a cartilage in a ripe child; and is only an epiphyſe after, till the full growth of the body.

All the difference of the *phalanges* of the ſeveral fingers conſiſts in their magnitude ——The bones of the *middle finger* (*r*) being the longeſt and largeſt,——thoſe of the *fore-finger* (*s*) come next to that in thickneſs, but not in length, for thoſe of the *ring-finger* (*t*) are a little longer. The *little finger* (*u*) has the ſmalleſt bones. Which diſpoſition is the beſt contrivance for holding the largeſt bodies; becauſe the longeſt fingers are applied to the middle largeſt periphery of ſuch ſubſtances as are of a ſpherical figure (*x*).

The uſes of all the parts of our *ſuperior extremities* are ſo evident in the common actions of life, that it is needleſs to enumerate them here; and therefore I ſhall proceed to the laſt part of the ſkeleton. Only, leſt I ſhould ſeem to have forgot the ſmall bones at the joints of the hand, I deſire now to refer to the deſcription of them, under the common title of *ſeſamoid bones*, which I have placed after the bones of the feet.

SECT. IV.

OF THE INFERIOR EXTREMITIES.

THE INFERIOR EXTREMITIES depend from the acetabula of the oſſa innominata: and are commonly divided into three parts, viz. the thigh, leg, and foot.

§ 1.

(*r*) Κατατυγων, σφακελος, Infamis, impudicus, verpus, famoſus, obſcœnus.

(*s*) Δεικτικος, Indicator, λιχανος, demonſtrativus, ſalutaris.

(*t*) Ἰατρικος, παραμεσος, δακτυλιωτης, επιβατης, Annularis, medicus, cordi-digitus.

(*u*) Μυωψ, ωτικης, Auricularis, minimus.

(*x*) Galen. de uſu part. lib. I. cap. 24.

§ 1. *The* THIGH.

THE thigh (*y*) has only one bone; which is the longeſt in the whole body, and the largeſt and ſtrongeſt of any of the cylindrical bones. The ſituation of it is not perpendicular; for the lower end is inclined conſiderably inwards; ſo that the knees are almoſt contiguous, while there is a conſiderable diſtance between the thigh-bones above; which is of great uſe, ſince ſufficient ſpace is thereby left for the external parts of generation, the two great cloacae of urine and faeces, and for the large thick muſcles that move the thigh inwards. At the ſame time this ſituation of the thigh-bones renders our progreſſion quicker, ſurer, ſtraighter, and in leſs room: For, had the knees been at a greater diſtance from each other, we muſt have been obliged to deſcribe ſome part of a circle with the trunk of our body in making a long ſtep; and when one leg was raiſed from the ground, our centre of gravity would have been too far from the baſe of the other, and we ſhould conſequently have been in danger of falling; ſo that our ſteps would neither have been ſtraight nor firm, nor would it have been poſſible to walk in a narrow path, had our thigh-bones been otherwiſe placed. In conſequence, however, of the weight of the body bearing ſo obliquely on the joint of the knee by this ſituation of the thigh-bones, weak ricketty children become inn-knee'd.

The upper end of the thigh-bone is not continued in a ſtraight line with the body of it, but is ſet off obliquely inwards and upwards, whereby the diſtance here between theſe two bones at their upper part is conſiderably increaſed.——This end is formed into a large ſmooth round head (*z*), which is the greater portion of a ſphere unequally divided.——Towards its lower internal part a round rough

(*y*) *Μηρος*, Femen, coxa, agis, anchae os, crus, femur.

(*z*) Vertebrum.

rough ſpongy pit is obſervable, where the ſtrong ligament, commonly, but unjuſtly, called the *round one*, is fixed, to be extended from thence to the lower internal part of the receiving cavity, where it is conſiderably broader than near to the head of the thigh-bone.——The ſmall part below the head, called the *cervix*, of the os femoris, has a great many large holes, into which the fibres of the ſtrong ligament continued from the capſular, enter, and are thereby firmly united to it; and round the root of the neck, where it riſes from the bones, a rough ridge is found, where the capſular ligament of the articulation itſelf is connected.——Below the back-part of this root, the large unequal protuberante, called *trochanter major* (*a*), ſtands out; the external convex part of which is diſtinguiſhed into three different ſurfaces: Whereof the one on the fore-part is ſcabrous and rough, for the inſertion of the glutaeus minimus; the ſuperior one is ſmooth, and has the glutaeus medius inſerted into it; and the one behind is made flat and ſmooth, by the tendon of the glutaeus maximus paſſing over it.——The upper edge of this proceſs is ſharp and pointed at its back-part, where the glutaeus medius is fixed; but forwards it is more obtuſe, and has two ſuperficial pits formed in it: Into the ſuperior of theſe the pyriformis is implanted; and the obturator internus and gemini are fixed into the lower one.——From the hindmoſt prominent part of this great trochanter, a rough ridge runs backwards and downwards, into which the quadratus is inſerted.——In the deep hollow, at the internal upper ſide of this ridge, the obturator externus is implanted.——More internally, a conoid proceſs, called *trochanter minor* (*b*), riſes for the inſertion of the muſculus pſoas and iliacus internus; and the pectineus is implanted into a rough hollow below its internal

(*a*) Γλουτος, Rotator natis, malum granatum teſticulorum.

(*b*) Rotator minor.

nal root. The muſcles inſerted into thoſe two proceſſes being the principal inſtruments of the rotatory motion of the thigh, have occaſioned the name of *trochanters* to be given to thoſe proceſſes.——The tendons that are fixed into or paſs over the great *trochanter*, cauſe bruiſes, by falls on this part, to be attended with great pain and weakneſs of the limb, which generally remain long.

The body of the os femoris is convex on the fore-part, and made hollow behind, by the action of the muſcles that move it and the leg, and for the conveniency of ſitting, without bearing too much on theſe muſcles; and probably the weight of the legs depending from the thighs in that poſture contributes to this curvature. The fore-part of the thigh-bone is a little flatted above by the beginning of the cruraeus muſcle, as it is alſo below by the ſame muſcle and the rectus. ——Its external ſurface is likewiſe made flat below by the vaſtus externus, where it is ſeparated from the former by an obtuſe ridge. The vaſtus internus depreſſes a little the lower part of the internal ſurface.——The poſterior concave ſurface has a ridge riſing in its middle, commonly called *linea aſpera*, into which the triceps is inſerted, and the ſhort head of the biceps flexor tibiae riſes from it.——At the upper part of it the medullary veſſels enter by a ſmall hole that runs obliquely upwards. A little above which there is a rough foſſa or two, where the tendon of the glutaeus maximus is fixed. ——The lower end of the linea aſpera divides into two, which deſcend towards each ſide.——The two vaſti muſcles have part of their origin from theſe ridges; and the long tendon of the triceps is fixed to the internal, by a part of the faſcia aponeurotica of the thigh.——Near the beginning of the internal ridge, there is a diſcontinuation of the ridge, where the crural artery paſſes through the aponeuroſis.—— Between theſe two rough lines, the bone is made flat by the large blood-veſſels and nerves which paſs upon it; and near

the

the end of each of theſe ridges, a ſmall ſmooth protuberance may often be remarked, where the two heads of the external gaſtrocnemius muſcle take their riſe, and where ſeſamoid bones are ſometimes found (*c*); and from the fore-part of the internal tubercle a ſtrong ligament is extended to the inſide of the tibia.

The lower end of the os femoris is larger than any other part of it, and is formed into a great protuberance on each ſides, called its *condyles;* between which a conſiderable cavity is found, eſpecially at the back-part, in which the crural veſſels and nerves lie immerſed in fat.——The internal condyle is longer than the external, which muſt happen from the oblique poſition of this bone, to give leſs obliquity to the leg. Each of theſe proceſſes ſeem to be divided into its plain ſmooth ſurface. The mark of diviſion on the external is a notch, and on the internal a ſmall protuberance. The fore-part of this diviſion, on which the rotula moves, is formed like a pulley, the external ſide of which is higheſt. Behind, there are two oblong large heads, whoſe greateſt extent is backwards, for the motion of the tibia: and from the rough cavity between them, but near to the baſe of the internal condyle, the ſtrong ligament, commonly called the *croſs one*, has its riſe.——A little above which a protuberance gives inſertion to the tendon of the triceps. The condyles, both on the outer and inner ſide of the knee, are made flat by the muſcles paſſing along them. On the back-part of the internal, a ſlight depreſſion is made by the tendons of the gracilis and ſartorius; and on the external ſuch another is formed by the biceps flexor cruris; behind which a deep foſſa is to be obſerved, where the poplitaeus muſcle has its origin. From the tubercle immediately before this cavity, a ſtrong ligament goes out to the upper part of the fibula.——Round this lower end

(*c*) Veſal. lib. 1. cap. 28. & 30.

end of the thigh-bone, large holes are found, into which the ligaments for the ſecurity of the joint are fixed, and blood-veſſels paſs to the internal ſubſtance of the bone.

All the proceſſes of the femur are cartilaginous in new-born children; and afterwards become ſmall apophyſes, with large epiphyſes.

The thigh-bone being articulated above with the acetabulum of the oſſa innominata, which affords its round head a ſecure and extenſive play, can be moved to every ſide; but is reſtrained in its motion outwards by the high brims of the cavity, and by the round ligament; for otherwiſe the head of the bone would have been frequently thruſt out at the breach of the brims on the inſide, which allows the thigh to move conſiderably inwards.——The body of this bone enjoys little or no rotatory motion, though the head moſt commonly moves round its own axis; becauſe the oblique progreſs of the neck and head from the bone is ſuch, that the rotatory motion of the head can only bring the body of the bone forwards and backwards. Nor is the head, as in the arm, ever capable of being brought to a ſtraight direction with its body; ſo far, however, as the head can move within the cavity backwards and forwards, the reſt of the bone may have a partial rotation.——When the thigh-bone reſiſts the actions of its muſcles more than the trunk of the body can then do, as in ſtanding, theſe muſcles have their effect on the trunk, cauſing it to bend forward, raiſing it up, inclining it to the one or the other ſide, twiſting it obliquely, &c. for which the rolling of the acetabula of the oſſa innominata on the round heads of the thigh-bones is well fitted.——The os femoris is articulated below to the tibia and rotula in the manner afterwards to be deſcribed.

The nearneſs of the ſmall neck to the round head of the

the thigh-bone, and its upper end being covered with very thick muſcles, make greater difficulty in diſtinguiſhing between a luxation and fracture here, than in any other part of the body.

§ 2. *The* LEG.

THE leg (*d*) is compoſed, according to the common account, of two bones, *tibia* and *fibula*, though it ſeems to have a very good title to a third, the *rotula*; which bears a ſtrong analogy to the olecranon of the ulna, and moves always with the other two.

TIBIA.

TIBIA (*e*), ſo called from its reſemblance to an old muſical pipe or flute, is the long thick triangular bone, ſituated at the internal part of the leg, and continued in almoſt a ſtraight line from the thigh-bone.

The upper end of the tibia is large, bulbous, and ſpongy, and is divided into two cavities by a rough irregular protuberance (*f*), which is hollow at its moſt prominent part, as well as before and behind. The anterior of the two ligaments that compoſe the great croſs one is inſerted into the middle cavity, and the depreſſion behind receives the poſterior ligament.——The two broad cavities at the ſides of this protuberance are not equal: for the internal is oblong and deep, to receive the internal condyle of the thigh-bone; while the external is more ſuperficial and rounder, for the external condyle.——In each of theſe two cavities of a recent ſubject, a ſemilunar cartilage is placed,

(*d*) Κνημη Crus, tibia.

(*e*) Προκνημιον, αντικνημιον, Focile majus, arundo major, canna major, canna domeſtica cruris.

(*f*) Διαφυσις, εξοχη κυρτοκοιλωδης, Tuber, tuberculum.

placed, which is thick at its convex edge, and becomes gradually thinner towards the concave or interior edge. The middle of each of these cartilages is broad, and the ends of them turn narrower and thinner as they approach the middle protuberance of the tibia. The thick convex edge of each cartilage is connected to the capsular and other ligaments of the articulation: but so near to their rise from the tibia, that the cartilages are not allowed to change place far; while the narrow ends of the cartilages becoming almost ligaments, are fixed at the insertion of the strong cross ligament into the tibia; and seem to have their substance united with it; therefore a circular hole is left between each cartilage and the ligament, in which the most prominent convex part of each condyle of the thigh-bone moves. The circumference of these cavities is rough and unequal, for the firm connection of the ligaments of the joint. Immediately below the edge at its back-part, two rough flatted protuberances stand out; into the internal, the tendon of the semimembranosus muscle is inserted; and a part of the cross ligament is fixed to the external.——On the outside of this last tubercle, a smooth slightly-hollowed surface is formed by the action of the poplitaeus muscle.

Below the fore-part of the upper end of the tibia, a considerable rough protuberance (*g*) rises, to which the strong tendinous ligament of the rotula is fixed.——On the internal side of this, there is a broad scabrous slightly-hollowed surface, to which the internal long ligament of the joint, the aponeurosis of the vastus internus, and the tendons of the seminervosus, gracilis, and sartorius, are fixed. The lowest part of this surface is therefore the place where the tibia ought to be sawn through in an amputation, so as not to have too long and troublesome a stump,

(*g*) Αντικνημιον, Anterior tuber.

and at the ſame time to preſerve its motions, by ſaving the proper muſcles. Below the external edge of the upper end of the tibia, there is a flat circular ſurface, covered in a recent ſubject with cartilage, for the articulation of the fibula; between which and the anterior knob, a rough hollow affords origins to the tibialis anticus, and extenſor digitorum longus. From this ſmooth flat ſurface, a ridge runs obliquely downwards and inwards, to give riſe to part of the ſolaeus, tibialis poſticus, and flexor digitorum longus, and inſertion to the aponeuroſis of the ſemimembranoſus which covers the poplitaeus, and to ſome of the external fibres of this laſt-named muſcle. At the inſide of this ridge an oblique plain ſurface is left, where the greateſt part of the muſculus poplitaeus is inſerted. The remaining body of the tibia is triangular. The anterior angle is very ſharp, and is commonly called the *ſpine* or *ſhin (b)*. This ridge is not ſtraight; but turns firſt inwards, then outwards, and laſtly inwards again. The plain internal ſide is ſmooth and equal, being little ſubjected to the actions of muſcles; but the external ſide is hollowed above by the tibialis anticus, and below by the extenſor digitorum longus and extenſor pollicis longus. The two angles behind theſe ſides are rounded by the action of the muſcles; the poſterior ſide comprehended between them is not ſo broad as thoſe already mentioned, but is more oblique and flatted by the action of the tibialis poſticus and flexor digitorum longus.——A little above the middle of the bone, the internal angle terminates, and the bone is made round by the preſſure of the muſculus ſolaeus. Near to this the paſſage of the medullary veſſels is ſeen ſlanting obliquely downwards.

The lower end of the tibia is hollowed, ſo as to occaſion a ſmall protuberance to riſe in the middle. The internal ſide

(b) Ακανθα, Spina, cerca, linea prima tibiae, angulus acutus.

ſide of this cavity, which is ſmooth, and in a recent ſubject is covered with cartilage, is produced into a conſiderable proceſs, commonly named *malleolus internus (i)*; the point of which is divided by a notch, and from it ligaments are ſent out to the foot. We ought to obſerve here, that this internal malleolus is ſituated more forwards than the internal condyle of the upper end of this bone; which is neceſſary to be remembered in reducing a fracture of the leg (*k*). The external ſide of this end of the tibia has a rough irregular ſemilunar cavity formed in it for receiving the lower end of the fibula. The poſterior ſide has two lateral grooves, and a ſmall middle protuberance. In the internal depreſſion, the tendons of the muſculus tibialis poſticus and flexor digitorum longus are lodged; and in the external, the tendon of the flexor longus pollicis plays. From the middle protuberance, ligamentous ſheaths go out, for tying down theſe tendons.

The articulations and motions of the tibia ſhall be explained, after all the three bones of the leg are deſcribed.

Both the ends of the tibia are cartilages at birth, and become afterwards epiphyſes.

FIBULA.

Fibula (*l*), is the ſmall long bone, placed on the outſide of the leg, oppoſite to the external angle of the tibia; the ſhape of it is irregularly triangular.

The head of the fibula has a ſuperficial circular cavity formed on its inſide, which, in a recent ſubject, is covered with a cartilage, and it is ſo cloſely connected to the tibia by ligaments, as to allow only a ſmall motion backwards and

(*i*) Σφυρον, πεσσος, Talus, clavicula, clavilla interior, clavilla domeſtica.

(*k*) Winſlow. Expoſition anatomique des os ſecs, § 865.

(*l*) Παραχνημιον, Perone, ſocile minus, arundo minor, canna minor cruris, ſura, radius.

and forwards. This head is protuberant and rough on its outside, where a ſtrong round ligament and the muſculus biceps are inſerted; and, below the back-part of its internal ſide, a tubercle may be remarked, that gives riſe to the ſtrong tendinous part of the ſolaeus muſcle.

The body of this bone is a little crooked inwards and backwards: which figure is owing to the actions of the muſcles; but is ſtill farther increaſed by nurſes, who often hold children careleſsly by the legs. The ſharpeſt angle of the fibula is forwards; on each ſide of which the bone is conſiderably, but unequally, depreſſed by the bellies of the ſeveral muſcles that riſe from or act upon it; and in old people, theſe muſcles make diſtinct ſinuoſities for themſelves. The external ſurface of the fibula is depreſſed obliquely from above downwards and backwards by the two peronaei. Its internal ſurface is unequally divided into two narrow longitudinal planes, by an oblique ridge extended from the upper part of the anterior angle, to join with the lower end of the internal angle. To this ridge the ligament ſtretched between the two bones of the leg is connected. The anterior of the two two planes is very narrow above, where the extenſor longus digitorum and extenſor longus pollicis ariſe from it; but is broader below, where it has the print of the nonus Veſalii. The poſterior plane is broad and hollow, giving origin to the larger ſhare of the tibialis poſticus. The internal angle of this bone has a tendinous membrane fixed to it, from which ſome fibres of the flexor digitorum longus take their riſe. The poſterior ſurface of the fibula is the plaineſt and ſmootheſt: but is made flat above by the ſolaeus, and is hollowed below by the flexor pollicis longus. In the middle of this ſurface, the canal for the medullary veſſels may be ſeen ſlanting downwards.

I have taken particular notice of the entry and direction of

of the medullary veſſels of the large bones of the extremities (*m*); becauſe in ſeveral chirurgical caſes, a ſurgeon who is ignorant of their ſituation and courſe, may do miſchief to his patient. Thus, for example, if theſe veſſels are opened very near their entry into the bone, or while they are in the oblique paſſage through it, an obſtinate haemorrhagy may enſue: For the arteries being connected to the bony paſſage, ſtyptics, and other like corrugators, are applied in vain; compreſſing inſtruments can do no ſervice, and ligatures cannot be employed. There ſeems to be a particular deſign in the contrivance of theſe canals; thoſe in the os humeri, tibia, and fibula running obliquely downwards from their external entry; whereas in the radius, ulna, and os femoris, they ſlant upwards, whereby the arteries and nerves which are ſent into theſe three laſt bones muſt ſuffer a conſiderable reflexion before they come at the cancelli. The reaſon of this diverſity may perhaps be, that the arteries, which are ſo ſmall within the bones as to have no ſtrong contractile propelling force in their coats, and where they are not aſſiſted by the action of any moving neighbouring organ, ſhould have, at leaſt in their paſſage through the bone, a favourable deſcent for their liquids: Which, it is evident, they have in the deſcending oblique paſſages formed for them in the firſt claſs of bones, to wit, the os humeri, tibia, and fibula, which are generally depending; and they alſo moſt frequently acquire the like advantage in the radius, ulna, and os femoris; becauſe the hand, in the moſt natural poſture, is higher than the elbow: And when we ſit or lie, the lower end of the thigh-bone comes to be at leaſt raiſed as high as the upper end. In ſtanding and walking, or when the arms are moved, the blood muſt indeed aſcend as it paſſes through the bones of the fore-arm and thigh; but the preſſure

(*m*) Havers, Oſteolog. Nov. diſ. 1. p. 59.

preſſure of the muſcles, then in action on the veſſels, before they enter the bones, is ſufficient to compenſate the diſadvantage of their courſe. This reaſoning ſeems to be ſtill enforced, by obſerving, that this paſſage is always nearer the upper than the lower ends of theſe bones.

The lower end of the fibula is extended into a ſpongy oblong head; on the inſide of which is a convex, irregular, and frequently a ſcabrous, ſurface, that is received by the external hollow of the tibia, and ſo firmly joined to it by a very thin intermediate cartilage and ſtrong ligaments, that it ſcarce can move.——Below this the fibula is ſtretched out into a ſmooth coronoid proceſs, covered with cartilage on its internal ſide, and is there contiguous to the outſide of the firſt bone of the foot, the aſtragalus, to ſecure the articulation. This proceſs, named *malleolus externus*, being ſituated farther back than the internal malleolus, and in an oblique direction, obliges us naturally to turn the fore-part of the foot outwards (*n*). At the lower internal part of this proceſs, a ſpongy cavity for mucilaginous glands may be remarked; from its point ligaments are extended to the aſtragalus, os calcis, and os naviculare, bones of the foot; and from its inſide ſhort ſtrong ones go out to the aſtragalus. On the back part of it a ſinuoſity is made by the tendons of the peronaei muſcles. When the ligament extended over theſe tendons from the one ſide of the depreſſion to the other is broken, ſtretched too much, or made weak by a ſprain, the tendons frequently ſtart forwards to the outſide of the fibula.

The conjunction of the upper end of the fibula with the tibia, is by plain ſurfaces tipped with cartilage; and at its lower end the cartilage ſeems to glue the two bones together; not, however, ſo firmly in young people, but that the motion at the other end of ſuch a long radius is very obſervable.

(*n*) Winſlow, Mem. de l'acad. des ſciences, 1722.

obſervable.—In old ſubjects I have often ſeen the two bones of the leg grown together at their lower ends.

The principal uſe of this bone is to afford origin and inſertion to muſcles; the direction of which may be a little altered on proper occaſions, by its upper part ſhuffling backwards and forwards.——It likewiſe helps to make the articulation of the foot more ſecure and firm.——The ends of the tibia and fibula being larger than their middle, a ſpace is here left, which is filled up with ſuch another ligament as I have deſcribed extended between the bones of the fore-arm; and which is alſo diſcontinued at its upper-part, where the tibialis anticus immediately adheres to the ſoleaus and tibialis poſticus; but every where elſe it gives origin to muſcular fibres (*o*).

Both the ends of this bone are cartilaginous in a ripe child, and aſſume the form of appendices before they are united to its body.

ROTULA.

Rotula (*p*) is the ſmall flat bone ſituated at the fore-part of the joint of the knee.—Its ſhape reſembles the common figure of the heart with its point downwards.—The anterior convex ſurface of the rotula is pierced by a great number of holes, into which fibres of the ſtrong ligament, that is ſpread over it, enter.—Behind, its ſurface is ſmooth, covered with cartilage, and divided by a middle convex ridge into two cavities, of which the external is largeſt; and both are exactly adapted to the pulley of the os femoris, on which they are placed in the moſt ordinary unſtraining poſtures of the leg: But when the leg is much bent, the rotula deſcends far down

(*o*) Weitbrecht, Syndeſmolog. p. 156.

(*p*) Επιμυλις, μυλακρις, κογχος, επιγονατις, πλακουντοειδες, Patella, mola, genu, ſcutiformę os, cartilaginoſum, diſciforme, oculus genu.

down on the condyles; and when the leg is fully extended, the rotula rises higher in its upper-part than the pulley of the thigh-bone.——The plain smooth surface is surrounded by a rough prominent edge, to which the capsular ligament adheres: Below, the point of the bone is scabrous, where the strong tendinous ligament from the tubercle of the tibia is fixed. The upper horizontal part of this bone is flatted and unequal, where the tendons of the extensors of the leg are inserted.

The substance of the rotula is cellular, with very thin external firm plates; but then these cells are so small, and such a quantity of bone is employed in their formation, that scarce any bone of its bulk is so strong. Besides, it is covered all over with a thick ligament, (as it was observed that this sort of bones generally is), to connect its substance, and is moveable to one side or other: Therefore it is sufficiently strong to resist the ordinary actions of the large muscles that are inserted into it, or any common external force applied to it; while a fixed process, such as the olecranon, would not have been sufficient to bear the whole weight of our bodies, which frequently falls on it, and would have hindered the rotatory motion of the leg. Notwithstanding these precautions to preserve this bone from such injuries, yet I have seen a transverse fracture in it, when, by the report of the patient, and of the people about him, and by the want of swelling, discolouring, or other mark of bruise or contusion, it was plain the bone was broken by the violent straining effort of the muscles(q). Though my patient recovered the use of the joint of the knee, yet I think it reasonable to believe, that this sort of fracture is commonly attended with difficulty of motion after the broken parts of the rotula are reunited; because the callous matter probably extends itself into the cavity of the joint, where it either grows to some of the parts, or makes such

(q) See Ruysch. Observ. anat. chirurg. obs. 3.

ſuch an inequality on the ſurface of this bone, as does not allow it to perform the neceſſary motions on the condyles of the femur (r).

At the ordinary time of birth, the rotula is entirely cartilaginous, and ſcarcely aſſumes a bony nature ſo ſoon as moſt epiphyſes do.

The parts which conſtitute the joint of the knee being now deſcribed, let us examine what are its motions, and how they are performed.—The two principal motions are flexion and extenſion. In the former of theſe, the leg may be brought to a very acute angle with the thigh, by the condyles of the thigh-bones being round and made ſmooth far backwards. In performing this, the rotula is pulled down by the tibia. When the leg is to be extended, the rotula is drawn upwards, conſequently the tibia forwards, by the extenſor muſcles; which, by means of the protuberant joint, and of this thick bone with its ligament, have the chord, with which they act, fixed to the tibia at a conſiderable angle, act, on that account, with advantage; but they are reſtrained from pulling the leg farther than to a ſtraight line with the thigh, by the poſterior part of the croſs ligament, that the body might be ſupported by a firm perpendicular column: For at this time the thigh and leg are as little moveable in a rotatory way, or to either ſide, as if they were one continued bone.—But when the joint is a little bent, the rotula is not tightly braced, and the poſterior ligament is relaxed; therefore this bone may be moved a little to either ſide, or with a ſmall rotation in the ſuperficial cavities of the tibia; which is done by the motion of the external cavity backwards and forwards, the internal ſerving as a ſort of axis (s). Seeing, then, one part of the croſs ligament is ſituated perpendicularly, and the poſ-

(r) Pare, liv. 15. cap. 22.

(s) Winſlow, Expoſition anatomique du corps humain, traite des os ſecs, § 976.

terior part is ſtretched obliquely from the internal condyle of the thigh outwards, that poſterior part of the croſs ligament prevents the leg from being turned at all inwards; but it could not hinder it from turning outwards almoſt round, was not that motion confined by the lateral ligaments of this joint, which can yield little. This rotation of the leg outwards is of good advantage to us in croſſing our legs, and turning our feet outwards, on ſeveral neceſſary occaſions; though it is altogether fit this motion ſhould not be very large, to prevent frequent luxations here.—While all theſe motions are performing, the part of the tibia that moves immediately on the condyles is only ſo much of it as is within the cartilaginous ring, which by the thickneſs on their outſides make the cavities of the tibia more horizontal; by raiſing their external ſide where the ſurface of the tibia ſlants downwards. By this means the motions of this joint are more equal and ſteady than otherwiſe they would have been. The cartilages being capable of changing a little their ſituation, are fit for doing this good office in the different motions and poſtures of the limb, and likewiſe contribute to make the motions larger and quicker.

On account of the very large ſurface of the bones forming the joint of the knee, and the many ſtrong ligaments connecting them, luxations ſeldom happen here. But theſe very ligaments, the aponeuroſis paſſing over this joint, and the quantity of fat and mucilaginous glands neceſſary for lubricating it, make it more ſubject to white ſwellings, dropſies, and ſuch other diſorders, than any other joint of the body.

§ 3. *The* FOOT.

THE foot is divided, as well as the hand, into three part, viz. *tarſus*, *metatarſus*, and *toes*: In the deſcription of

of which, the ſeveral ſurfaces ſhall be named, according to their natural ſituation, viz. the broad upper part of the foot ſhall be called *ſuperior*; the ſole, *inferior*; the ſide on which the great toe is, *internal*; and the other *external*.

TARSUS.

The tarſus (*t*) conſiſts of ſeven ſpongy bones; to wit, the aſtragalus, os calcis, naviculare, cuboides, cuneiforme externum, cuneiforme medium, and cuneiforme internum.

The *aſtragalus* is the uppermoſt of theſe bones.——The *os calcis* is below the aſtralagus, and is conſiderably prominent backwards beyond the other bones, to form the heel.——The *os naviculare* is in the middle of the internal ſide of the tarſus.——The *os cuboides* is the moſt external of the row of four bones at its fore-part.——The *os cuneiforme externum* is placed at the inſide of the cuboid.——The *cuneiforme medium* is between the external and internal cuneiform bones; and the *internal cuneiforme* is put at the internal ſide of the foot.

That the deſcription of theſe bones may not be immoderately ſwelled with repetition, I deſire, once for all, to obſerve, That wherever a ridge is mentioned, without a particular uſe aſſigned, a ligament is underſtood to be fixed to it; or where a ſpongy rough cavity, depreſſion, or foſſa, is remarked, without naming its uſe, a ligament is inſerted, and mucilaginous glands are lodged: for ſuch will occur in the detail of each of theſe bones.

Aſtragalus. The upper part of the aſtragalus (*u*) is formed into a large ſmooth head (*x*), which is ſlightly hollowed in the middle; and therefore reſembles a ſuperficial pulley, by which it is fitted to the lower end of the tibia.——The internal ſide of this head is flat and ſmooth, to play

(*t*) Raſſetta.

(*u*) Ἀστρις, Talus, balliſtae os, malleolus, chaib, quatrio, os teſſarae, claviculae, unciforme.

(*x*) Τροχιτος.

play on the internal malleolus.——The external ſide has alſo ſuch a ſurface, but larger, for its articulation with the external malleolus. Round the baſe of this head there is a rough foſſa; and immediately before the head, as alſo below its internal ſmooth ſurface, we find a conſiderable rough cavity.

The lower ſurface of the aſtralagus is divided by an irregular deep rough foſſa; which at its internal end is narrow, but gradually widens as it ſtretches obliquely outwards and forwards. The ſmooth ſurface, covered with cartilage, behind this foſſa, is large, oblong, extended in the ſame oblique ſituation with the foſſa, and concave for its conjunction with the os calcis. The back-part of the edge of this cavity is produced into two ſharp-pointed rough proceſſes; between which is a depreſſion made by the tendon of the flexor pollicis longus.——The lower ſurface before the foſſa is convex, and compoſed of three diſtinct ſmooth planes.——The long one behind, and the exterior or ſhorteſt, are articulated with the heel-bone; while the internal, which is the moſt convex of the three, reſts and moves upon a cartilaginous ligament, that is continued from the calcaneum to the os ſcaphoides. Without which ligament the aſtralagus could not be ſuſtained; but would be preſſed out of its place by the great weight it ſupports, and the other bones of the tarſus would be ſeparated. Nor would a bone be fit here, becauſe it muſt have been thicker than could conveniently be allowed; otherwiſe it would break, and would not prove ſuch an eaſy bending baſe, to leſſon the ſhock which is given to the body in leaping, running, &c.

The fore-part of this bone is formed into a convex oblong ſmooth head, called by ſome its *proceſs*, which is received by the os naviculare. Round the root of this head, eſpecially

eſpecially on the upper ſurface, a rough foſſa may be remarked.

The aſtragalus is articulated above to the tibia and fibula, which together form one cavity. Though in this articulation the bones have prominences and cavities, ſo ſmall as might allow motions in all directions, yet the flexion and extenſion are the moſt conſiderable, the other motions being confined by the malleoli, and, by the ſtrong ligaments which go out from the points of theſe proceſſes, to the aſtragalus and os calcis.——When the foot is bent, ſo far as it commonly is when we ſtand, no lateral or rotatory motion is allowed in this joint; for then the head of the aſtragalus is ſunk deep between the malleoli, and the ligaments are tenſe: but when the foot is extended, the aſtragalus can move a little to either ſide, and with a ſmall rotation. By this contrivance the foot is firm, when the weight of the body is to be ſupported on it; and when a foot is raiſed, we are at liberty to direct it more exactly to the place we intend next to ſtep upon.—The aſtragalus is joined below to the os calcis; and before to the os naviculare, in the manner to be explained when theſe bones are deſcribed.

A conſiderable ſhare of this bone is oſſified in a new-born infant.

Calcaneum (*w*) is the largeſt bone of the ſeven.———Behind, it is formed into a large knob, commonly called the *heel:* the ſurface of which is rough behind, where the tendo Achillis is inſerted; and above that part it is hollow and ſpongy. Farther forwards, on the upper ſurface of the calcaneum, there is an irregular oblong ſmooth convexity, adapted to the concavity at the back part of the aſtragalus: and beyond this a narrow foſſa is ſeen, which divides it from two ſmall concave ſmooth ſurfaces, that are joined to

(*w*) Os calcis, πτερνα, calcar pedis.

to the fore-part of the aſtragalus.——Behind the poſterior of theſe ſmooth ſurfaces, which is the largeſt, a ſmall ſinuoſity is made by the tendon of the flexor digitorum longus; at the fore-part of which a ſmall rough protuberance appears, that gives riſe to the muſculus extenſor digitorum brevis.

The external ſide of this bone is flat, with a ſuperficial foſſa running horizontally, in which the tendon of the muſculus peronaeus longus is lodged. The internal ſide of the heel-bone is hollowed, for lodging the origin of the maſſa cornea Jac. Sylvii, and for the ſafe paſſage of tendons, nerves, and arteries. Under the ſide of the internal ſmooth concavity, a particular groove is made by the tendon of the flexor pollicis longus; and from the thin protuberance of this internal ſide the cartilaginous ligament that ſupports the aſtragalus, goes out to the os naviculare; on which ligament, and on the edge of this bone to which it is fixed, the groove is formed for the tendon of the flexor digitorum profundus.

The lower ſurface of this bone is preſſed flat at the back-part, by the weight of our bodies; and immediately before this plane, there are two tubercles, from the internal of which the muſculus abductor pollicis, flexor digitorum ſublimis, as alſo part of the aponeuroſis plantaris, and of the abductor minimi digiti, have their origin; and the other part of the abductor minini digiti and aponeuroſis plantaris riſes from the external. Before theſe protuberances this bone is concave, for lodging the flexor muſcles; and at its fore-part we may obſerve a rough depreſſion, from which, and a tubercle behind it, the ligament goes out that prevents this bone from being ſeparated from the os cuboides.

The fore-part of the os calcis is formed into an oblong pulley-like ſmooth ſurface, which is circular at its upper external

external end, but is pointed below. The ſmooth ſurface is fitted to the os cuboides.

Though the ſurfaces by which the aſtragalus and os calcis are articulated, ſeem fit enough for motion; yet the very ſtrong ligaments by which theſe bones are connected, prevent much motion, and render this principal part of our baſe, which reſts on the ground, firm.

A large ſhare of the heel-bone is oſſified at the ordinary time of birth, and the large knob appears afterwards in form of an epiphyſe.

Os naviculare (*y*), is ſomewhat circular.——It is formed into an oblong concavity behind, for receiving the anterior head of the aſtragalus.——On the upper ſurface there is a rough foſſa.——Below, the os naviculare is very unequal and rough; but hollow for the ſafety of the muſcles.——On its inſide a large knob riſes out, from which the abductor pollicis takes in part its origin, the tendon of the tibialis poſticus is inſerted into it, and to it two remarkable ligaments are fixed; the firſt is the ſtrong one, formerly mentioned, which ſupports the aſtragalus; the ſecond is ſtretched from this bone obliquely croſs the foot, to the metatarſal bones of the middle toe, and of the toe next to the little one.——On the outſide of the os naviculare there is a ſemicircular ſmooth ſurface, where it is joined to the os cuboides. The fore-part of this bone is all covered with cartilage, and is divided into three ſmooth planes, fitted to the three oſſa cuneiformia.

The os naviculare and aſtragalus are joined as a ball and ſocket; and the naviculare moves in all directions in turning the toes inwards, or in raiſing or depreſſing either ſide of the foot, though the motions are greatly reſtrained by the ligaments which connect this to the other bones of the tarſus.

(*y*) Σκαφοειδης, Os cymbae.

tarſus. A weakneſs of theſe ligaments cauſes ſometimes an unnatural turn of the fore-part of the foot inwards.

The os naviculare is wholly cartilaginous in a new-born infant.

Os cuboides (z) is a very irregular cube.——Behind, it is formed into an oblong unequal concavity, adapted to the fore-part of the os calcis. On its internal ſide, there is a ſmall ſemicircular ſmooth cavity, to join the os naviculare.—Immediately before which, an oblong ſmooth plane is made by the os cuneiforme externum. Below this the bone is hollow and rough.—On the internal ſide of the lower ſurface a round protuberance and foſſa are found, where the muſculus adductor pollicis has its origin. On the external ſide of this ſame ſurface, there is a round knob, covered with cartilage; immediately before which a ſmooth foſſa may be obſerved, in which the tendon of the peronaeus primus runs obliquely croſs the foot; and on the knob the thin flat cartilage proper to this muſcle plays; in place of which ſometimes a bone is found: More externally than the knob, a rough hollow is made, for the ſtrong ligaments ſtretched between this bone and the os calcis.—Before, the ſurface of the os cuboides is flat, ſmooth, and ſlightly divided into two planes, for ſuſtaining the os metatarſi of the little toe, and of the toe next to it.

The form of the back-part of the os cuboides, and the ligaments connecting the joint with os calcis, both concur in allowing little motion in this part.

The oſſification of this bone is ſcarcely begun at the birth.

Os cuneiforme externum (*a*), if we regard its ſituation or medium by its bulk, is much of the ſhape of a wedge, being

(z) Πολυμορφον, Cubiforme, quadratum, grandinoſum varium teſſarae, multiforme.

(*a*) Chalcoideum externum.

ing broad and flat above, with long ſides running obliquely downwards, and terminating in a ſharp edge. The upper ſurface of this bone is an oblong ſquare. The one behind is nearly a triangle, but not complete at the inferior angle, and is joined to the os naviculare. The external ſide is an oblong ſquare divided as it were by a diagonal; the upper half of it is ſmooth, for its conjunction with the os cuboides: The other is a ſcabrous hollow, and in its ſuperior anterior angle a ſmall ſmooth impreſſion is made by the os metatarſi of the toe next to the little one. The internal ſide of this bone is alſo quadrangular, with the fore-part of its edge made flat and ſmooth by the os metatarſi of the toe next to the great one, and the back-part is alſo flat and ſmooth where the os cuneiforme medium is contiguous to it. The fore-part of this bone is an oblong triangle, for ſuſtaining the os metatarſi of the middle toe.

Os cuneiforme medium, or *minimum*, is ſtill more exactly the ſhape of a wedge than the former. Its upper part is ſquare;——its internal ſide has a flat ſmooth ſurface above and behind, for its conjunction with the following bone; with a ſmall rough foſſa below; and a conſiderable ſhare of it is rough and hollow. The external ſide is ſmooth and a little hollowed, where it is contiguous to the laſt deſcribed bone.——Behind, this bone is triangular, where it is articulated with the os naviculare; and it is alſo triangular at its fore-part, where it is contiguous to the os metatarſi of the toe next to the great one.

Os cuneiforme maximum, or *internum*, differs from the two former in its ſituation, which is more oblique than theirs.—Beſides, its broad thick part is placed below, and the ſmall thin point is above and outwards; while its under broad ſurface is concave, for allowing a ſafe paſſage to the flexor of the great toe.——The ſurface of this os cuneiforme behind,

where it is joined to the os naviculare, is hollow, ſmooth, and of a circular figure below, but pointed above. The external ſide conſiſts of two ſmooth and flat ſurfaces, whoſe direction is nearly at right angles with each other. With the poſterior, that runs obliquely from below forwards and upwards, the os cuneiforme minimum is joined; and with the anterior, whoſe direction is longitudinal, the os metatarſi of the toe next to the great one is connected. The fore-part of this bone is ſemilunar, but flat and ſmooth, for ſuſtaining the os metatarſi of the great toe. The internal ſide is ſcabrous, with two remarkable tubercles below, from which the muſculus abductor pollicis riſes, and the tibialis anticus is inſerted into its upper part.

The three cuneiform bones are all ſo ſecured by ligaments, that very little motion is allowed in any of them, and they are cartilaginous in a foetus of nine months.

The ſeven bones of the *tarſus*, when joined, are convex above, and leave a concavity below, for lodging ſafely the ſeveral muſcles, tendons, veſſels, and nerves, that lie in the ſole of the foot. In the recent ſubject, their upper and lower ſurfaces are covered with ſtrong ligaments, which adhere firmly to them; and all the bones are ſo tightly connected by theſe and the other ligaments, which are fixed to the rough ridges and foſſae mentioned in the preceding deſcription of the particular bones, that notwithſtanding the many ſurfaces covered with cartilage, ſome of which are of the form of the very moveable articulations, no more motion is here allowed, than only to prevent too great a ſhock of the fabric of the body in walking, leaping, &c. by falling on too ſolid a baſe; which, if it was one continued bone, would likewiſe be much more liable to be broken; and, in order to make our foot accommodate itſelf to the ſurfaces we tread on, by becoming more or leſs hollow, or by raiſing or depreſſing

depreſſing either ſide of it, as might be judged by what was ſaid of the particular bones.

Sprains occaſion here, as in the wriſt, great pain and obſtinate tumours, which too often cauſe carious bones.

METATARSUS.

METATARSUS (*a*) is compoſed of five bones, which, in their general characters, agree with the metacarpal bones; but may be diſtinguiſhed from them by the following marks: 1. They are longer, thicker, and ſtronger. 2. Their anterior round ends are not ſo broad, and are leſs in proportion to their baſes. 3. Their bodies are ſharper above and flatter on their ſides, with their inferior ridge inclined more to the outſide. 4. The tubercles at the lower part of the round head are larger.

The firſt or internal metatarſal bone is eaſily diſtinguiſhed from the reſt by its thickneſs. The one next to it is the longeſt, and with its ſharp edge almoſt perpendicular. The others are ſhorter and more oblique, as their ſituation is more external. Which general remarks, with the deſcription I am now to give of each, may teach us to diſtinguiſh them from each other.

Os metatarſi pollicis is by far the thickeſt and ſtrongeſt, as having much the greateſt weight to ſuſtain. Its baſe is oblong, irregularly concave, and of a ſemilunar figure, to be adapted to the os cuneiforme maximum. The inferior edge of this baſe is a little prominent and rough, where the tendon of the peronaeus primus muſcle is inſerted. On its outſide an oblique circular depreſſion is made by the ſecond metatarſal bone. Its round head has generally on

(*a*) Στηθος, πεδιον, Planta, planum, veſtigium, ſolium, pectus, precordium, pectuſculum.

on its fore-part a middle ridge, and two oblong cavities, for the ossa sesamoidea; and on the external side a depression is made by the following bone.

Os metatarsi of the *second toe*, is the longest of the five, with a triangular base supported by the os cuneiforme medium and the external side produced into a process; the end of which is an oblique smooth plane, joined to the os cuneiforme externum.—Near the internal edge of the base, this bone has two small depressions, made by the os cuneiforme maximum, between which is a rough cavity.—Farther forwards we may observe a smooth protuberance, which is joined to the foregoing bone.——On the outside of the base are two oblong smooth surfaces, for its articulation with the following bone; the superior smooth surface being extended longitudinally, and the inferior perpendicularly; between which there is a rough fossa.

Os metatarsi of the *middle toe*, is the second in length.—— Its base, supported by the os cuneiforme externum, is triangular, but slanting outwards, where it ends in a sharp-pointed little process; and the angle below is not completed.

The internal side of this base is adapted to the preceding bone; and the external side has also two smooth surfaces covered with cartilage, but of a different figure; for the upper one is concave, and being round behind, turns smaller as it advances forwards; and the lower surface is little, smooth, convex, and very near the edge of the base.

Os metatarsi of the *fourth* toe, is near as long as the former, with a triangular slanting base joined to the os cuboides, and made round at its external angle; having one hollow smooth surface on the outside, where it is pressed upon by the following bone; and two on the internal side, corresponding to the former bone; behind which is a long narrow surface impressed by the os cuneiforme externum.

Os

Os metatarsi of the *little toe*, is the shortest, situated with its two flat sides above and below, and with the ridges laterally. ——The base of it, part of which rests on the os cuboides, is very large, tuberous, and produced into a long-pointed process externally, where part of the abductor minimi digiti is fixed; and into its upper part the peronaeus secundus is inserted. Its inside has a flat conoidal surface, where it is contiguous to the preceding bone.

When we stand, the fore-ends of these metatarsal bones, and the os calcis, are our only supporters; and therefore it is necessary that they should be strong, and should have a confined motion.

TOES.

The bones of the toes are nearly similar to those of the thumb and fingers; particularly the two of the great toe are precisely formed as the two last of the thumb; only their position, in respect of the other toes, is not oblique; and they are proportionally much stronger, because they are subjected to a greater force; for they sustain the force with which our bodies are pushed forwards by the foot behind at every step we make; and on them principally the weight of the body is supported, when we are raised on our tip-toes.

The three bones in each of the other four toes, compared to those of the fingers, differ from them in these particulars. They are less and smaller in proportion to their lengths; Their bases are much larger than their anterior ends: Their bodies are more narrow above than below, and flatter on the sides. The first phalanx is proportionally much longer than the bones of the second and third, which are very short.

Of the four, the toe next to the great one has the largest bones in all dimensions, and more externally the toes are less.

leſs.——The little toe, and frequently that next to it, have the ſecond and third bones intimately united into one; which may be owing to their little motion, and the great preſſure they are ſubject to.

The toes are of great uſe to us in walking; for, when the ſole is raiſed, they bring our body, with its centre of gravity, perpendicular to the advanced foot.

The bones of the METATARSUS and TOES, are in the ſame condition in children as thoſe of the metacarpus and fingers.

THE only bones now remaining to complete the deſcription of the ſkeleton, are the ſmall ones which are found at the joints of the fingers and toes, and in ſome other parts, called

OSSA SESAMOIDEA.

Theſe are of very different figures and ſizes, though they are generally ſaid to reſemble the ſeed of the ſeſamum.——They ſeem to me nothing elſe than the ligaments of the articulations, or the firm tendons of ſtrong muſcles, or both, become bony by the compreſſion which they ſuffer. Thus the ſeſamoid bones at the beginning of the gaſtrocnemii muſcles, are evidently compoſed of the tendinous fibres only. Theſe, at the firſt joint of the great toe, are as plainly the ſame continued ſubſtance with the ligaments and the tendons of the adductor, flexor, brevis, and abductor.——That which is ſometimes double at the ſecond joint of that toe is part of the capſular ligament; and if we enumerate the other ſeſamoid bones that are at any time found, we may obſerve all of them formed in this manner. Their number, figure, ſituation, and magnitude, are ſo uncertain, that it were in vain to inſiſt on

the

the difference of each; and therefore I ſhall only in general remark,

1. That wherever the tendons and ligaments are firmeſt, the actions of the muſcles ſtrongeſt, and the compreſſion greateſt, there ſuch bones are moſt commonly found.

2. That, *caeteris paribus*, the older the ſubject is in which they are ſought, their number is greater, and their ſize larger.

3. The more labour any perſon is inured to, he has, *caeteris paribus*, the moſt numerous and largeſt oſſa ſeſamoidea.

However, as the two at the firſt joint of the great toe are much larger than any other, are early formed, and are ſeldom wanting in an adult, we may judge, that beſides the more forcible cauſe of their formation, there ſhould alſo be ſome particular advantage neceſſary at this place, rather than elſewhere; which may poſſibly be, to allow the flexor muſcles to ſend their tendons along this joint, ſecure from compreſſion in the hollow between the two oblong ſeſamoid bones; while by removing theſe tendons from the centre of motion, and giving them the advantage of an angle at their inſertion, the force of the muſcle is increaſed; and therefore the great ſuperincumbent weight of our body in progreſſion is more eaſily raiſed.

APPENDIX.

MARKS OF A FEMALE SKELETON.

TO finiſh the deſcription of the bones, is generally to conclude the oſteology: But, that no part of the ſubject may be left untouched, I think it neceſſary to ſubjoin the diſtinguiſhing marks of the Male and Female ſkeletons; and I have choſen to illuſtrate them principally in the latter; becauſe women having a more delicate conſtitution, and affording lodging and nouriſhment to their tender fœtuſes till they have ſufficient ſtrength and firmneſs to bear the injuries of the atmoſphere, and contact of other more ſolid ſubſtances, their bones are frequently incomplete, and always of a make in ſome parts of the body different from thoſe of the robuſt male; which agree to the deſcription already given, unleſs where the proper ſpecialities of the female were particularly remarked; which could not be done in all places where they occur, without perplexing the order of this treatiſe: Therefore I chooſe rather to ſum them up here by way of appendix.

The cauſes of the following ſpecialities of the female bones may be reduced to theſe three: 1. A weak lax conſtitution. 2. A ſedentary unactive life, increaſing that conſtitution. 3. A proper frame for being mothers.

The bones of women are ſmaller in proportion to their length than thoſe of men; becauſe the force of their muſcles is not ſo great, nor is ſuch ſtrong external force applied to them to prevent their ſtretching out in length.

The

The *depreſſions*, *ridges*, *ſcabrous ſurfaces*, and other inequalities made by the *muſcles*, are not ſo conſpicuous in them; becauſe their *muſcles* are neither ſo thick nor ſtrong, nor ſo much employed, as to make ſo ſtrong prints on their bones.

The *os frontis* is more frequently divided by a continuation of the ſagittal ſuture; which depends on the firſt and ſecond general cauſes aſſigned above for the ſpecialties in their bones, as will appear after reflecting on the account given formerly of the middle internal ſpine of this bone.

Their *clavicles* are leſs crooked; becauſe their arms have been leſs forcibly pulled forwards; which, in our European women, eſpecially thoſe of diſtinction, is more hindered by their garb.

Their *ſternum* is more raiſed by long cartilages below, that the thorax might be there widened in ſome proportion to what it is ſhortened by the preſſure upon the diaphragm when they are with child.

The defect of bone, or the hole, in the middle of the *ſternum*, is oftenest found in them; to allow the paſſage of the mammary veſſels, ſay ſome. But, in my opinion, this is owing to a lax conſtitution, by which the oſſification is not ſo ſoon completed as in men, where the action of the ſolids is vigorous, and the circulation of the fluids is briſk; for a much ſmaller hole might have ſerved this purpoſe; and the branches of the internal mammary veſſels which are ſent to the external parts of the thorax, do not paſs here, but between the cartilages of the ribs, before theſe are joined to the ſternum.

The *cartilago xiphoides* is oftener bifurcated in women than in men, for the reaſon aſſigned in the preceding paragraph, viz. a leſs forcible power of oſſification.

The

The ſuperior cartilages of the ribs ſooner oſſify, to ſupport the weight of the mammae.

The middle cartilages are more flat and broad by the weight of the breaſts.

The inferior cartilages are longer, for enlarging the cheſt.

Weak women, who have borne many children when young, often have the *vertebræ* of their back bent forwards, and their *ſternum* depreſſed, or become round-ſhouldered and flat-breaſted (*b*), by the preſſure and weight of the impregnated uterus, and by the ſtrong action of the abdominal muſcles.

The *os ſacrum* is broader, and turned much more backwards, for enlarging the pelvis.

The *os coccygis* is more moveable, and much leſs bent forwards, to facilitate the birth.

The *oſſa ilium* are more hollow, and more reflected outwards, and conſequently further removed from each other, in order to widen the lower part of their abdomen, and for the better ſupport of the impregnated uterus.

The *ridge* on the upper part of the *os pubis* is larger in ſuch women as have borne children, being extended by the ſtrong action of the muſculi recti abdominis.

The *cartilage* between the two oſſa pubis, eſpecially in women who have borne children, is thicker than in men, by which the pelvis is more capacious in females.

The *conjoined ſurfaces* of the *oſſa pubis*, and of the *oſſa innominata* and *ſacrum*, are leſs, the angle under the *ſymphyſis* of the *oſſa pubis* is much larger, and the arches formed below and behind by the *oſſa ilium* and *iſchium* are wider, which, with the ſtraighter *os ſacrum*, and more diſtant *tubera iſchii*, leave a larger paſſage for the excluſion of the child in birth.

The great *tuberoſity* of the *oſſa iſchium* is flatter in women than in men, becauſe it is more preſſed upon in the ſedentary life which females enjoy.

In

(*b*) Cheſelden, Anatomy, book I. chap. 2.

In consequence of the *pelvis* of women being wider, the articulations of their thigh-bones must be farther removed from each other; and therefore a larger space is left for the procreation and birth of children (*c*); which distance of the thighs may be one reason why women in running shuffle more from one side to the other than men, to preserve the centre of gravity of their bodies from falling too far to a side of the joint of the thigh that supports them when the other is raised, which would endanger their tumbling to the ground.

(*c*) Albin, De ossib. § 339.

A

SYSTEM OF ANATOMY.

PART II.

CONTAINING A DESCRIPTION OF THE

HUMAN MUSCLES,

Chiefly as they appear on Diſſection.

Together with their ſeveral Uſes, and the Synonyma of the beſt Authors.

By JOHN INNES.

TO

ALEXANDER MONRO, M. D.

Profeſſor of Anatomy and Medicine, in the Univerſity of Edinburgh.

SIR,

HAVING been repeatedly ſolicited, for ſeveral years paſt, by many of your Pupils, to publiſh a ſhort Deſcription of the Muſcles as they appear upon Diſſection of the Human Body, I have at laſt ventured to comply with their requeſt. Your anxiety to promote the Science of Anatomy, and to encourage every thing that may be uſeful to the gentlemen who attend your Theatre, was the principal motive which induced me to undertake this taſk. I have no knowledge of the ſubject but what I derived from you. If, therefore, this Tre aiſe, which you have never ſeen till I now preſent it to you, ſhould communicate any advantage to the gentlemen under your care, it is to you alone they are indebted for the obligation. I am, Sir, with reſpect, gratitude, and eſteem, your much obliged and very humble ſervant,

EDINBURGH, 1776.

JOHN INNES.

PRE-

PREFACE.

SEVERAL full and accurate deſcriptions of the muſcles have already been publiſhed. But their ſize and prolixity have rendered them of leſs value to the diſſector than the ſmall treatiſe of Dr Douglas, which was firſt publiſhed about the beginning of this century; and ſince that time has undergone various impreſſions, without receiving any improvement, excepting the addition of the ſynonyma from Albinus. It is, therefore, preſumed, that a ſimple and conciſe deſcription of the muſcles, which ſhould contain all the improvements of the moderns, is ſtill wanting.

To claſs the muſcles according to their uſes, may do very well in a large work, or in deſcribing their compound actions; but this method can never anſwer the purpoſes of diſſection. To remedy this inconvenience, the muſcles, in the following Treatiſe, are deſcribed chiefly as they appear in diſſecting the human body.

The deſcribing of the muſcles according to their origins and inſertions prevents much circumlocution. This is the method purſued by Dr Douglas; and wherever his deſcriptions ſeemed tolerably accurate, they have been followed with little alteration. But Dr Douglas's book is peculiarly defective with regard to the muſcles of the abdomen, back, and neck: In deſcribing theſe, therefore, the method of Albinus has been preferred.

Thoſe who have not opportunity, or are averſe from undergoing the labour of diſſecting, may derive conſiderable advantages from comparing the deſcriptions now given with the beautiful and correct tables of Albinus; and, to facilitate ſtill more the ſtudy of theſe intricate organs, I have cauſed eight of Albinus's tables to be publiſhed, with conciſe explanations, on a ſmall ſcale adapted for the pocket.

For the benefit of thoſe who wiſh to examine the hiſtory of the muſcles more minutely, the ſynonyma of the beſt authors are added; and, for the ſake of brevity, the compound action of the muſcles, and the origin and inſertion of ſeveral inconſiderable fibres, are omitted.

The reader will obſerve, that, in general, the muſcles of one ſide only are deſcribed; becauſe all the muſcles of the body, with very few exceptions, have correſpondent ones on the oppoſite ſide.

A

SHORT ACCOUNT

OF THE

AUTHOR,

AND

OF HIS WRITINGS.

MR JOHN INNES, author of the following Treatiſe, was born at Callart, an obſcure village in the Highlands of Scotland. He came to Edinburgh at an early period of life, where he obtained the patronage and protection of Dr Alexander Monro, Profeſſor of Anatomy; who inſtructed him in the knowledge of the human body, and in the art of diſſection. When he was about the age of eighteen years, Dr Monro appointed him diſſector to the Anatomical Theatre. The functions of this important and difficult office, he continued to perform with much reputation for near twenty years.

But his abilities were not confined to the dexterity of diſſecting the moſt minute parts of the human frame. He deſcribed the various organs with eaſe and with perſpicuity. This happy talent attracted the notice of the ſtudents; and, at their ſolicitation, and by the approbation of Dr Monro, he opened an evening courſe of anatomical demonſtrations.

The number of pupils who annually attended theſe demonſtrations afforded the beſt evidence of his abilities, and of the advantages derived from his labours. During his laſt courſe, he was attended by near two hundred ſtudents.

For ſome time before his death, he was troubled with an affection of the lungs, which terminated in a phthiſis pulmonalis, and proved fatal to him on the 12th of January 1777.

On the 15th of the ſame month, the following account of him appeared in the public papers:

" Mr John Innes, at an early period of life, had been educated in the diſſecting art: He made a rapid progreſs in his profeſſion; and his genius and induſtry were rewarded with the privilege of giving private lectures for his own emolument. The utility of his leſſons was ſoon perceived. Numbers

of

of ſtudents reſorted to him for inſtruction; and all of them acknowledged the advantages they had received. At that ſtage of life when men are moſt capable of benefiting themſelves, and of being uſeful to the public, death hurried him out of the world He has given two ſmall ſpecimens of what was to be expected from his anatomical ſkill In the courſe of laſt year, he publiſhed a ſhort deſcription of the Human Muſcles as they appear on diſſection, together with their ſeveral uſes, and the ſynonyma of the beſt authors. The merit of this work was univerſally acknowledged. Some months after, he publiſhed, as a *vade mecum* for ſtudents, eight anatomical tables, containing the principal parts of the ſkeleton and muſcles repreſented in the large tables of Albinus, with accurate explanations. Theſe are all the monuments he has left, by which the public are to judge of his ability. To his numerous friends and acquaintance, it is unneceſſary to mention the warmth of his heart, or the integrity of his diſpoſition."

ADVERTISEMENT.

DURING the illneſs of which Mr INNES died, he put into my hands the firſt edition of his Deſcription of the Muſcles, with a few, chiefly verbal, corrections of it.

ON peruſing that work lately, at the requeſt of the Bookſeller, I have found it neceſſary to make a very conſiderable number of alterations in what relates to the deſcription, as well as to the uſes, of the Muſcles.

EDIN. Sept. 5. 1777.

ALEX. MONRO.

CHAP.

CHAP. I.

Muscles *of the* Teguments *of the* Cranium.

THE ſkin that covers the cranium is moved by a ſingle broad digaſtric muſcle, and one ſmall pair.

1. Occipito Frontalis,

Ariſes fleſhy from the tranſverſe protuberant ridge near the middle of the os occipitis laterally, where it joins with the temporal bone; and tendinous from the reſt of that ridge backwards, oppoſite to the lateral ſinus; it ariſes after the ſame manner on the other ſide: From thence it comes ſtraight forwards, by a broad thin tendon, which covers the upper part of the cranium at each ſide, as low down as the attollens aurem, to which it is connected, as alſo to the zygoma, and covers a part of the aponeuroſis of the temporal muſcles; when it comes as far forwards as near the hair of the front, it becomes fleſhy, and deſcends with ſtraight fibres.

Inſerted into the orbicularis palpebrarum of each ſide, and into the ſkin of the eye-brows, ſending down a fleſhy ſlip between them, as far as the compreſſor naris and levator labii ſuperioris alæque naſi.

Uſe. Pulls the ſkin of the head backwards; raiſes the eye-brows upwards; and, at the ſame time, it draws up and wrinkles the ſkin of the forehead.

Epicranius, Albinus.

Frontalis et occipitalis, Winſlow.

2. Cor.

2. Corrugator Supercilii,

Arises fleshy from the internal angular process of the os frontis, above the joining of the os nasi, and nasal process of the superior maxillary bone; from thence it runs outwards, and a little upwards.

Inserted into the inner and inferior fleshy part of the occipito-frontalis muscle, where it joins with the orbicularis palpebrarum, and extends outwards as far as the middle of the superciliary ridge.

Use. To draw the eye-brow of that side towards the other, and make it project over the inner canthus of the eye: When both act, they pull down the skin of the forehead, and make it wrinkle, particularly between the eye-brows.

Musculus Supercilii, Winslow.

Musculus Frontalis verus, seu Corrugator, Douglas.

CHAP. II.

Of the Muscles *of the* Ear.

THE muscles of the ear may be divided into three classes, viz. the common, proper, and internal. The common move the whole ear; the proper only affect the particular parts to which they are connected; and the internal, the small bones within the tympanum.

The common muscles are,

1. Attollens Aurem,

Arises thin, broad, and tendinous, from the tendon of the occipito-

occipito-frontalis, from which it is almoſt inſeparable, where it covers the aponeuroſis of the temporal muſcle.

Inſerted into the upper part of the ear, oppoſite to the antihelix.

Uſe. To draw the ear upwards, and make the parts into which it is inſerted tenſe.

Superior auris, Winſlow.

2. Anterior Auris,

Ariſes thin and membranous near the poſterior part of the zygoma.

Inſerted into a ſmall eminence on the back of the helix, oppoſite to the concha.

Uſe. To draw this eminence a little forwards and upwards.

3. Retrahentes Auris,

Ariſes, ſometimes by three, but always by two diſtinct ſmall muſcles, from the external and poſterior part of the root of the maſtoid proceſs, immediately above the inſertion of the ſterno-cleido-maſtoid muſcle.

Inſerted into that part of the back of the ear which is oppoſite to the ſeptum that divides the ſcapha and concha.

Uſe. To draw the ear back, and ſtretch the concha.

Poſterior auris, Winſlow.

The proper muſcles are,

1. Helicis Major,

Ariſes from the upper and acute part of the helix anteriorly.

Inſerted into its cartilage a little above the tragus.

Uſe.

Uſe. To depreſs that part from which it ariſes a little downwards and forwards.

2. Helicis Minor,

Ariſes from the inferior and anterior part of the helix.

Inſerted into the crus of the helix, near the fiſſure in the cartilage oppoſite to the concha.

Uſe. To contract the fiſſure.

3. Tragicus,

Ariſes from the middle and outer part of the concha, at the root of the tragus, along which it runs.

Inſerted into the point of the tragus.

Uſe. Pulls the point of the tragus a little forwards.

4. Antitragicus,

Ariſes from the internal part of the cartilage that ſupports the antitragus; and, running upwards, is

Inſerted into the tip of the antitragus, as far as the inferior part of the antihelix, where there is a fiſſure in the cartilage.

Uſe. Turns the tip of the antitragus a little outwards, and depreſſes the extremity of the antihelix towards it.

5. Transversus Auris,

Ariſes from the prominent part of the concha on the dorſum of the ear; the fibres not ſo fleſhy as in the former.

Inſerted oppoſite to the outer ſide of the antihelix.

Uſe. Draws the parts to which it is connected towards each other, and ſtretches the ſcapha and concha.

The muſcles of the internal ear are three:

1. Laxator Tympani,

Ariſes by a ſmall beginning from the extremity of the ſpinous

ſpinous proceſs of the ſphenoid bone, behind the entry of the artery of the dura mater; then runs backwards, and a little upwards, along with the nerve called *chorda tympani*, in a fiſſure of the os temporis near the foſſa that lodges the condyle of the lower jaw.

Inſerted into the long proceſs of the malleus, within the tympanum, where it reſts upon the edge of the fiſſure between the pars ſquamoſa and petroſa.

Uſe. To draw the malleus obliquely forwards towards its origin, and conſequently the membrana tympani; by which means that membrane is made leſs concave, or is relaxed.

Externus mallei, Albinus. *Anterior mallei*, Winſlow.

Obliquus auris, Douglas.

2. Tensor Tympani,

Ariſes by a very ſmall beginning, from the cartilaginous extremity of the Euſtachian tube, juſt where it begins to be covered by the pars petroſa, and ſpinous proceſs of the ſphenoid bone, near the entry of the artery of the dura mater; from thence running backwards near the oſſeous part of the Euſtachian tube, it forms a very diſtinct fleſhy belly, below a thin oſſeous plate, between the pars ſquamoſa and labyrinth; and ſends off a ſlender tendon, which makes a turn into the tympanum along with the nerve called *chorda tympani*.

Inſerted into the poſterior part of the handle of the malleus, a little lower than the root of its long proceſs.

Uſe. To pull the malleus and membrana tympani inwards towards the pars petroſa, by which the membrane is made more concave and tenſe.

Internus mallei, Winſlow. *Internus auris*, Douglas.

3. Stapedius,

Ariſes, by a ſmall fleſhy belly, from a little cavern in the

the pars petrosa, near the cells of the mastoid process, before the inferior part of the passage for the portio dura of the auditory nerve; its tendon passes straight through a small round hole in the same cavern, enters the anterior part of the tympanum, and is

Inserted into the posterior part of the head of the stapes.

Use. To draw the stapes obliquely upwards towards the cavern, by which the posterior part of its base is moved inwards, and the anterior part outwards.

Musculus stapedis, Winslow. *Stapidaeus*, Douglas.

CHAP. III.

Of the MUSCLES *of the* EYE-LIDS.

THE palpebrae or eye-lids, have one muscle common to both, and the upper eye-lid one proper to itself.

1. ORBICULARIS PALPEBRARUM.

Arises, by a number of fleshy fibres, from the outer edge of the orbitar process of the superior maxillary bone, and from a tendon near the inner angle of the eye: these run a little downwards, then outwards, over the upper part of the cheek, below the orbit, covering the under eye-lid, and surround the external angle, being loosely connected only to the skin and fat: run over the superciliary ridge of the os frontis, towards the inner canthus, where they intermix with those of the occipito frontalis and corrugator supercilii; then covering the upper eye-lid, they descend to the inner angle opposite to the inferior origin of this muscle, firmly adhering to the internal angular process of the os frontis, and to the

short

ſhort round tendon which ſerves to fix the palpebrae and muſcular fibres ariſing from it.

Inſerted, by the ſhort round tendon, into the naſal proceſs of the ſuperior maxillary bone, covering the anterior and upper part of the lachrymal ſac; which tendon can be eaſily felt at the inner canthus of the eye.

Uſe. To ſhut the eye, by drawing both lids cloſe together, the fibres contracting from the outer angle towards the inner, preſs the eye-ball, ſqueeze the lachrymal gland, and convey the tears towards the puncta lachrymalia.

The *ciliaris* of ſome authors is only a part of this muſcle covering the cartilages of the eye-lids, called *cilia* or *tarſi*.

There is often a ſmall fleſhy ſlip which runs down from the outer and inferior part of this muſcle above the zygomaticus minor, and joins with the levator labii ſuperioris alaeque naſi.

2. Levator Palpebrae Superioris,

Ariſes from the upper part of the foramen opticum of the ſphenoid bone, through which the optic nerve paſſes, above the levator oculi, near the trochlearis muſcle.

Inſerted, by a broad thin tendon, into the cartilage that ſupports the upper eye-lid, named *tarſus*.

Uſe. To open the eye, by drawing the eye-lid upwards; which it does completely, by being fixed to the tarſus, pulling it below the eye-brow, and within the orbit.

Aperiens palpebram rectus, Douglas.

CHAP.

CHAP. IV.

Muscles *of the* Eye-Ball.

THE muſcles which move the globe of the eye are ſix, viz.

Four *ſtraight* and two *oblique*.

The four ſtraight muſcles very much reſemble each other: all

Ariſing by a narrow beginning, a little tendinous and fleſhy, from the bottom of the orbit around the foramen opticum of the ſphenoid bone, where the optic nerve enters, ſo that they may be taken out adhering to this nerve; and all having ſtrong fleſhy bellies.

Inſerted at the fore-part of the globe of the eye into the anterior part of the tunica ſclerotica, and under the tunica adnata, at oppoſite ſides, which indicates both their names and *Uſe*; ſo that they ſcarcely require any further deſcription, but to name them ſingly.

1. Levator Oculi,

Ariſes from the upper part of the foramen opticum of the ſphenoid bone, below the levator palpebrae ſuperioris; and runs forwards to be

Inſerted into the ſuperior and fore-part of the tunica ſclerotica, by a broad thin tendon.

Uſe. To raiſe up the globe of the eye.

Attolens, Albinus. *Elevator*, Douglas.

2. De-

2. Depressor Oculi,

Arises from the inferior part of the foramen opticum.

Inserted oppoſite to the former.

Uſe. To pull the globe of the eye down.

Deprimens, Albinus.

3. Adductor Oculi,

Ariſes, as the former, between the obliquus ſuperior and depreſſor, being, from its ſituation, the ſhorteſt.

Inſerted oppoſite to the inner angle.

Uſe. To turn the eye towards the noſe.

4. Abductor Oculi,

Ariſes from the bony partition between the foramen opticum and lacerum, being the longeſt from its ſituation; and is

Inſerted into the globe oppoſite to the outer canthus.

Uſe. To move the globe outwards.

The oblique muſcles are two:

Obliquus Superior, *ſeu* Trochlearis,

Ariſes, like the ſtraight muſcles, from the edge of the foramen opticum at the bottom of the orbit, between the levator and adductor oculi; from thence runs ſtraight along the pars plana of the ethmoid bone to the upper part of the orbit, where a cartilaginous trochlea is fixed to the inſide of the internal angular proceſs of the os frontis, through which its tendon paſſes, and runs a little downwards and outwards, incloſed in a looſe membranous ſheath.

Inſerted, by a broad thin tendon, into the tunica ſclerotica, about half way between the inſertion of the attolens oculi and optic nerve.

Uſe.

Uſe. To roll the globe of the eye, and turn the pupil downwards and outwards, ſo that the upper ſide of the globe is turned inwards, and the inferior part to the outſide of the orbit, and the whole globe drawn forwards towards the inner canthus.

Obliquus major, Winſlow.

2. OBLIQUUS INFERIOR,

Ariſes, by a narrow beginning, from the outer edge of the orbitar proceſs of the ſuperior maxillary bone, near its juncture with the os unguis; and running obliquely outwards, is

Inſerted into the ſclerotica, in the ſpace between the abductor and optic nerve, by a broad thin tendon.

Uſe. To draw the globe of the eye forwards, inwards, and downwards; and, contrary to the ſuperior, to turn the pupil upwards, towards the inner extremity of the eye-brow; at the ſame time, the external part of the globe is turned towards the inferior ſide, and the internal rolls towards the upper part.

Obliquus minor, Winſlow.

CHAP. V.

Of the MUSCLES *of the* NOSE.

THERE is only one muſcle on each ſide that can be called proper to the noſe, though it is affected by ſeveral muſcles of the face.

COMPRESSOR NARIS,

Ariſes, by a narrow beginning, from the root of the ala naſi externally, where part of the levator labii ſuperioris alaeque

alaeque naſi is connected to it; it ſpreads into a number of thin ſeparate fibres, which run up along the cartilage in an oblique manner towards the dorſum of the noſe, where it joins with its fellow, and is

Inſerted ſlightly into the anterior extremity of the os naſi and naſal proceſs of the ſuperior maxillary bone, where it meets with ſome of the fibres deſcending from the occipito-frontalis muſcle.

Uſe. To compreſs the ala towards the ſeptum naſi, particularly when we want to ſmell acutely; but, if the fibres of the frontal muſcle, which adhere to it, act, the upper part of this thin muſcle aſſiſts to pull the ala outwards. It alſo corrugates the ſkin of the noſe, and aſſiſts in expreſſing certain paſſions.

Rinaeus, vel naſalis, Douglas.

CHAP. VI.

Muscles *of the* Mouth *and* Lips.

THE mouth has nine pair of muſcles, which are inſerted into the lips, and a common one formed by the termination of theſe, viz. three *above*, three *below*, three *outwards*, and the common muſcle ſurrounds the mouth.

The three above are,

1. Levator Anguli Oris,

Ariſes, thin and fleſhy, from the hollow of the ſuperior maxillary

maxillary bone, between the root of the ſocket of the firſt dens molaris and the foramen infra-orbitarium.

Inſerted into the angle of the mouth and under lip, where it joins with its antagoniſt.

Uſe. To draw the corner of the mouth upwards, and make that part of the cheek oppoſite to the chin prominent, as in ſmiling.

Elevator labiorum communis, Douglas.

Caninus, Winſlow.

2. Levator Labii Superioris Alaeque Nasi,

Ariſes by two diſtinct origins; the firſt broad and fleſhy, from the external part of the orbitar proceſs of the ſuperior maxillary bone which forms the lower part of the orbit, immediately above the foramen infra orbitarium; the ſecond portion ariſes from the naſal proceſs of the ſuperior maxillary bone, where it joins the os frontis at the inner canthus, deſcending along the edge of the groove for the lacrymal ſac. The firſt and ſhorteſt portion is

Inſerted into the upper lip and orbicularis labiorum; the ſecond and longeſt, into the upper lip and outer part of the ala naſi.

Uſe. To raiſe the upper lip towards the orbit, and a little outwards; the ſecond portion ſerves to draw the ſkin of the noſe upwards and outwards, by which the noſtril is dilated.

Elevator labii ſuperioris proprius, Douglas.

Inciſivus lateralis, Firſt portion; *Pyramidalis*, Second portion; Winſlow.

3. Depressor Labii Superioris Alaeque Nasi.

Ariſes, thin and fleſhy, from the os maxillare ſuperius, immediately above the joining of the gums with the two dentes

dentes incisivi, and the dens caninus: from thence it runs up under part of the levator labii superioris alaeque nasi.

Inserted into the upper lip and root of the ala nasi.

Use. To draw the upper lip and ala nasi downwards and backwards.

Depressor alae nasi, Albinus.

Incisivus medius, Winslow.

Depressor labii superioris proprius, Douglas.

The three below are,

1. Depressor Anguli Oris,

Arises, broad and fleshy, from the lower edge of the maxilla inferior, at the side of the chin, being firmly connected to that part of the platysma myoides, which runs over the maxilla to the angle of the mouth, to the depressor labii inferioris within, and to the skin and fat without, gradually turning narrower; and is

Inserted into the angle of the mouth, joining with the zygomaticus major and levator anguli oris.

Use. To pull down the corner of the mouth.

Triangularis, Winslow.

Depressor labiorum communis, Douglas.

2. Depressor Labii Inferioris,

Arises, broad and fleshy, intermixed with fat, from the inferior part of the lower jaw next the chin; runs obliquely upwards; and is

Inserted into the edge of the under lip, extends along one half of the lip, and is lost in its red part.

Use. To pull the under lip and the skin of the side of the chin downwards, and a little outwards.

Quadratus, Winslow.

Depressor labii inferioris proprius, Douglas.

3. Le-

3. Levator Labii Inferioris,

Arises, from the lower jaw, at the roots of the alveoli of two dentes incisivi and of the caninus; is

Inserted into the under lip and skin of the chin.

Use. To pull the parts into which it is inserted upwards.

Levator menti, Albinus.

Incisivus inferior, Winslow.

Elevator labii inferioris proprius, Douglas.

The three outward are,

1. Buccinator,

Arises, tendinous and fleshy, from the lower jaw, as far back as the last dens molaris and fore-part of the root of the coronoid process; fleshy from the upper jaw, between the last dens molaris and pterygoid process of the sphenoid bone; from the extremity of which it arises tendinous, being continued between both jaws to the constrictor pharyngis superior, with which it joins; from thence proceeding with straight fibres, and adhering close to the membrane that lines the mouth, it is

Inserted into the angle of the mouth within the orbicularis oris.

Use. To draw the angle of the mouth backwards and outwards, and to contract its cavity, by pressing the cheek inwards, by which the food is thrust between the teeth.

Retractor anguli oris, Albinus.

2. Zygomaticus Major,

Arises, fleshy, from the os malae, near the zygomatic suture.

Inserted into the angle of the mouth, appearing to be lost in the depressor anguli oris and orbicularis oris.

Use.

Uſe. To draw the corner of the mouth and under lip towards the origin of the muſcle, and make the cheek prominent as in laughing.

Zygomaticus, Douglas.

3. Zygomaticus Minor,

Ariſes from the upper prominent part of the os malæ above the origin of the former muſcle; and, deſcending obliquely downwards and forwards, is

Inſerted into the upper lip, near the corner of the mouth, along with the levator anguli oris.

Uſe. To draw the corner of the mouth obliquely outwards, and upwards, towards the external canthus of the eye.

The common muſcle is the

Orbicularis Oris.

This muſcle is, in a great meaſure, formed by the muſcles that move the lips; the fibres of the ſuperior deſcending, thoſe of the inferior aſcending, and, decuſſating each other about the corner of the mouth, run along the lip to join thoſe of the oppoſite ſide, ſo that the fleſhy fibres appear to ſurround the mouth like a ſphincter.

Uſe. To ſhut the mouth, by contracting and drawing both lips together, and to counteract all the muſcles that aſſiſt in forming it.

Sphincter labiorum, Douglas.

Semi orbicularis, Winſlow.

Conſtrictor oris, Cowper.

There is another ſmall muſcle deſcribed by Albinus, which he calls *Naſalis labii ſuperioris;* but it ſeems to be only ſome fibres of the former connected to the ſeptum naſi.

CHAP.

CHAP. VII.

Muscles *of the* Lower Jaw.

THE lower jaw has four pair of muſcles for its elevation or lateral motions, viz. two, which are ſeen on the ſide of the face, and two concealed by the angle of the jaw.

1. Temporalis,

Ariſes, fleſhy, from a ſemicircular ridge of the lower and lateral part of the parietal bone, from all the pars ſquamoſa of the temporal bone, from the external angular proceſs of the os frontis, from the temporal proceſs of the ſphenoid bone, and from an aponeuroſis which covers it; from theſe different origins the fibres deſcend like radii towards the jugum, under which they paſs; and are

Inſerted, by a ſtrong tendon, into the upper part of the coronoid proceſs of the lower jaw; in the duplicature of which tendon this proceſs is incloſed as in a ſheath, being continued down all its fore-part to near the laſt dens molaris.

Uſe. To pull the lower jaw upwards, and preſs it againſt the upper, at the ſame time drawing it a little backwards.

N. B. This muſcle is covered by a tendinous membrane, called its *aponeuroſis*, which ariſes from the bones that give origin to the upper and ſemicircular part of the muſcle; and deſcending

deſcending over it, is inſerted into all the jugum, and the adjoining part of the os frontis.

The uſe of this membrane is to give room for the origin of a greater number of fleſhy fibres, to fortify the muſcle in its action, and to ſerve as a defence to it.

Crotaphite muſcle, Winſlow.

2. Masseter,

Ariſes, by ſtrong, tendinous, and fleſhy fibres, which run in different directions, from the ſuperior maxillary bone, where it joins the os malæ, and from the inferior and anterior part of the zygoma, its whole length, as far back as the tubercle before the ſocket for the condyle of the lower jaw; the external fibres ſlanting backwards, and the internal forwards.

Inſerted into the angle of the lower jaw, and from that upwards to near the top of its coronoid proceſs.

Uſe. To pull the lower to the upper jaw, and by means of its oblique decuſſation, a little forwards and backwards.

3. Pterygoideus Internus,

Ariſes, tendinous and fleſhy, from the inner and upper part of the internal plate of the pterygoid proceſs, filling all the ſpace between the two plates; and from the pterygoid proceſs of the os palati between theſe plates.

Inſerted into the angle of the lower jaw internally.

Uſe. To draw the jaw upwards, and obliquely towards the oppoſite ſide.

Pterygoideus major, Winſlow.

4. Pterygoideus Externus,

Ariſes, from the outer ſide of the external plate of the pterygoid proceſs of the ſphenoid bone, from part of the tuberoſity

tuberosity of the os maxillare adjoining to it, and from the root of the temporal process of the sphenoid bone.

Inserted into a cavity in the neck of the condyloid process of the lower jaw; some of its fibres are inserted into the ligament that connects the moveable cartilage and that process to each other.

Use. To pull the lower jaw forwards, and to the opposite side; and to pull the ligament from the joint, that it may not be pinched during these motions: When both external pterygoid muscles act, the fore-teeth of the under jaw are pushed forwards beyond those of the upper jaw.

Pterygoideus minor, Winslow.

CHAP. VIII.

The MUSCLES *which appear about the anterior part of the* NECK.

ON the side of the neck are two muscles or layers.

1. MUSCULUS CUTANEUS, VULGO PLATYSMA MYOIDES,

Arises, by a number of slender separate fleshy fibres, from the cellular substance that covers the upper parts of the deltoid and pectoral muscles; in their ascent, they all unite to form a thin muscle, which runs obliquely upwards along the side of the neck, adhering to the skin.

Inserted into the lower jaw, between its angle and the origin of the depressor anguli oris, to which it is firmly con-

nected, and but slightly to the skin that covers the inferior part of the masseter muscle and parotid glands.

Use. To assist the depressor anguli oris in drawing the skin of the cheek downwards; and when the mouth is shut, it draws all that part of the skin, to which it is connected, below the lower jaw, upwards.

Platysma myoides, Galen.

Musculus cutaneus, Winslow.

Quadratus genæ, vel Latissimus colli, Douglas.

Latissimus colli, Albinus.

2. STERNO-CLEIDO-MASTOIDEUS,

Arises, by two distinct origins; the anterior, tendinous and a little fleshy, from the top of the sternum near its junction with the clavicle; the posterior, fleshy, from the upper and anterior part of the clavicle; both unite a little above the anterior articulation of the clavicle, to form one muscle, which runs obliquely upwards and outwards, to be

Inserted, by a thick strong tendon, into the mastoid process, which it surrounds; and, gradually turning thinner, is inserted as far back as the lambdoid suture.

Use. To turn the head to one side, and bend it forwards.

Sterno-mastoideus and *Cleido-mastoideus*, Albinus.

Mastoideus, Douglas.

CHAP.

CHAP. IX.

Muscles *situated between the* Lower Jaw *and* Os Hyoides.

THERE are four layers before, and two muscles at the side.

The four layers are,

1. Digastricus,

Arises, by a fleshy belly, intermixed with tendinous fibres, from the fossa at the root of the mastoid process of the temporal bone, and soon becomes tendinous; runs downwards and forwards: The tendon passes generally through the stylo-hyoideus muscle; then it is fixed by a ligament to the os hyoides: And having received from that bone an addition of tendinous and muscular fibres, runs obliquely forwards, turns fleshy again, and is

Inserted, by its anterior belly, into a rough sinuosity at the inferior and anterior edge of that part of the lower jaw called the *chin*.

Use. To open the mouth, by pulling the lower jaw downwards, and backwards; and, when the jaws are shut, to raise the larynx, and consequently the pharynx, upwards, as in deglutition.

Biventer maxillae inferioris, Albinus.

2. Mylo-Hyoideus,

Arises, fleshy, from all the inside of the lower jaw, between

tween the laſt dens molaris and the middle of the chin, where it joins with its fellow.

Inſerted into the lower edge of the baſis of the os hyoides, and joins with its fellow.

Uſe. To pull the os hyoides forwards, upwards, and to a ſide.

3. Genio-Hyoideus,

Ariſes, tendinous, from a rough protuberance in the middle of the lower jaw internally, or on the inſide of the chin.

Inſerted into the baſis of the os hyoides.

Uſe. To draw this bone forwards to the chin.

4. Genio-Hyo-Glossus,

Ariſes, tendinous, from a rough protuberance in the ſide of the middle of the lower jaw; its fibres run, like a fan, forwards, upwards, and backwards; and are

Inſerted into the tip, middle, and root of the tongue, and baſe of the os hyoides, near its cornu.

Uſe. According to the direction of its fibres, to draw the tip of the tongue backwards into the mouth, the middle downwards, and to render its dorſum concave; to draw its root and os hyoides forwards, and to thruſt the tongue out of the mouth.

The two muſcles at the ſide are,

1. Hyo-Glossus,

Ariſes, broad and fleſhy, from the baſe, cornu, and appendix of the os hyoides; the fibres run upwards and outwards, to be

Inſerted into the ſide of the tongue, near the ſtylo-gloſſus.

Uſe. To pull the tongue inwards and downwards.

Baſio-cerato-chondro-gloſſus, Albinus.

Cerato-gloſſus, Douglas.

2. Lin-

2. Linguālis,

Ariſes, from the root of the tongue laterally; runs forwards between the hyo-gloſſus and genio-gloſſus, to be

Inſerted into the tip of the tongue, along with part of the ſtylo-gloſſus.

Uſe. To contract the ſubſtance of the tongue, and bring it backwards.

CHAP. X.

Muscles *ſituated between the* Os Hyoides *and* Trunk.

THESE may be divided into two layers.

The firſt layer conſiſts of two muſcles.

1. Sterno-Hyoideus,

Ariſes, thin and fleſhy, from the cartilaginous extremity of the firſt rib, the upper and inner part of the ſternum, and from the clavicle where it joins with the ſternum.

Inſerted into the baſe of the os hyoides.

Uſe. To pull the os hyoides downwards.

2. Omo-Hyoideus,

Ariſes, broad, thin, and fleſhy, from the ſuperior coſta of the ſcapula, near the ſemilunar nitch, and from the ligament that runs acroſs it; thence aſcending obliquely, it becomes tendinous below the ſterno-cleido-maſtoid muſcle; and growing fleſhy again, is

Inſerted into the baſe of the os hyoides, between its cornu and the inſertion of the ſterno-hyoides.

Uſe.

Use. To pull the os hyoides obliquely downwards.

Coraco-hyoideus, Albinus and Douglas.

The second layer consists of three muscles.

1. STERNO-THYROIDEUS,

Arises, fleshy, from the whole edge of the uppermost bone of the sternum internally, opposite to the cartilage of the first rib, from which it receives a small part of its origin.

Inserted into the surface of the rough line at the external part of the inferior edge of the thyroid cartilage.

Use. To draw the larynx downwards.

2. THYREO-HYOIDEUS,

Arises from the rough line, opposite to the former.

Inserted into part of the bases, and almost all the cornu of the os hyoides.

Use. To pull the os hyoides downwards, or the thyroid cartilage upwards.

Thyro-hyoideus vel Hyo-thyroideus, Winslow.

3. CRICO-THYROIDEUS,

Arises from the side and fore-part of the cricoid cartilage, running obliquely upwards.

Inserted by two portions; the first, into the lower part of the thyroid cartilage; the second into its inferior cornu.

Use. To pull forwards and depress the thyroid, or to elevate and draw backwards the cricoid cartilage.

CHAP.

CHAP. XI.

Muscles *situated between the* Lower Jaw *and* Os Hyoides *laterally.*

THEY are five in number. Three proceed from the ſtyloid proceſs of the temporal bone, from which they have half of their names; and two from the pterygoid proceſs of the ſphenoid bone.

The three from the ſtyloid proceſs are,

1. Stylo-Glossus,

Ariſes, tendinous and fleſhy, from the ſtyloid proceſs, and from a ligament that connects that proceſs to the angle of the lower jaw.

Inſerted into the root of the tongue, runs along its ſide, and is inſenſibly loſt near its tip.

Uſe. To draw the tongue laterally and backwards.

2. Stylo-Hyoideus,

Ariſes by a round tendon, from the middle and inferior part of the ſtyloid proceſs.

Inſerted into the os hyoides at the junction of the baſe and cornu.

Uſe. To pull the os hyoides to one ſide, and a little upwards.

N. B. Its fleſhy belly is generally perforated by the tendon of the digaſtric muſcle, on one or both ſides.——

There

There is often another accompanying it, called *ſtylo-hyoideus alter*; and has the ſame origin, inſertion, and uſe.

3. Stylo-Pharyngeus,

Ariſes, fleſhy, from the root of the ſtyloid proceſs.

Inſerted into the ſide of the pharynx and back-part of the thyroid cartilage.

Uſe. To dilate and raiſe the pharynx and thyroid cartilage upwards.

The two from the pterygoid proceſs are,

1. Circumflexus, or Tensor Palati,

Ariſes from the ſpinous proceſs of the ſphenoid bone, behind the foramen ovale, which tranſmits the third branch of the fifth pair of nerves; from the Euſtachian tube, not far from its oſſeous part: it then runs down along the pterygoideus internus, paſſes over the hook of the internal plate of the pterygoid proceſs by a round tendon, which ſoon ſpreads into a broad membrane.

Inſerted into the velum pendulum palati, and the ſemilunar edge of the os palati, and extends as far as the ſuture which joins the two bones. Generally ſome of its poſterior fibres join with the conſtrictor pharyngis ſuperior, and palato-pharyngeus.

Uſe. To ſtretch the velum, to draw it downwards, and to a ſide towards the hook. It has little effect upon the tube, being chiefly connected to its oſſeous part.

Circumflexus palati, Albinus.

Spheno-ſalpingo ſtaphylinus, ſeu *Staphylinus externus*, Winſlow.

Muſculus tubae novus, Valſalva; vel *Palato-ſalpingeus*, Douglas.

2. Levator

2. Levator Palati,

Ariſes, tendinous and fleſhy, from the extremity of the pars petroſa of the temporal bone, where it is perforated by the Euſtachian tube, and alſo from the membranous part of the ſame tube.

Inſerted into the whole length of the velum pendulum palati, as far as the root of the uvula, and unites with its fellow.

Uſe. To draw the velum upwards and backwards, ſo as to ſhut the paſſage from the fauces into the mouth and noſe.

Levator palati mollis, Albinus.

Petro-ſalpingo-ſtaphylinus, vel *Salpingo ſtaphylinus internus*, *vulgo*, Winſlow.

Salpingo-ſtaphylinus, Valſalva. *Pterigo-ſtaphylinus externus*, *vulgo*, Douglas.

Spheno-ſtaphylinus, Cowper.

Previous to the deſcription of the muſcles ſituated about the paſſage into the throat, it will be neceſſary to mention the principal parts to which they are connected.

Upon looking into any perſon's mouth, when wide opened, we ſee a ſoft curtain hanging from the palate bones, named *velum pendulum palati.* In the middle of which we likewiſe obſerve a papilla projecting from the velum, named *uvula*, or *pap of the throat.* From each ſide of the uvula, at its root, two arches, or columns are ſent down; the anterior to the root of the tongue, the poſterior to the pharynx. Between theſe arches, on each ſide, the cellular glands, called *amygdalae*, or *almonds of the ears*, are ſituated.

The common opening behind the anterior arch may be named *fauces*, or *top of the throat*; from which there are ſix paſſages, viz. two upwards, being one to each noſtril; two

at the ſides, or one to each ear, called the *Euſtachian tubes*: two downwards; the anterior is the paſſage through the *glottis* and *larynx*, into the *trachea*, which terminates in the lungs; the poſterior is the largeſt, named *pharynx*, or *top of the oeſophagus*, which leads to the ſtomach.

CHAP. XII.

Muscles *ſituated about the entry to the* Fauces.

THERE are two on each ſide, and a ſingle one in the middle.

The two on each ſide are,

1. Constrictor Isthmi Faucium,

Ariſes, by a ſlender beginning, from the ſide of the tongue, near its root; from thence running upwards within the anterior arch, before the amygdala; it is

Inſerted into the middle of the velum pendulum palati, at the root of the uvula anteriorly, being connected with its fellow, and with the beginning of the palato pharyngeus.

Uſe. Draws the velum toward the root of the tongue, which it raiſes at the ſame time, and, with its fellow, contracts the paſſage between the two arches, by which it ſhuts the opening into the fauces.

Gloſſo ſtaphylinus, Winſlow and Douglas.

2. Palato-Pharyngeus,

Ariſes, by a broad beginning, from the middle of the velum

velum pendulum palati, at the root of the uvula posteriorly, and from the tendinous expansion of the circumflexus palati. The fibres are collected within the posterior arch behind the amygdalae, and run backwards to the top and lateral part of the pharynx, where the fibres are scattered, and mix with those of the stylo-pharyngeus.

Inserted into the edge of the upper and back-part of the thyroid cartilage; some of its fibres being lost between the membrane of the pharynx, and the two inferior constrictors.

Use. Draws the uvula and velum downwards and backwards; and at the same time pulls the thyroid cartilage and pharynx upwards, and shortens it; with the constrictor superior and tongue, it assists in shutting the passage into the nostrils; and, in swallowing, it thrusts the food from the fauces into the pharynx.

Thyro staphylinus, Douglas.

Thyro pharyngo-staphylinus, Winslow.

Salpingo Pharyngeus of Albinus is composed of a few fibres of this muscle, which

Arise from the anterior and lower part of the cartilaginous extremity of the Eustachian tube; and are

Inserted into the inner part of the last-mentioned muscle.

Use. To assist the former, and to dilate the mouth of the tube.

The one in the middle is the

Azygos Uvulae,

Arises, fleshy, from the extremity of the suture which joins the palate-bones; runs down the whole length of the velum

velum and uvula, reſembling a ſmall earth-worm, and adhering to the tendons of the circumflexi.

Inſerted into the tip of the uvula.

Uſe. Raiſes the uvula upwards and forwards, and ſhortens it.

Palato-ſtaphylinus, Douglas.

Staphylinus, or *Epiſtaphylinus*, Winſlow.

CHAP. XIII.

MUSCLES *ſituated on the poſterior part of the* PHARYNX.

OF theſe there are three pair.

1. CONSTRICTOR PHARYNGIS INFERIOR,

Ariſes from the ſide of the thyroid cartilage, near the attachment of the ſterno-hyoideus and thyreo-hyoideus muſcles; and from the cricoid cartilage, near the crico-thyroideus. This muſcle is the largeſt of the three; and is

Inſerted into the white line, where it joins with its fellow; the ſuperior fibres running obliquely upwards, covering nearly one half of the middle conſtrictor, and terminating in a point; the inferior fibres run more tranſverſely, and cover the beginning of the oeſophagus.

Uſe. To compreſs that part of the pharynx which it covers, and to raiſe it with the larynx a little upwards.

Thyro-pharyngeus, *Crico-pharyngeus*, Douglas.

2. CON-

2. Constrictor Pharyngis Medius,

Arises from the appendix of the os hyoides, from the cornu of that bone, and from the ligament which connects it to the thyroid cartilage; the fibres of the superior part running obliquely upwards, and, covering a considerable part of the superior constrictor, terminate in a point.

Inserted into the middle of the cuneiform process of the os occipitis, before the foramen magnum, and joined to its fellow at a white line in the middle back-part of the pharynx. The fibres at the middle part run more transversely than those above or below.

Use. To compress that part of the pharynx which it covers, and to draw it and the os hyoides upwards.

Hyo pharyngeus, *Syndesmo pharyngeus*, Douglas.

3. Constrictor Pharyngis Superior,

Arises, above, from the cuneiform process of the os occipitis, before the foramen magnum, near the holes where the ninth pair of nerves passes out; lower down, from the pterygoid process of the sphenoid bone; from the upper and under jaw, near the roots of the last dentes molares; and between the jaws, it is continued with the buccinator muscle; and with some fibres from the root of the tongue, and from the palate.

Inserted into a white line in the middle of the pharynx, where it joins with its fellow, and is covered by the constrictor medius.

Use. To compress the upper part of the pharynx, and draw it forwards and upwards.

Cephalo-pharyngeus, *Pterygo-pharyngeus*, *Mylo-pharyngeus*, *Glosso-pharyngeus*, Douglas.

CHAP.

CHAP. XIV.

Muscles *situated about the* Glottis.

THEY consist generally of four pair of small muscles, and a single one.

1. Crico-Arytaenoideus Posticus,

Arises, fleshy, from the back-part of the cricoid cartilage, and is

Inserted into the posterior part of the base of the arytenoid cartilage.

Use. To open the rima glottidis a little, and, by pulling back the arytenoid cartilage, to stretch the ligament so as to make it tense.

2. Crico-Arytaenoideus Lateralis,

Arises, fleshy, from the cricoid cartilage, laterally, where it is covered by part of the thyroid, and is

Inserted into the side of the base of the arytenoid cartilage near the former.

Use. To open the rima glottidis, by pulling the ligaments from each other.

3. Thyreo-Arytaenoideus,

Arises from the under and back-part of the middle of the thyroid cartilage; and, running backwards and a little upwards, along the side of the glottis, is

Inserted into the arytenoid cartilage, higher up and farther forwards than the crico-arytaenoideus lateralis.

Use.

Uſe. To pull the arytenoid cartilage forwards nearer to the middle of the thyroid, and conſequently to ſhorten and relax the ligament of the larynx or glottis vera.

4. Arytænoideus Obliquus,

Ariſes from the baſe of one arytenoid cartilage; and, croſſing its fellow, is

Inſerted near the tip of the other arytenoid cartilage.

Uſe. When both act, they pull the arytenoid cartilages towards each other.

N. B. One of theſe is very often wanting.

Arytaenoideus minor, Douglas.

The ſingle muſcle is the

Arytænoideus Transversus,

Ariſes from the ſide of one arytenoid cartilage, from near its articulation with the cricoid to near its tip. The fibres run ſtraight acroſs, and are

Inſerted, in the ſame manner, into the other arytenoid cartilage.

Uſe. To ſhut the rima glottidis, by bringing theſe two cartilages, with the ligaments, nearer one another.

Arytaenoideus major, Douglas.

Beſides theſe, there are a few ſeparate muſcular fibres on each ſide; which, from their general direction, are named,

1. Thyreo-Epiglottideus.

Ariſes, by a few pale ſeparated fibres, from the thyroid cartilage; and is

Inſerted into the epiglottis laterally.

Uſe. To draw the epiglottis obliquely downwards, or, when

when both act, directly downwards; and, at the same time, it expands that soft cartilage.

2. Arytæno-Epiglottideus,

Arises, by a number of small fibres, from the lateral and upper part of the arytenoid cartilage; and, running along the outer side of the external rima, is

Inserted into the epiglottis along with the former.

Use. To pull that side of the epiglottis towards the external rima; or, when both act, to pull it close upon the glottis. It is counteracted by the elasticity of the epiglottis.

CHAP. XV.

Muscles *situated on the anterior part of the* Abdomen.

THEY consist of three broad layers on each side of the belly; always a long one, and generally also a short one, on each side of the linea alba.

The three layers are,

1. Obliquus Descendens Externus.

Arises, by eight heads, from the lower edges of an equal number of inferior ribs, at a little distance from their cartilages: It always intermixes, in a serrated manner, with portions of the serratus major anticus; and generally coheres to the pectoralis major, intercostals, and latissimus dorsi: Which last covers the edge of a portion of it extend-

ed

ed from the laſt rib to the ſpine of the os ilium. From theſe origins the fibres run down obliquely forwards, and terminate in a thin broad tendon, whoſe fibres are continued in the ſame direction.

Inſerted into the whole length of the *linea alba* *; becomes thicker towards the lower part of the abdomen, and is perforated in the middle by the umbilicus †. On the outſide of the rectus muſcle, the tendon of the external oblique appears whiter than elſewhere, by its being there connected with the tendons of the internal oblique and tranſverſe muſcles; ſo that this part has been called *linea ſemilunaris*, from its curved ſhape. The under part of the tendon divides into two columns, which leaves an oval ſpace between them, named the *ring* ‡ of the external oblique muſcle, for the paſſage of the ſpermatic cord in the male, or round ligament of the womb: The anterior ſuperior column paſſes over the cartilage between the oſſa pubis, and is fixed to the oppoſite os pubis; the other is fixed to the os pubis of the ſame ſide. It is alſo inſerted, tendinous and fleſhy, into the middle of the ſpine of the ilium.

From that part, which is named its *anterior ſuperior ſpinous proceſs*, it is ſtretched tendinous to the os pubis, and is

* The *linea alba* is formed by the tendinous fibres of the two oblique and tranſverſe muſcles, interlaced with thoſe of the oppoſite ſide, the whole way from the cartilago enſiformis to the os pubis; ſo that ſome anatomiſts think they ſhould be called three digaſtric muſcles, with a broad middle tendon and two fleſhy bellies.

† The *umbilicus* was originally the paſſage for the veſſels that connected the foetus to the ſecundines; and is really a hole through the tegumens and tendons, filled up only by a cellular ſubſtance, and covered within by the peritoneum.

‡ The *ring* of the external oblique muſcle is made ſomewhat circular by a thin tendinous or rough cellular ſubſtance, which helps to fill it up, and though a few muſcular fibres of the internal are ſeparated, yet the ſtricture in herniae only happens in the tendon of the external.

named *Poupart's* or *Fallopius's* ligament *. From this ligament it ſends a tendinous layer, which is loſt in the membranous faſcia of the thigh.

Uſe. Supports and compreſſes the peritonaeum and abdomen; aſſiſts the evacuations of faeces and urine, and likewiſe in the excluſion of the foetus; thruſts the diaphragm upwards, and draws down the ribs in expiration; bends the body obliquely when the ribs are fixed, and raiſes the pelvis obliquely.

Obliquus externus abdominis, Albinus.

Obliquus deſcendens, Douglas.

2. Obliquus Ascendens Internus,

Ariſes from the ſpine of the ilium, the whole length between the poſterior and ſuperior anterior ſpinous proceſs; from the os ſacrum and the three undermoſt lumbar vertebrae, by a tendon common to it and to the ſerratus poſticus inferior muſcle; from Poupart's ligament, at the middle of which it ſends off the beginning of the cremaſter muſcle; and the ſpermatic cord in the male, or round ligament of the womb in the female, paſſes under its thin edge, except a few detached fibres.

Inſerted into the cartilago enſiformis, into the cartilages of the ſeventh, and thoſe of all the falſe ribs; but, at the upper part, it is extremely thin, reſembling a cellular membrane,

* *Poupart's* or *Fallopius's ligament* is the inferior part of the tendon of the externus obliquus, extending from the anterior ſuperior ſpinous proceſs of the ilium to the os pubis, where it is thickeſt in order to ſtrengthen the inferior part of the abdomen: Here it is not inſerted into any bone, but paſſes over the blood veſſels of the inferior extremity; and in women, from the greater ſize of the pelvis, it is longer and looſer, by which they are more ſubject to crural herniae; but, by the ſize of the ſpermatic cord, men are more liable to the inguinal.

brane, and only becomes fleſhy at the cartilage of the tenth rib. Here its tendon divides into two layers *; the anterior layer, with a great portion of the inferior part of the poſterior layer, joins the tendon of the external oblique, and runs over the rectus to be inſerted into the whole length of the linea alba. The poſterior layer joins the tendon of the tranſverſalis muſcle as low as half-way between the umbilicus and os pubis; but, below this place, only a few fibres of the poſterior layer are ſeen, and the reſt of it paſſes before the rectus muſcle, and is inſerted into the linea alba; ſo that the whole tendon of the external oblique muſcle, with the anterior layer of the internal oblique, paſſes before the rectus muſcle; and the whole poſterior layer of the internal oblique, together with the whole tendon of the tranſverſalis muſcle, excepting at the inferior part, paſs behind the rectus, and are inſerted into the linea alba. At its undermoſt part it is inſerted into the fore-part of the os pubis.

Uſe. To aſſiſt the former; but it bends the trunk in the reverſe direction.

Obliquus internus abdominis, Albinus and Winſlow.

Obliquus aſcendens, Douglas.

3. TRANSVERSALIS,

Ariſes tendinous, but ſoon becoming fleſhy, from the inner or back part of the cartilages of the ſeven lower ribs, where

* To obtain a proper view of the two layers of the tendon of the internal oblique muſcles, both the oblique muſcles ſhould be raiſed as far forwards as their joining near the linea ſemilunaris: then the tendon before the rectus muſt be cut parallel to the linea alba, and turned outwards as far as the outer edge of the rectus; by which the whole of the rectus is brought into view, and the tendons are preſerved. But Douglas directs to cut the poſterior layer of the internal oblique, where it joins with the tranſverſalis; by this method the rectus is laid bare; but the ſtructure of the tendinous ſheath, which incloſes it, is deſtroyed.

where ſome of its fibres are continued with thoſe of the diaphragm and the intercoſtal muſcles; by a broad thin tendon, connected to the tranſverſe proceſſes of the laſt vertebra of the back and the four ſuperior vertebra of the loins: fleſhy, from the whole ſpine of the os ilium internally, and from the tendon of the external oblique muſcle, where it intermixes with ſome fibres of the internal oblique.

Inſerted into the cartilago enſiformis, and into the whole length of the linea alba, excepting its lowermoſt part.

Uſe. To ſupport and compreſs the abdominal bowels; and it is ſo particularly well adapted for the latter purpoſe, that it might be called the *proper conſtrictor* of the abdomen.

Tranſverſus abdominis, Albinus.

The long muſcle in the middle is named,

RECTUS ABDOMINIS.

Ariſes, by two heads, from the ligament of the cartilage which joins the two oſſa pubis to each other; runs upwards the whole length of, and parallel to, the linea alba, growing broader and thinner as it aſcends.

Inſerted into the cartilages of the three inferior true ribs, and often intermixed with ſome fibres of the pectoral muſcle.

It is generally divided by three tendinous interſections; the firſt is at the umbilicus, the ſecond where it runs over the cartilage of the ſeventh rib, and the third in the middle between theſe; and there is commonly a half interſection below the umbilicus: Theſe interſections ſeldom penetrate through the whole thickneſs of the muſcle; they adhere firmly to the anterior part of the ſheath, but very ſlightly to the poſterior layer.

Uſe. To compreſs the fore-part, but more particularly the

the lower part of the belly; to bend the trunk forwards, or to raiſe the pelvis. By its tendinous interſections, it is enabled to contract at any of the intermediate ſpaces; and, by its connection with the tendons of the other muſcles, it is prevented from changing place, and from riſing into a prominent form when in action.

The ſhort muſcle in the middle is named

PYRAMIDALIS.

Ariſes along with the rectus; and, running upwards, within the ſame ſheath, is

Inſerted, by an acute termination, near half-way between the os pubis and umbilicus, into the linea alba and inner edge of the rectus muſcle.

As it is frequently wanting in both ſides, without any inconveniency, its

Uſe ſeems to be, to aſſiſt the inferior part of the rectus.

CHAP. XVI.

MUSCLES *about the Male Organs of* GENERATION.

THE *teſticles* are ſaid to have a thin muſcle common to both, and one proper to each.

The ſuppoſed common muſcle is called the

DARTOS.

This appears to be no more than a condenſation of the cellular membrane lining the ſcrotum; yet the ſkin here

is

is capable of being corrugated and relaxed in a greater degree than in other places.

The muſcle proper to each teſticle is the

Cremaster.

Ariſes from the internal oblique, where a few fibres of that muſcle intermix with the tranſverſalis, near the junction of the os ilium and pubis, over which part it paſſes, after having pierced the ring of the externus obliquus; and then it deſcends upon the ſpermatic cord.

Inſerted into the tunica vaginalis of the teſticle, upon which it ſpreads, and is inſenſibly loſt.

Uſe. To ſuſpend and draw up the teſticle, and to compreſs it in the act of coition.

The penis has three pair of muſcles,

1. Erector Penis,

Ariſes, tendinous and fleſhy, from the tuberoſity of the os iſchium, and runs upwards, embracing the whole crus of the penis.

Inſerted into the ſtrong tendinous membrane that covers the corpora cavernoſa penis, near as far up as the union of theſe bodies.

Uſe. To compreſs the crura penis, by which the blood is puſhed from it into the fore-part of the corpora cavernoſa; and the penis is by that means more completely diſtended. The erectores ſeem likewiſe to keep the penis in its proper direction.

Iſchio-cavernoſus, Winſlow.

2. Accelerator Urinae, *ſeu* Ejaculator Seminis,

Ariſes, fleſhy, from the ſphincter ani and membranous part of the urethra; and tendinous from the crus, nearly as far

far forwards as the beginning of the corpus cavernosum penis; the inferior fibres run more transversely, and the superior descend in an oblique direction.

Inserted into a line in the middle of the bulb, where it joins with its fellow, by which the bulb is completely incloſed.

Use. To drive the urine or semen forwards; and, by graſping the bulb of the urethra, to push the blood towards its corpus cavernosum and the glans, by which they are distended.

Bulbo-cavernosus, Winslow.

3. Transversus Perinei.

Arises from the tough fatty membrane that covers the tuberosity of the os ischium; from thence it runs transversely inwards, and is

Inserted into the accelerator urinae, and into that part of the sphincter ani which covers the bulb.

Use. To dilate the bulb, and draw the perineum and verge of the anus a little outwards and backwards.

Transversalis urethrae, Winslow.

Transversus perinei, Albinus.

Levator parvus, seu *externus*, Douglas.

There is often a fourth muscle, named

Transversus Perinei Alter.

Arises behind the former, runs more obliquely forwards, and is

Inserted into that part of the accelerator urinae which covers the anterior part of the bulb of the urethra.

Use. To assist the former.

Inferior prostate, Winslow.

Transversus perinei alter, Albinus.

CHAP.

CHAP. XVII.

Muscles *of the* Anus.

THE *anus* has a ſingle muſcle, and one pair.

The ſingle muſcle is

Sphincter Ani.

Ariſes from the ſkin and fat that ſurround the verge of the anus on both ſides, near as far out as the tuber of the os iſchium; the fibres are gradually collected into an oval form, and ſurround the extremity of the rectum.

Inſerted, before, by a narrow point, into the perineum, acceleratores urinae, and tranſverſi perinei; behind, by an acute termination, into the extremity of the os coccygis.

Uſe. Shuts the paſſage through the anus into the rectum: pulls down the bulb of the urethra, by which it aſſiſts in ejecting the urine and ſemen.

Sphincter externus, Albinus and Douglas.

Sphincter cutaneus, Winſlow.

N. B. The ſphincter internus of Albinus and Douglas is only that part of the circular fibres of the muſcular coat of the rectum, which ſurrounds its extremity.

Levator Ani,

Ariſes from the os pubis within the pelvis, as far up as the upper edge of the foramen thyroideum, and joining of the

the os pubis with the os ischium; from the thin tendinous membrane that covers the obturator internus and coccygeus muscles; from the spinous process of the os ischium; and its fibres run down like rays from a circumference to a centre.

Inserted into the sphincter ani, acceleratores urinae, and anterior part of the two last bones of the coccygis; surrounds the extremity of the rectum, neck of the bladder, prostate gland, and part of the vesiculae seminales; so that its fibres behind and below the os coccygis joining it with its fellow, they together very much resemble the shape of a funnel.

Use. To draw the rectum upwards after the evacuation of the faeces, and to assist in shutting it; to sustain the contents of the pelvis, and to help in ejecting the semen, urine, and contents of the rectum; and, perhaps, by pressing upon the veins, to contribute greatly to the erection of the penis.

CHAP. XVIII.

Muscles *of the Female Organs of* Generation.

THE *clitoris* has one pair,

Erector Clitoridis,

Arises from the crus of the os ischium internally, and in its ascent covers the crus of the clitoris as far up as the os pubis.

Inſerted into the upper part of the crus and body of the clitoris.

Uſe. Draws the clitoris downwards and backwards; and may ſerve to make the body of the clitoris more tenſe, by ſqueezing the blood into it from its crus.

Firſt muſcle of the clitoris, Douglas.

The vagina has one pair,

Sphincter Vaginæ,

Ariſes from the ſphincter ani, and from the poſterior ſide of the vagina, near the perineum: From thence it runs up the ſide of the vagina, near its external orifice, oppoſite to the nymphae, and covers the corpus cavernoſum vaginae.

Inſerted into the crus and body, or union of the crura clitoridis.

Uſe. Contracts the mouth of the vagina, and compreſſes its corpus cavernoſum.

Conſtrictor cunni, Albinus.

Second muſcle of the clitoris, Douglas.

The perineum has one pair,

Transversus Perinei,

Ariſes, as in the male, from the fatty cellular membrane which covers the tuberoſity of the os iſchium.

Inſerted into the upper part of the ſphincter ani, and into a white hardiſh tough ſubſtance in the perineum, between the lower part of the pudendum and anus.

Uſe. To ſuſtain and keep the perineum in its proper place.

The anus, as in the male, has a ſingle muſcle, and one pair.

Sphincter

Sphincter Ani,

Arifes, as in the male, from the ſkin and fat ſurrounding the extremity of the rectum.

Inſerted, above, into the white tough ſubſtance of the perineum; and below, into the point of the os coccygis.

Uſe. To ſhut the paſſage into the rectum; and, by pulling down the perineum, to aſſiſt in contracting the mouth of the vagina.

Levator Ani,

Arifes, as in the male, within the pelvis, and deſcends along the inferior part of the vagina and rectum.

Inſerted into the perineum, ſphincter ani, extremity of the vagina, and rectum.

Uſe. To raiſe the extremity of the rectum upwards, to contract the inferior part of the rectum, and to aſſiſt in contracting and ſupporting the vagina; and, perhaps, by preſſing on the veins, to contribute to the diſtention of the cells of the clitoris and corpus cavernoſum of the vagina.

CHAP. XIX.

Muscles *ſituated within the* Pelvis.

OF theſe there are two pair.

1. Obturator Internus,

Arifes from more than one half of the internal circumference of the foramen thyroideum, formed by the os pubis and iſchium: Its inſide is covered by a portion of the levator

tor ani; and appears to be divided into a number of fasciculi, which unite and form a roundish tendon, that passes out of the pelvis, between the posterior sacro ischiatic ligament and tuberosity of the os ischium; where it passes over the capsular ligament of the thigh-bone, it is inclosed as in a sheath, by the gemini muscles.

Inserted, by a round tendon, into the large pit at the root of the trochanter major.

Use. To roll the os femoris obliquely outwards.

Marsupialis, seu *Obturator internus*, Douglas,

N. B. The insertion of this muscle should not be prosecuted, until the muscles of the thigh, to which it belongs, are dissected. *Vid.* Chap. xxix.

2. Coccygeus,

Arises, tendinous and fleshy, from the spinous process of the os ischium, and covers the inside of the posterior sacro-ischiatic ligament; from this narrow beginning, it gradually increases, to form a thin fleshy belly, interspersed with tendinous fibres.

Inserted into the extremity of the os sacrum, and near the whole length of the os coccygis laterally.

Use. To support and move the os coccygis forwards, and to tie it more firmly to the sacrum.

CHAP. XX.

Muscles *situated within the Cavity of the* Abdomen.

These consist of a single muscle, and four pair.

Diaphragma.

This broad thin muscle, which makes a complete septum between the thorax and abdomen, is concave below and convex

convex above; the middle of it on each ſide reaching as high within the thorax of the ſkeleton as the fourth rib; it is commonly divided into two portions.

1. The ſuperior or Greater Muſcle of the DIAPHRAGM,

Ariſes, by diſtinct fleſhy fibres, from the cartilago enſiformis, from the cartilages of the ſeventh, and of all the inferior ribs on both ſides. The fibres from the cartilago enſiformis, and from the ſeventh and eighth ribs, run obliquely upwards and backwards; from the ninth and tenth, tranſverſely inwards and upwards; and from the eleventh and twelfth, obliquely upwards. From theſe different origins the fibres run, like radii from the circumference to the centre of a circle; and are

Inſerted into a cordiform tendon, of a conſiderable breadth, which is ſituated in the middle of the diaphragm, and in which, therefore, the fibres from oppoſite ſides are interlaced. Towards the right ſide the tendon is perforated, by a triangular hole for the paſſage of the vena cava inferior; and to the upper convex part of it the pericardium and mediaſtinum are connected.

2. The inferior, Leſſer Muſcle, or Appendix of the DIAPHRAGM,

Ariſes from the ſecond, third, and fourth lumbar vertebrae, by eight heads; of which two in the middle, commonly called its *crura*, are the longeſt, and begin tendinous. Between the crura, the aorta and thoracic duct paſs; and, on the outſide of theſe, the great ſympathetic nerves and branches of the vena azygos perforate the ſhorter heads. The muſcular fibres run obliquely upwards and forwards, and form in the middle two fleſhy columns, which decuſſate and leave an oval ſpace between them for the paſſage of the oeſophagus and eighth pair of nerves.

Inſerted,

Inſerted, by ſtrong fleſhy fibres, into the poſterior part of the middle tendon.

Uſe. The diaphragm is the principal agent in reſpiration, particularly in inſpiration: for when it is in action, the fibres, from their different attachments, endeavour to bring themſelves into a plane towards the middle tendon, by which the cavity of the thorax is enlarged, particularly at the ſides, where the lungs are chiefly ſituated; and as the lungs muſt always be contiguous to the inſide of the thorax and upper ſide of the diaphragm, the air ruſhes into them, in order to fill up the increaſed ſpace. This muſcle is aſſiſted by the two rows of intercoſtals, which elevate the ribs, and the cavity of the thorax is more enlarged. In time of violent exerciſe, or whatever cauſe drives the blood with unuſual celerity towards the lungs, the pectoral muſcles, the ſerrati antici majores, the ſerrati poſtici ſuperiores, and ſcaleni muſcles, are brought into action. And in laborious inſpiration, the muſcles which ariſe from the upper part of the thorax, when the parts into which they are inſerted, are fixed, likewiſe aſſiſt. In exſpiration, the diaphragm is relaxed and puſhed up by the preſſure of the abdominal muſcles upon the viſcera of the abdomen; and at the ſame time that they preſs it upwards, they alſo, together with the ſterno-coſtales and ſerrati poſtici inferiores, pull down the ribs, and are aſſiſted in a powerful manner by the elaſticity of the cartilages that join the ribs to the ſternum; by which the cavity of the thorax is diminiſhed, and the air ſuddenly puſhed out of the lungs: and, in laborious exſpiration, the quadrati lumborum, ſacro-lumbales, and longiſſimi dorſi, concur in pulling down the ribs.

The four pair are,

1. Qua-

1. QUADRATUS LUMBORUM,

Ariſes, ſomewhat broad, tendinous, and fleſhy from the poſterior part of the ſpine of the os ilium.

Inſerted into the tranſverſe proceſſes of all the vertebrae of the loins, into the laſt rib near the ſpine, and by a ſmall tendon into the ſide of the laſt vertebra of the back.

Uſe. To move the loins to one ſide, pull down the laſt rib, and, when both act, to bend the loins forwards.

Quadratus, ſeu *Lumbaris externus*, Winſlow.

2. PSOAS PARVUS,

Ariſes, fleſhy, from the ſides of the two upper vertebrae of the loins, and ſends off a ſmall long tendon, which ends thin and flat, and is

Inſerted into the brim of the pelvis, at the junction of the os ilium and pubis.

Uſe. To aſſiſt the pſoas magnus in bending the loins forwards; and, in certain poſitions, to aſſiſt in raiſing the pelvis.

N. B. This muſcle is very often wanting.

3. PSOAS MAGNUS,

Ariſes, fleſhy, from the ſide of the body, and tranſverſe proceſs of the laſt vertebra of the back; and, in the ſame manner, from all thoſe of the loins, by as many diſtinct ſlips.

Inſerted, tendinous, into the trochanter minor of the os femoris; and fleſhy into that bone, a little below the ſame trochanter.

Uſe. To bend the thigh forwards; or, when the inferior extremity is fixed, to aſſiſt in bending the body.

Pſoas, ſeu *Lumbaris internus*, Winſlow.

4. ILIACUS

4. Iliacus Internus,

Arifes, fleſhy, from the tranſverſe proceſs of the laſt vertebra of the loins, from all the inner lip of the ſpine of the os ilium, from the edge of that bone between its anterior ſuperior ſpinous proceſs and the acetabulum, and from moſt of the hollow part of the ilium. It joins with the pſoas magnus, where it begins to become tendinous; and is

Inſerted along with it.

Uſe. To aſſiſt the pſoas in bending the thigh, and to bring it directly forwards.

N. B. The inſertion of the two laſt muſcles ſhould not be proſecuted till the muſcles of the thigh are diſſected.

CHAP. XXI.

Muscles *ſituated on the Anterior Part of the* Thorax.

THESE may be divided into two layers. The firſt layer conſiſts of one muſcle, named

Pectoralis Major,

Arifes, from the cartilaginous extremities of the fifth and ſixth ribs, where it always intermixes with the external oblique muſcle of the abdomen; from almoſt the whole length of the ſternum, and from near half of the anterior part of the clavicle: the fibres run towards the axilla in a folding manner.

Inſerted, by two broad tendons, which croſs each other at

at the upper and inner part of the os humeri, above the inſertion of the deltoid muſcle, and outer ſide of the groove for lodging the tendon of the long head of the biceps.

Uſe. To move the arm forwards, and obliquely upwards, towards the ſternum.

Pectoralis, Albinus.

The ſecond layer conſiſts of three muſcles:

1. Sueclavius,

Ariſes tendinous from the cartilage that joins the firſt rib to the ſternum.

Inſerted, after becoming fleſhy, into the inferior part of the clavicle, which it occupies from within an inch or ſo of the ſternum, as far outwards as to its connection, by ligament, with the coracoid proceſs of the ſcapula.

Uſe. To pull the clavicle downwards and forwards.

2. Pectoralis Minor,

Ariſes, tendinous and fleſhy, from the upper edge of the third, fourth, and fifth ribs, near where they join with their cartilages.

Inſerted, tendinous, into the coracoid proceſs of the ſcapula; but ſoon grows fleſhy and broad.

Uſe. To bring the ſcapula forwards and downwards, or to raiſe the ribs upwards.

Serratus anticus, Albinus.

Serratus minor anticus, Douglas.

3. Serratus Magnus,

Ariſes from the nine ſuperior ribs, by an equal number of fleſhy digitations, reſembling the teeth of a ſaw.

Inſerted, fleſhy, into the whole baſe of the ſcapula internally, between the inſertion of the rhomboid and the origin

of the ſubſcapularis muſcles, being folded about the two angles of the ſcapula.

Uſe. To move the ſcapula forwards; and, when the ſcapula is forcibly raiſed, to draw upwards the ribs.

Serratus major anticus, Douglas.

CHAP. XXII.

MUSCLES *ſituated between the Ribs, and within the* THORAX.

BETWEEN the ribs, on each ſide, there are eleven double rows of muſcles, which are therefore named *intercoſtals*. Theſe decuſſate each other like the ſtrokes of the letter X.

1. INTERCOSTALES EXTERNI,

Ariſes from the inferior acute edge of each ſuperior rib, and run obliquely forwards, the whole length from the ſpine to near the joining of the ribs with their cartilages; from which, to the ſternum, there is only a thin membrane covering the internal intercoſtals.

Inſerted into the upper obtuſe edge of each inferior rib, as far back as the ſpine, into which the poſterior portion is fixed.

2. INTERCOSTALES INTERNI,

Ariſes in the ſame manner as the external: But they begin at the ſternum, and run obliquely backwards, as far as the angle of the rib; and from that to the ſpine they are wanting.

Inſerted in the ſame manner as the external.

Uſe.

Uſe. By means of theſe muſcles, the ribs are equally raiſed upwards during inſpiration. Their fibres being oblique, give them a greater power of bringing the ribs near each other, than could be performed by ſtraight ones. But, by the obliquity of the fibres, they are almoſt brought contiguous: And as the fixed points of the ribs are before and behind, if the external had been continued forwards to the ſternum, and the internal backwards to the ſpine, it would have hindered their motion, which is greateſt in the middle, though the obliquity of the ribs renders it leſs perceptible; and, inſtead of raiſing the fibres fixed to the ſternum and ſpine, would have depreſſed the ribs.

N. B. The portions of the external intercoſtals which ariſe from the tranſverſe proceſſes of the vertebrae where the ribs are fixed to them, and other portions that paſs over one rib and terminate in the next below it, Albinus calls *Levatores coſtarum longiores et breviores.*

The portions of the internal that paſs over one rib, and are inſerted into the next below it, are by Douglas called, *Coſtarum depreſſores proprii Cowperii.*

Theſe portions of both rows aſſiſt in raiſing the ribs in the ſame manner as the reſt of the intercoſtals.

Supra coſtales, and *Infra coſtales*, Winſlow.

The muſcles within the thorax are one pair, viz.

TRIANGULARIS, *or* STERNOCOSTALIS,

Ariſes, fleſhy, and a little tendinous, from all the length of the cartilago enſiformis laterally, and from the edge of the lower half of the middle bone of the ſternum, from whence its fibres aſcend obliquely upwards and outwards.

Inſerted, generally by three triangular terminations, into the lower edge of the cartilages of the third, fourth, and fifth ribs, near where theſe join with the ribs.

Uſe.

Uſe. To depreſs theſe cartilages, and the extremities of the ribs; and conſequently to aſſiſt in contracting the cavity of the thorax.

This muſcle often varies; and is ſometimes inſerted into the cartilage of the ſecond rib, ſometimes into the cartilage of the ſixth rib.

CHAP. XXIII.

Muscles *ſituated on the Anterior Part of the* Neck *cloſe to the* Vertebræ.

THESE conſiſt of one layer formed by four muſcles.

1. Longus Colli,

Ariſes, tendinous and fleſhy, from the bodies of the three vertebrae of the back laterally; and from the tranſverſe proceſs of the third, fourth, fifth, and ſixth vertebrae of the neck, near their roots.

Inſerted into the fore-part of the bodies of all the vertebrae of the neck, by as many ſmall tendons, which are covered with fleſh.

Uſe. To bend the neck gradually forwards, and to one ſide.

2. Rectus Capitis Internus Major,

Ariſes from the anterior points of the tranſverſe proceſſes of the third, fourth, fifth, and ſixth vertebrae of the neck, by four diſtinct beginnings.

Inſerted into the cuneiform proceſs of the os occipitis, a little before the condyloid proceſs.

Uſe.

Uſe. To bend the head forwards.

Rectus anterior longus, Winſlow.

3. Rectus Capitis Internus Minor,

Ariſes, fleſhy, from the fore-part of the body of the firſt vertebra of the neck, oppoſite to the ſuperior oblique proceſs.

Inſerted near the root of the condyloid proceſs of the os occipitis, under, and a little farther outwards than the former muſcle.

Uſe. To bend the head forwards.

Rectus anterior brevis, Winſlow.

4. Rectus Capitis Lateralis,

Ariſes, fleſhy, from the anterior part of the point of the tranſverſe proceſs of the firſt vertebra of the neck.

Inſerted into the os occipitis, oppoſite to the foramen ſtylo-maſtoideum of the temporal bone.

Uſe. To bend the head a little to one ſide.

Tranſverſalis anticus primus, Winſlow.

CHAP. XXIV.

Muscles *ſituated on the Poſterior Part of the* Trunk.

THESE may be divided into four layers, and a ſingle pair.

The firſt layer conſiſts of two muſcles, which cover almoſt the whole poſterior part of the trunk.

1. Tra-

1. Trapezius *seu* Cucularis,

Ariſes, by a ſtrong round tendon, from the lower part of the protuberance in the middle of the os occipitis behind; and, by a thin membranous tendon, which covers part of the ſplenius and complexus muſcles, from the rough curved line that extends from the protuberance towards the maſtoid proceſs of the temporal bone; runs down along the nape of the neck; where it ſeems to ariſe from its fellow, and covers the ſpinous proceſſes of the ſuperior vertebrae of the neck; but riſes from the ſpinous proceſſes of the two inferior, and from the ſpinous proceſſes of all the vertebrae of the back; adhering, tendinous, to its fellow, the whole length of its origin.

Inſerted, fleſhy, into the poſterior half of the clavicle; tendinous and fleſhy, into the acromion, and into almoſt all the ſpine of the ſcapula.

Uſe. Moves the ſcapula according to the three different directions of its fibres; for the upper deſcending fibres draw it obliquely upwards, the middle tranſverſe ſtraight fibres draw it directly backwards, and the inferior aſcending fibres draw it obliquely downwards and backwards.

N. B. Where it is inſeparably united to its fellow in the nape of the neck, it is named *Ligamentum Nuchae* or *Colli.*

2. Latissimus Dorsi,

Ariſes, by a broad thin tendon, from the poſterior part of the ſpine of the os ilium, from all the ſpinous proceſſes of the os ſacrum and vertebrae of the loins, and from the ſeven inferior ones of the vertebrae of the back; alſo, tendinous and fleſhy, from the extremities of the three or four inferior ribs, a little beyond their cartilages, by as many diſtinct ſlips. The inferior fibres aſcend obliquely, and the ſuperior run tranſverſely, over the inferior angle of

of the ſcapula, towards the axilla, where they are all collected, twiſted, and folded.

Inſerted, by a ſtrong thin tendon, into the inner edge of the groove for lodging the tendon of the long head of the biceps.

Uſe. To pull the arm backwards and downwards, and to roll the os humeri.

N. B. The inſertion of this muſcle ſhould not be proſecuted till the muſcles of the os humeri, to which it belongs, are diſſected.

The ſecond layer conſiſts of three pair, two on the back, and one on the neck.

On the back,

1. Serratus Posticus Inferior,

Ariſes, by a broad thin tendon, in common with that of the latiſſimus dorſi, from the ſpinal proceſſes of the two inferior vertebrae of the back, and from the three ſuperior vertebrae of the loins.

Inſerted into the lower edges of the four inferior ribs, at a little diſtance from their cartilages, by as many diſtinct fleſhy lips.

Uſe. To depreſs the ribs into which it is inſerted.

2. Rhomboideus.

This muſcle is divided into two portions.

1. *Rhomboideus major*, *ariſes*, tendinous, from the ſpinous proceſſes of the five ſuperior vertebrae of the back.

Inſerted into all the baſis of the ſcapula below its ſpine.

Uſe. To draw the ſcapula obliquely upwards, and directly inwards.

2. *Rhomboideus minor*, *ariſes*, tendinous, from the ſpinous

nous proceſſes of the three inferior vertebrae of the neck, and from the ligamentum nuchae.

Inſerted into the baſe of the ſcapula, oppoſite to its ſpine.

Uſe. To aſſiſt the former.

On the neck,

3. SPLENIUS,

Ariſes, tendinous, from the four ſuperior ſpinous proceſſes of the vertebrae of the back; tendinous and fleſhy, from the five inferior of the neck, and adheres firmly to the ligamentum nuchae. At the third vertebra of the neck, the ſplenii recede from each other, ſo that part of the complexus muſcle is ſeen.

Inſerted, by as many tendons, into the five ſuperior tranſverſe proceſſes of the vertebrae of the neck; and tendinous and fleſhy, into the poſterior part of the maſtoid proceſs, and into the os occipitis, where it joins with the root of that proceſs.

Uſe. To bring the head and upper vertebrae of the neck backwards laterally; and, when both act, to pull the head directly backwards.

N. B. Albinus divides this muſcle into two; viz. That portion which ariſes from the five inferior ſpinous proceſſes of the neck, and is inſerted into the maſtoid proceſs and os occipitis, he calls *Splenius Capitis*: and that portion which ariſes from the third, and fourth of the back, and is inſerted into the five ſuperior tranſverſe proceſſes of the neck, is called by him *Splenius Colli*.

The ſingle pair,

SERRATUS SUPERIOR POSTICUS,

Ariſes, by a broad thin tendon, from the ſpinous proceſſes

ceſſes of the three laſt vertebrae of the neck, and the two uppermoſt of the back.

Inſerted into the ſecond, third, fourth, and fifth ribs, by as many fleſhy ſlips.

Uſe. To elevate the ribs, and dilate the thorax.

The third layer conſiſts of three pair on the back, and three on the neck.

Thoſe on the back are,

1. Spinalis Dorsi,

Ariſes from the ſpinous proceſſes of the two uppermoſt vertebrae of the loins, and the three inferior of the back by as many tendons.

Inſerted into the ſpinous proceſſes of the nine uppermoſt vertebrae of the back, except the firſt, by as many tendons.

Uſe. To erect and fix the vertebrae, and to aſſiſt in raiſing the ſpine.

2. Longissimus Dorsi,

Ariſes, tendinous without, and fleſhy within, from the ſide, and all the ſpinous proceſſes of the os ſacrum: From the poſterior ſpine of the os ilium: from all the ſpinous proceſſes; and from the roots of the tranſverſe proceſſes of the vertebrae of the loins.

Inſerted into all the tranſverſe proceſſes of the vertebrae of the back, chiefly by ſmall double tendons; alſo by a tendinous and fleſhy ſlip, into the lower edge of all the ribs, except the two inferior, at a little diſtance from their tubercles.

Uſe. To extend the vertebrae, and to raiſe and keep the trunk of the body erect.

N. B. From the upper part of this muſcle there runs up

a round fleshy portion which joins with the cervicalis descendens.

3. Sacro-Lumbalis,

Arises in common with the longissimus dorsi.

Inserted into all the ribs, where they begin to be curved forwards, by as many long and thin tendons; and,

From the upper part of the six or eight lower ribs, arise as many bundles of thin fleshy fibres, which soon terminate in the inner side of this muscle, and are named *Musculi ad Sacro-lumbalem Accessorii.*

Use. To pull the ribs down, and assist to erect the trunk of the body.

N. B. There is a fleshy slip which runs from the upper part of this muscle into the fourth, fifth, and sixth transverse processes of the vertebrae of the neck, by three distinct tendons: It is named *Cervicalis Descendens;* and *its* use is to turn the neck obliquely backwards, and to one side.

On the neck are,

1. Complexus,

Arises from the transverse processes of the seven superior vertebrae of the back, and four inferior of the neck, by as many distinct tendinous origins; in its ascent it receives a fleshy slip from the spinous process of the first vertebra of the back: From these different origins it runs upwards, and is every where intermixed with tendinous fibres.

Inserted, tendinous and fleshy, into the inferior edge of the protuberance in the middle of the os occipitis, and into a part of the curved line that runs forwards from that protuberance.

Use. To draw the head backwards, and to one side; and, when both act, to draw the head directly backwards.

N. B.

N. B. The long portion of this muſcle that is ſituated next the ſpinous proceſſes, lies more looſe, and has a roundiſh tendon in the middle of it; for which reaſon Albinus calls it *Bivrenter cervicis.*

2. Trachelo-Mastoideus,

Ariſes from the tranſverſe proceſſes of the three uppermoſt vertebrae of the back, and from the five lowermoſt of the neck, where it is connected to the tranſverſalis cervicis, by as many thin tendons, which unite into a belly, and run up under the ſplenius.

Inſerted into the middle of the poſterior ſide of the maſtoid proceſs, by a thin tendon.

Uſe. To aſſiſt the complexus; but it pulls the head more to a ſide.

Complexus minor, ſeu *Maſtoideus lateralis*, Winſlow.

Trachelo maſtoideus, ſeu *Capitis par tertium Fallopii*, Douglas.

3. Levator Scapulae,

Ariſes, tendinous and fleſhy, from the tranſverſe proceſſes of the five ſuperior vertebrae of the neck, by as many diſtinct ſlips, which ſoon unite to form a muſcle that runs downwards and outwards.

Inſerted, fleſhy, into the ſuperior angle of the ſcapula.

Uſe. To pull the ſcapula upwards, and a little forwards.

Angularis, vel *Levator proprius*, Winſlow.

Elevator ſeu *Muſculus patientiae*, Douglas.

The fourth layer conſiſts of two pair on the back, two on the poſterior part of the neck, four ſmall pair ſituated immediately below the poſterior part of the occiput, and three on the ſide of the neck.

On the back are,

1. Semi-spinalis Dorsi,

Ariſes, from the tranſverſe proceſſes of the ſeventh, eighth, ninth, and tenth vertebrae of the back, by as many diſtinct tendons, which ſoon grow fleſhy, and then become tendinous again; and are

Inſerted, into the ſpinous proceſſes of all the vertebrae of the back above the eighth, and into the two lowermoſt of the neck, by as many tendons.

Uſe. To extend the ſpine obliquely backwards.

Semi-ſpinalis externus, ſeu *Tranſverſo-ſpinalis dorſi*, Winſlow.

2. Multifidus Spinae,

Ariſes from the ſide and ſpinous proceſſes of the os ſacrum, and from the poſterior part of the os ilium, where it joins with the ſacrum; from all the oblique and tranſverſe proceſſes of the vertebrae of the loins; from all the tranſverſe proceſſes of the vertebrae of the back, and from thoſe of the neck, except the three firſt, by as many diſtinct tendons, which ſoon grow fleſhy, run in an oblique direction; and are

Inſerted, by diſtinct tendons, into all the ſpinous proceſſes of the vertebrae of the loins, of the back, and of the neck, except the firſt.

Uſe. When the different portions of this muſcle act on one ſide, they extend the back obliquely, or move it laterally; but if they act together on both ſides, they extend the vertebrae backwards.

Tranſverſo-ſpinalis lumborum, veterib. Sacer.

Semi ſpinalis internus, ſive *Tranſverſo-ſpinalis dorſi.*

Semi-

Semi-ſpinalis, ſive *Tranſverſo-ſpinalis colli*, *Pars interna*, Winſlow.

Tranſverſalis lumborum, *vulgo Sacer*.

Tranſverſalis dorſi.

Tranſverſalis colli, Douglas.

On the poſterior part of the neck are,

1. SEMI-SPINALIS COLLI,

Ariſes, from the tranſverſe proceſſes of the uppermoſt ſix vertebrae of the back, by as many diſtinct tendons, aſcending obliquely under the complexus.

Inſerted into the ſpinous proceſſes of all the vertebrae of the neck, except the firſt and the laſt.

Uſe. To extend the neck obliquely backwards.

Semi-ſpinalis, ſive *Tranſverſo-ſpinalis colli*, Winſlow.

Spinalis cervicis, Albinus.

Spinalis, Douglas.

2. TRANSVERSALIS COLLI,

Ariſes from the tranſverſe proceſſes of the five uppermoſt vertebrae of the back, by as many tendinous and fleſhy origins; runs between the trachelo-maſtoideus, and ſplenius colli and cervicalis deſcendens.

Inſerted into the tranſverſe proceſſes of all the cervical vertebrae, except the firſt and the laſt.

Uſe. To turn the neck obliquely backward, and a little to one ſide.

Below the poſterior part of the occiput are,

1. RECTUS CAPITIS POSTICUS MAJOR,

Ariſes, fleſhy, from the external part of the ſpinous proceſs of the ſecond vertebra of the neck; and grows broader

broader in its aſcent, which is not ſtraight, but obliquely outwards.

Inſerted, tendinous and fleſhy, into the os occipitis, near the rectus capitis lateralis, and the inſertion of the obliquus capitis ſuperior.

Uſe. To pull the head backwards, and to aſſiſt a little in its rotation.

Rectus major, Winſlow and Douglas.

2. Rectus Capitis Posticus Minor,

Ariſes, by a narrow beginning, cloſe to its fellow, from a little protuberance in the middle of the back part of the firſt vertebra of the neck, its outer edge being covered by the rectus major.

Inſerted, ſomewhat broad, into the ſides of a dimple in the os occipitis, near its foramen magnum.

Uſe. To aſſiſt the rectus major in moving the head backwards.

Obliquus minor, Winſlow and Douglas.

3. Obliquus Capitis Superior,

Ariſes from the tranſverſe proceſs of the firſt vertebra of the neck.

Inſerted, tendinous and fleſhy, into the os occipitis behind the back part of the maſtoid proceſs of the temporal bone, and under the inſertion of the complexus muſcle.

Uſe. To draw the head backwards.

Obliquus major, Winſlow.

Obliquus ſuperior, Douglas.

4. Obliquus Capitis Inferior,

Ariſes, fleſhy, from the ſpinous proceſs of the ſecond vertebra of the neck, its whole length; and, forming a thick fleſhy belly, is

Inſerted

Inserted into the tranſverſe proceſs of the firſt vertebra of the neck.

Uſe. To give a rotatory motion to the head.

On the ſide of the neck are,

1. Scalenus Anticus,

Ariſes, from the fourth, fifth, and ſixth tranſverſe proceſſes of the firſt vertebra of the neck, by as many tendons.

Inſerted, tendinous and fleſhy, into the upper ſide of the firſt rib, near its cartilage.

Scalenus prior, Albinus.

Anterior portion of the firſt ſcalenus, Winſlow.

Firſt ſcalenus, Douglas.

2. Scalenus Medius,

Ariſes, from all the tranſverſe proceſſes of the vertebrae of the neck, by as many ſtrong tendons; the nerves to the ſuperior extremity paſs between it and the former.

Inſerted into the upper and outer part of the firſt rib, from its root, to within the diſtance of an inch from its cartilage.

Poſterior portion of the firſt ſcalenus, Winſlow.

Second ſcalenus, Douglas.

3. Scalenus Posticus,

Ariſes from the fifth and ſixth tranſverſe proceſſes of the vertebrae of the neck.

Inſerted into the upper edge of the ſecond rib, not far from the ſpine.

Poſterior portion of the ſecond ſcalenus, Winſlow.

Third ſcalenus, Douglas.

Uſe of the three ſcaleni: To bend the neck to one ſide;

or,

or, when the neck is fixed, to elevate the ribs, and to dilate the thorax.

There are a number of ſmall muſcles ſituated between the ſpinous and tranſverſe proceſſes of contiguous vertebrae; which are accordingly named,

1. Interspinales Colli,

The ſpace between the ſpinous proceſſes of the vertebrae of the neck, moſt of which are bifurcated, is filled up with fleſhy portions; which

Ariſe, double, from the ſpinous proceſs of the inferior vertebrae of the neck; and aſcend to be

Inſerted, in the ſame manner, into the ſpinous proceſs of the ſuperior vertebra. They are five in number.

Uſe. To draw theſe proceſſes nearer to each other.

2. Intertransversales Colli.

They begin from the tranſverſe proceſs of the firſt vertebra of the back, and fill up the ſpaces between the tranſverſe proceſſes of the vertebrae of the neck, which are likewiſe bifurcated; and, conſequently, there are ſix diſtinct double muſcles, which

Ariſe from the inferior tranſverſe proceſs of each vertebra of the neck, and firſt of the back, and are

Inſerted into the ſuperior tranſverſe proceſſes.

Uſe. To draw theſe proceſſes towards each other, and turn the neck a little to one ſide.

3, 4, 5. Interspinales Dorsi et Lumborum, and the Intertransversales Dorsi,

Are rather ſmall tendons than muſcles, ſerving to connect the ſpinal and tranſverſe proceſſes.

6. Inter-

6. Intertransversales Lumborum,

Are four diſtinct ſmall bundles of fleſh, which fill up the ſpaces between the tranſverſe proceſſes of the vertebrae of the loins, and ſerve to draw them towards each other.

CHAP. XXV.

Muscles *of the* Superior Extremities.

THESE may be divided into the muſcles that are ſituated on the ſcapula, on the os humeri, on the cubit or fore-arm, and on the hand.

Muſcles ſituated on the ſcapula.

Theſe are called *muſcles* of the *os humeri*; and are three behind, one along its inferior coſta, two before, and one beneath it.

Behind are,

1. Supraspinatus,

Ariſes, fleſhy, from all that part of the baſe of the ſcapula that is above its ſpine; alſo from the ſpine and ſuperior coſta; paſſes under the acromion, and adheres to the capſular ligament of the os humeri.

Inſerted, tendinous, into that part of the large protuberance on the head of the os humeri that is next the groove for lodging the tendon of the long head of the biceps.

Uſe. To raiſe the arm upwards; and, at the ſame time, to pull the capſular ligament from between the bones, that it may not be pinched.

2. Infraspinatus,

Arifes, fleſhy, from all that part of the baſe of the ſcapula that is between its ſpine and inferior angle; from the ſpine as far as the cervix of the ſcapula. The fibres aſcend and deſcend obliquely towards a tendon in the middle of the muſcle, which runs forwards, and adheres to the capſular ligament.

Inſerted, by a thick and ſhort tendon, into the upper and middle part of the large protuberance on the head of the os humeri.

Uſe. To roll the humerus outwards; to aſſiſt in raiſing, and in ſupporting it when raiſed; and to pull the ligament from between the bones.

N. B. Theſe two muſcles are covered with a tendinous membrane, from which a number of their fleſhy fibres ariſe. It ſerves beſides to ſtrengthen their actions, and keeps them from ſwelling too much outwardly when in action.

3. Teres Minor,

Arifes, fleſhy, from all the round edge of the inferior coſta of the ſcapula, and runs forwards along the inferior edge of the infraſpinatus muſcle, and adheres to the ligament.

Inſerted, tendinous, into the back-part of the large protuberance on the head of the os humeri, a little behind and below the termination of the laſt named muſcles.

Uſe. To roll the humerus outwards; to draw the humerus backwards; and to prevent the ligament from being pinched between the bones.

Along the inferior coſta of the ſcapula is,

Teres

TERES MAJOR,

Arifes, fleſhy, from the inferior angle of the ſcapula, and from all that portion of its inferior coſta that is rough and thicker than the reſt; its fleſhy fibres are continued over part of the infraſpinatus muſcle, to which they firmly adhere.

Inſerted, by a broad, ſhort, and thin tendon, into the ridge at the inner ſide of the groove for lodging the tendon of the long head of the biceps, along with the latiſſimus dorſi.

Uſe. To roll the humerus inwards, and to draw it backwards and downwards.

The two before the ſcapula are,

1. DELTOIDES,

Arifes, fleſhy, from all the poſterior part of the clavicle that the pectoralis major does not poſſeſs; tendinous and fleſhy, from the acromion, and lower margin of almoſt the whole ſpine of the ſcapula oppoſite to the inſertion of the cucullaris muſcle: From theſe origins it runs in three different directions, i. e. from the clavicle outwards and downwards; from the ſpine of the ſcapula outwards, forwards, and downwards; and from the acromion ſtraight downwards; and is compoſed of a number of faſciculi, which form a ſtrong fleſhy muſcle that covers the anterior part of the joint of the os humeri.

Inſerted, tendinous, into a rough protuberance in the outer ſide of the os humeri, near its middle, where the fibres of this muſcle intermix with ſome part of the brachialis externus.

Uſe. To pull the arm directly outwards and upwards, and a little forwards or backwards, according to the different directions of its fibres.

2. Co-

2. Coraco-Brachialis,

Ariſes, tendinous and fleſhy, from the fore-part of the coracoid proceſs of the ſcapula; adhering, in its deſcent, to the ſhort head of the biceps.

Inſerted, tendinous and fleſhy, about the middle of the internal part of the os humeri, near the origin of the third head of the triceps, called *brachialis externus*, where it ſends down a thin tendinous expanſion to the internal condyle of the os humeri.

Uſe. To raiſe the arm upwards and forwards.

N. B. There paſſes a nerve through this muſcle, called *Muſculo cutaneus*.

The one beneath the ſcapula is,

Subscapularis,

Ariſes, fleſhy, from all the baſe of the ſcapula internally, and from its ſuperior and inferior coſtae, being compoſed of a number of tendinous and fleſhy faſciculi, which make prints on the bone; they all join together, fill up the hollow of the ſcapula, and paſs over the joint adhering to the capſular ligament.

Inſerted, tendinous, into the upper part of the internal protuberance at the head of the os humeri.

Uſe. To roll the humerus inwards, and to draw it to the ſide of the body; and to prevent the capſular ligament from being pinched.

CHAP.

CHAP. XXVI.

Muscles *ſituated on the* Os Humeri.

THESE are called

Muſcles of the *Cubit* or *Fore-arm.*

They conſiſt of two before, and two behind.

Before are,

1. Biceps Flexor Cubiti,

Ariſes, by two heads. The firſt and outermoſt, called *longus*, begins tendinous from the upper edge of the glenoid cavity of the ſcapula; paſſes over the head of the os humeri within the joint; and, in its deſcent without the joint, is incloſed in a groove near the head of the os humeri, by a membranous ligament that proceeds from the capſular ligament and adjacent tendons. The ſecond, or innermoſt head, called *brevis*, ariſes, tendinous and fleſhy, from the coracoid proceſs of the ſcapula, in common with the coraco-brachialis muſcle. A little below the middle of the fore part of the os humeri, theſe heads unite.

Inſerted, by a ſtrong roundiſh tendon, into the tubercle of the upper end of the radius internally.

Uſe. To turn the hand ſupine, and to bend the fore-arm.

N. B. At the bending of the elbow, where it begins to grow tendinous, it ſends off an aponeuroſis, which covers all

all the muſcles on the inſide of the fore-arm, and joins with another tendinous membrane, which is ſent off from the triceps extenſor cubiti, and covers all the muſcles on the outſide of the fore-arm, and a number of the fibres, from oppoſite fibres, decuſſate each other. It ſerves to ſtrengthen the muſcles, by keeping them from ſwelling too much outwardly, when in action; and a number of their fleſhy fibres take their origin from it.

Biceps brachii, Albinus.

Coraco-radialis, ſeu *biceps*, Winſlow.

Biceps-internus, Douglas.

2. Brachialis Internus,

Ariſes, fleſhy, from the middle of the os humeri, at each ſide of the inſertion of the deltoid muſcle, covering all the inferior and fore-part of this bone, runs over the joint, and adheres firmly to the ligament.

Inſerted, by a ſtrong ſhort tendon, into the coronoid proceſs of the ulna.

Uſe. To bend the fore-arm, and to prevent the capſular ligament of the joint from being pinched.

Brachialis, Winſlow.

Behind are,

1. Triceps Extensor Cubiti,

Ariſes, by three heads; the firſt, called *longus*, ſomewhat broad and tendinous, from the inferior coſta of the ſcapula, near its cervix. The ſecond head called *brevis*, ariſes by an acute, tendinous, and fleſhy beginning, from the back-part of the os humeri, and a little below its head, outwardly. The third, called *brachialis externus*, ariſes by an acute beginning, from the back-part of the os humeri. Theſe three heads unite lower than the inſertion of the te-

res

res major, and cover the whole posterior part of the humerus, from which they receive addition in their descent.

Inserted into the upper and external part of the process of the ulna, called *olecranon*, and partly into the condyles of the os humeri, adhering firmly to the ligament.

Use. To extend the fore-arm.

Anconeus major, *Anconeus externus*, and *Anconeus internus*, Winslow.

Biceps externus, and *Brachialis externus*, Douglas.

3. Anconeus,

Arises, tendinous, from the posterior part of the external condyle of the os humeri; it soon grows fleshy, and is continued from the third head of the triceps.

Inserted, fleshy, and thin, into a ridge on the outer and posterior edge of the ulna, being continued some way below the olecranon, and covered with a tendinous membrane.

Use. To assist in extending the fore-arm.

Anconeus minor, Winslow.

Anconeus, vel *Cubitalis Riolani*, Douglas.

CHAP. XXVII.

Muscles *situated on the* Cubit *or* Fore-Arm *.

THESE may be divided into three classes; *first*, flexors and extensors of the whole *hand*; *second*, flexors and extensors

* In the following description, the arm is supposed to hang by the side with the palm turned forwards; so that the radius and thumb are upon its outer side, and the ulna and little finger upon its inner side; whereas, when the muscles are described in the less straining posture of pronation, as has been generally done by authors, the utmost confusion is necessarily introduced in the application of the terms *outer* and *inner*, from the decussation of the radius and ulna.

extenſors of the *fingers;* and *third*, ſupinators and pronators, or thoſe that roll the *radius on the ulna.*

Firſt claſs conſiſts of three flexors, and three extenſors.

Flexors:

1. Palmaris Longus,

Ariſes, tendinous, from the internal condyle of the os humeri, ſoon grows fleſhy, and after a ſhort progreſs, ſends off a long ſlender tendon.

Inſerted into the ligamentum carpi annulare, and into a tendinous membrane that is expanded on the palm of the hand, named *aponeuroſis palmaris;* which, above, begins at the tranſverſe or annular ligament of the wriſt, and, below, is fixed to the roots of the fingers.

Uſe. To bend the hand, and to ſtretch the membrane that is expanded on the palm.

Ulnaris gracilis, Winſlow.

N. B. This muſcle is ſometimes wanting; but the aponeuroſis palmaris is always to be found, and a ſmall muſcle named

Palmaris Brevis,

Ariſes from the ligamentum carpi annulare, and tendinous membrane that is expanded on the palm of the hand.

Inſerted, by ſmall bundles of fleſhy fibres, into the ſkin and fat that covers the abductor minimi digiti, and into the os piſiforme.

Uſe. To aſſiſt in contracting the palm of the hand.

Palmaris cutaneus, Winſlow.

2. Flexor

2. Flexor Carpi Radialis,

Ariſes, tendinous and fleſhy, from the internal condyle of the os humeri, and from the anterior part of the upper end of the ulna, where it firmly adheres to the pronator radii teres.

Inſerted, by a flat tendon, into the fore and upper part of the metacarpal bone that ſuſtains the fore finger, after running through a foſſa in the os trapezium.

Uſe. To bend the hand, and to aſſiſt in its pronation.

Radialis internus, Albinus and Winſlow.

3. Flexor Carpi Ulnaris,

Ariſes, tendinous, from the internal condyle of the os humeri. It has likewiſe a ſmall fleſhy beginning from the outer ſide of the olecranon; between which and the condyle the ulnar nerve paſſes to the fore-arm; and a number of its fleſhy fibres ariſe from the tendinous membrane which covers the fore-arm.

Inſerted, by a ſhort ſtrong tendon, into the os piſiforme; at a little diſtance from its inſertion, a ſmall ligament is ſent off to the metacarpal bone that ſuſtains the little finger.

Uſe. To aſſiſt the former in bending the arm.

Ulnaris internus, Albinus and Winſlow.

Extenſors are,

1. Extensor Carpi Radialis Longior,

Ariſes, broad, thin, and fleſhy, immediately below the ſupinator radii longus, from the lower part of the external ridge of the os humeri, above its external condyle.

Inſerted, by a round tendon, into the poſterior and upper part of the metacarpal bone that ſuſtains the fore-finger.

Uſe. To extend and bring the hand backwards.

Radialis externus longior, Albinus.

Radialis externus primus, Winſlow.

2. Extensor Carpi Radialis Brevior,

Ariſes, tendinous, from the external condyle of the os humeri, and from the ligament that connects the radius to it, and runs along the outſide of the radius.

Inſerted, by a round tendon, into the upper and back part of the metacarpal bone that ſuſtains the middle finger.

Uſe. To aſſiſt the laſt-mentioned muſcle.

Radialis externus brevior, Albinus.

Radialis ſecundus, Winſlow.

3. Extensor Carpi Ulnaris,

Ariſes, tendinous, from the external condyle of the os humeri; and, in its progreſs, fleſhy from the middle of the ulna, where it paſſes over the ulna. Its round tendon is incloſed by a membranous ſheath, in a groove which is ſituated at the extremity of the ulna.

Inſerted, by its round tendon, into the poſterior and upper part of the metacarpal bone that ſuſtains the little-finger.

Uſe. To aſſiſt the former in extending the hand.

Ulnaris externus, Albinus and Winſlow.

Second Claſs.

The flexors and extenſors of the four fingers are two long, and one ſmall flexor to each finger, and one extenſor.

1. Flexor Sublimis Perforatus,

Ariſes, tendinous and fleſhy, from the internal condyle of the os humeri; tendinous from the coronoid proceſs of the

the ulna, near the edge of the cavity that receives the head of the radius; fleshy from the tubercle of the radius; and membranous and fleshy from the middle of the fore-part of the radius, where the flexor pollicis longus arises. Its fleshy belly sends off four round tendons before it passes under the ligament of the wrist.

Inserted into the anterior and upper part of the second bone of each finger, being, near the extremity of the first bone, divided for the passage of the perforans.

Use. To bend the second joint or phalanx of the fingers.

Sublimus, Albinus.

Perforatus, Douglas.

2. Flexor Profundus Perforans,

Arises, fleshy, from the external side, and upper part of the ulna, for some way downwards, and from a large share of the interosseous ligament. It splits into four tendons, a little before it passes under the ligamentum carpi annulare; and these pass through the slits in the tendons of the flexor sublimis.

Inserted into the fore and upper part of the third or last bone of all the four fingers.

Use. To bend the last joint of the fingers.

Profundus, Albinus.

Perforans, Douglas.

The four small flexors are named

Lumbricales.

Arises, thin and fleshy, from the outside of the tendons of the flexor profundus, a little above the lower edge of the ligamentum carpi annulare.

Inserted, by long slender tendons, into the outer sides of the

the broad tendons of the interoſſei muſcles, about the middle of the firſt joint.

Uſe. To increaſe the flexion of the fingers while the long flexors are in full action.

Extenſors are,

EXTENSOR DIGITORUM COMMUNIS,

Ariſes, by an acute, tendinous, and fleſhy beginning, from the external condyle of the os humeri, where it adheres to the ſupinator radii brevis. Before it paſſes under the ligamentum carpi annulare externum, it ſplits into four tendons; ſome of which may be divided into ſeveral ſmaller: and about the fore-part of the metacarpal bones they remit tendinous filaments to each other.

Inſerted into the poſterior part of all the bones of the four fingers by a tendinous expanſion.

Uſe. To extend all the joints of the fingers.

Third Claſs,

Conſiſts of four muſcles, viz, two *ſupinators*, and two *pronators*.

Supinators are,

1. SUPINATOR RADII LONGUS,

Ariſes, by an acute and fleſhy origin, from the external ridge of the os humeri, above the external condyle, near as far up as the middle of that bone.

Inſerted into the outer-ſide of the inferior extremity of the radius.

Uſe. To roll the radius outwards, and conſequently the palm of the hand upwards.

Supinator longus, Albinus, Winſlow, and Douglas.

2. SU-

2. Supinator Radii Brevis,

Ariſes, tendinous, from the external condyle of the os humeri; tendinous and fleſhy, from the external and upper part of the ulna, and adheres firmly to the ligament that joins theſe two bones.

Inſerted into the head, neck, and tubercle of the radius, near the inſertion of the biceps, and ridge running from that downwards and outwards.

Uſe. To roll the radius outwards, and ſo bring the hand ſupine.

Pronators are,

1. Pronator Radii Teres,

Ariſes, fleſhy, from the internal condyle of the os humeri, and tendinous from the coronoid proceſs of the ulna.

Inſerted, thin, tendinous, and fleſhy, into the middle of the poſterior part of the radius.

Uſe. To roll the radius, together with the hand, inwards.

2. Pronator Radii Quadratus,

Ariſes, broad, tendinous, and fleſhy, from the lower and inner part of the ulna; the fibres run tranſverſely, to be

Inſerted into the lower and anterior part of the radius, oppoſite to its origin.

Uſe. To turn the radius, together with the hand, inwards.

CHAP.

CHAP. XXVIII.

Muscles *situated on the* Hand *chiefly.*

THESE may be divided into four classes, viz. muscles of the *thumb*, *fore-finger*, *little-finger*, and *metacarpal bones*.

Muscles of the Thumb.

These consist of *three flexors*, *three extensors*, *one abductor*, and *one adductor*.

Flexors are,

1. Flexor Longus Pollicis Manus,

Arises, by an acute fleshy beginning, from the upper part of the radius, immediately below its tubercle, and is continued down for some space on the fore-part of this bone. It has likewise generally another origin from the internal condyle of the os humeri, which forms a distinct fleshy slip that terminates near the upper part of the origin from the radius.

Inserted into the last joint of the thumb, after having passed its tendon under the ligament of the wrist.

Use. To bend the last joint of the thumb.

Flexor tertii internodii, Douglas.

2. Flexor Brevis Pollicis Manus,

Arises from the os trapezoides, magnum, and unciforme of

of the carpus, and is divided into two portions by the tendon of the flexor pollicis longus.

Inserted into the ossa sesamoidea and first bone of the thumb.

Use. To bend the first joint of the thumb.

Flexor secundi internodii, Douglas.

3. Flexor Ossis Metacarpi Pollicis, *or* Opponens Pollicis,

Arises, fleshy, from the os trapezium and ligamentum carpi annulare, lying under the abductor pollicis.

Inserted, tendinous and fleshy, into the under and anterior part of the metacarpal bone of the thumb.

Use. To bring the thumb inwards, opposite to the other fingers.

Flexor primi internodii, Douglas.

Extensors are,

1. Extensor Ossis Metacarpi Pollicis Manus,

Arises, fleshy, from the middle and posterior part of the ulna, immediately below the insertion of the anconaeus muscle, from the posterior part of the middle of the radius, and from the interosseous ligament.

Inserted, generally by two tendons, into the os trapezium, and upper back-part of the metacarpal bone of the thumb, and often joins with the abductor pollicis.

Use. To extend the metacarpal bone of the thumb outwardly.

Abductor longus pollicis manus, Albinus.

Extensor primi internodii, Douglas.

2. Ex-

2. Extensor Primi Internodii,

Ariſes, fleſhy, from the poſterior part of the ulna near the former muſcle, and from the interoſſeous ligament.

Inſerted, tendinous, into the poſterior part of the firſt bone of the thumb; and part of it may be traced as far as the ſecond bone.

Uſe. To extend the firſt bone of the thumb obliquely outwards.

Extenſor minor pollicis manus, Albinus.

This and the preceding muſcle is called

Extenſor pollicis primus, Winſlow.

Extenſor ſecundi internodii, Douglas.

3. Extensor Secundi Internodii,

Ariſes, by an acute, tendinous, and fleſhy beginning, from the middle back part of the ulna, and from the interoſſeous ligament; its tendon runs through a ſmall groove at the inner and back part of the lower end of the radius.

Inſerted into the laſt bone of the thumb.

Uſe. To extend the laſt joint of the thumb obliquely backwards.

Extenſor major pollicis manus, Albinus.

Extenſor pollicis ſecundus, Winſlow.

Extenſor tertii internodii, Douglas.

Abductor Pollicis Manus,

Ariſes, by a broad tendinous and fleſhy beginning, from the ligamentum carpi annulare, and from the os trapezium.

Inſerted, tendinous, into the outer ſide of the root of the firſt bone of the thumb.

Uſe. To draw the thumb from the fingers.

N. B.

N. B. Albinus names the inner portion of this muſcle *Abductor brevis alter.*

Abductor, Thenar Riolani, Douglas.

Adductor Pollicis Manus,

Ariſes, fleſhy, from almoſt the whole length of the metacarpal bone that ſuſtains the middle finger; from thence its fibres are collected together.

Inſerted, tendinous, into the inner part of the root of the firſt bone.

Uſe. To pull the thumb towards the fingers.

Adductor ad minimum digitum, Douglas.

Fore-finger:

Indicator,

Ariſes, by an acute fleſhy beginning, from the middle of the poſterior part of the ulna; its tendon paſſes under the ſame ligament with the extenſor digitorum communis, with part of which it is

Inſerted into the poſterior part of the fore-finger.

Extenſor ſecundi internodii indicis proprius, vulgo indicator, Douglas.

Abductor Indicis Manus,

Ariſes, from the os trapezium, and from the ſuperior part and inner ſide of the metacarpal bone of the thumb.

Inſerted, by a ſhort tendon, into the outer and back-part of the firſt bone of the fore-finger.

Uſe. To bring the fore-finger towards the thumb.

Semi-interoſſeus, Winſlow.

Little finger:

ABDUCTOR MINIMI DIGITI MANUS,

Ariſes, fleſhy, from the os piſiforme, and from that part of the ligamentum carpi annulare next it.

Inſerted, tendinous, into the inner ſide of the upper end of the firſt bone of the little finger.

Uſe. To draw this finger from the reſt.

Hypothenar minor, Winſlow.

Extenſor tertii internodii minimi digiti, Douglas.

ADDUCTOR METACARPI MINIMI DIGITI MANUS,

Ariſes, fleſhy, from the thin edge of the os unciforme, and from that part of the ligament of the wriſt next it.

Inſerted, tendinous, into the inner ſide and anterior part of the metacarpal bone of this finger.

Uſe. To bend and bring the metacarpal bone of this finger towards the reſt.

Metacarpus, Winſlow.

Flexor primi internodii minimi digiti, Douglas.

FLEXOR PARVUS MINIMI DIGITI,

Ariſes, fleſhy, from the outer ſide of the os unciforme, and from the ligament of the wriſt which joins with that bone.

Inſerted, by a roundiſh tendon, into the inner and anterior part of the upper end of the firſt bone of this finger.

Uſe. To bend the little finger, and aſſiſt the adductor.

Abductor minimi digiti, Hypothenar Riolani, Douglas.

Between the metacarpal bones, there are *four internal* and *three external* muſcles named *interoſſei*.

Interoſſei

Interoſſei interni:

1. Prior Indicis,

Ariſes, tendinous and fleſhy, from the upper and outer part of the metacarpal bone that ſuſtains the fore-finger.

Inſerted into the outſide of that part of the tendinous expanſion from the extenſor digitorum communis, which covers the poſterior part of the fore-finger.

Uſe. To draw the fore-finger inwards towards the thumb, and extend it obliquely.

Extenſor tertii internodii indicis, Douglas.

2. Posterior Indicis,

Ariſes, tendinous and fleſhy, from the root and inner part of the metacarpal bone that ſuſtains the fore-finger.

Inſerted into the inner ſide of the tendinous expanſion which is ſent off from the extenſor digitorum communis along the poſterior part of the fore-finger.

Uſe. To extend the fore-finger obliquely, and to draw it outwards.

Firſt interoſſeus, Douglas.

3. Prior Annularis,

Ariſes, from the root of the outſide of the metacarpal bone that ſuſtains the ring-finger.

Inſerted into the outſide of the tendinous expanſion of the extenſor digitorum communis which covers the ring-finger.

Uſe. To extend and pull the ring-finger towards the thumb.

Fourth interoſſeus, Douglas.

4. Interosseus Auricularis,

Ariſes from the root and outer ſide of the metacarpal bone of the little finger; and is

Inſerted

Inserted into the outside of the tendinous expansion of the extensor digitorum communis, which covers the posterior part of the little finger.

Use. To extend and draw the little finger outwards.

Sixth interosseus, Douglas.

Interossei externi, seu bicipites :

1. Prior Medii,

Arises, by two origins, from the roots of the metacarpal bones that sustain the fore and middle fingers externally, and next each other : Runs along the outside of the middle finger ; and, being conspicuous on both sides of the hand, is

Inserted into the outside of the tendinous expansion from the extensor digitorum communis, which covers the posterior part of the middle finger.

Use. To extend, and to draw the middle finger inwards.

Second interosseus, Douglas.

2. Posterior Medii,

Arises, by two origins, from the roots of the metacarpal bones, next each other, that sustain the middle and ring-fingers.

Inserted into the inside of the tendinous expansion from the extensor digitorum communis, which runs along the posterior part of the middle finger.

Use. To extend and draw the middle-finger outwards.

Third interosseus, Douglas.

3. Posterior Annularis,

Arises, by two origins, from the roots of the metacarpal bones that sustain the ring and little fingers next each other.

Inserted

Inserted into the inside of the tendinous expansion of the extensor digitorum communis, which runs along the posterior part of the ring-finger.

Use. To extend and draw the ring-finger inwards.

Fifth interosseus, Douglas.

N. B. The internal interossei are only conspicuous on the palm of the hand; but the external are apparent on both the palm and back of the hand.

CHAP. XXIX.

Muscles *of the* Inferior Extremities.

THESE may be divided into the muscles *situated* on the *outside* of the *pelvis*, on the *thigh*, on the *leg*, and on the *foot*.

Muscles on the outside of the *pelvis*, which are called *muscles of the thigh*.

These are composed of *one layer before* and *three layers behind*.

The layer before consists of five muscles:

1. Psoas Magnus.
2. Iliacus Internus. } These were described, p. 311 & 312.

3. Pectinalis,

Arises, broad and fleshy, from the upper and anterior part of the os pubis or pectinis, immediately above the foramen thyroideum.

Inserted into the anterior and upper-part of the linea aspera

pera of the os femoris, a little below the trochanter minor, by a flat and ſhort tendon.

Uſe. To bring the thigh upwards, and inwards, and to give it a degree of rotation outwards.

Pectineus, Albinus.

4. Triceps Adductor Femoris,

Under this appellation are comprehended three diſtinct muſcles:

1. Adductor Longus Femoris,

Ariſes, by a ſtrong roundiſh tendon, from the upper and interior part of the os pubis, and ligament of its ſynchondroſis, on the inner ſide of the pectinalis.

Inſerted, tendinous, near the middle of the poſterior part of the linea aſpera, being continued for ſome way down.

Adductor femoris primus, Douglas.

Triceps minus, Winſlow.

2. Adductor Brevis Femoris,

Ariſes, tendinous, from the os pubis near its joining with the oppoſite os pubis below and behind the former.

Inſerted, tendinous and fleſhy, into the inner and upper part of the linea aſpera, from a little below the trochanter minor, to the beginning of the inſertion of the adductor longus.

Adductor femoris ſecundus, Douglas.

Triceps ſecundus, Winſlow.

3. Adductor Magnus Femoris,

Ariſes, a little lower down than the former, near the ſymphaſis of the oſſa pubis; tendinous and fleſhy, from the tuberoſity of the os iſchium; the fibres run outwards and downwards.

Inſerted

Inſerted into almoſt the whole length of the linea aſpera; into a ridge above the internal condyle of the os femoris; and, by a roundiſh long tendon, into the upper part of that condyle, a little above which the femoral artery takes a ſpinal turn towards the ham, paſſing between this muſcle and the bone.

Uſe of theſe three muſcles or triceps. To bring the thigh inwards and upwards, according to the different directions of their fibres; and, in ſome degree, to roll the thigh outwards

Adductor femoris tertius, and
Adductor femoris quartus, Douglas.
Triceps tertius, Winſlow.

5. Obturator Externus,

Ariſes, fleſhy, from the lower fore-part of the os pubis, and fore-part of the inner crus of the iſchium; ſurrounds the foramen thyroideum; a number of its fibres, ariſing from the membrane which fills up that foramen, are collected like rays towards a centre, and paſs outwards around the root of the back-part of the cervix of the os femoris.

Inſerted, by a ſtrong tendon, into the cavity at the inner and back-part of the root of the trochanter major, adhering in its courſe to the capſular ligament of the thigh-bone.

Uſe. To roll the thigh-bone obliquely outwards, and to prevent the capſular ligament from being pinched.

Behind are,

Firſt Layer.

Gluteus Maximus,

Ariſes, fleſhy, from the poſterior part of the ſpine of the

os

os ilium, a little higher up than the joining of the ilium with the os ſacrum, from the whole external ſide of the os ſacrum, below the poſterior ſpinous proceſs of the os ilium; from the poſterior ſacro-iſchiatic ligament, over which part of the inferior edge of this muſcle hangs in a folded manner; from the os coccygis. All the fleſhy fibres run obliquely forwards, and a little downwards, to form a thick broad muſcle, which is divided into a number of ſtrong faſciculi. The upper part of it covers almoſt the whole of the trochanter major, between which and the tendon of this muſcle there is a large burſa mucoſa, and where it is inſeparably joined to the broad tendon of the *tenſor vaginae femoris*.

Inſerted, by a ſtrong, thick, and broad tendon, into the upper and outer part of the linea aſpera, which is continued from the trochanter major, for ſome way downwards.

Uſe. To extend the thigh, by pulling it directly backwards, and a little outwards.

Gluteus magnus, Albinus.

Gluteus major, Cowper.

Second Layer.

GLUTEUS MEDIUS,

Ariſes, fleſhy, from the anterior ſuperior ſpinous proceſs of the os ilium, and from all the outer edge of the ſpine of the ilium, except its poſterior part, where it ariſes from the dorſum of that bone.

Inſerted, by a broad tendon, into the outer and poſterior part of the trochanter major.

Uſe. To draw the thigh-bone outwards, and a little backwards; to roll the thigh-bone outwards, eſpecially when it is bended.

N. B.

N. B. The anterior and upper part of this muſcle is covered by a tendinous membrane, from which a number of its fleſhy fibres ariſe, and which joins with the broad tendons of the gluteus maximus, tenſor vaginae femoris, and latiſſimus dorſi.

Third Layer conſiſts of four Muſcles.

1. GLUTEUS MINIMUS,

Ariſes, fleſhy, from a ridge that is continued from the ſuperior anterior ſpinous proceſs of the os ilium, and from the middle of the dorſum of that bone as far back as its great niche.

Inſerted, by a ſtrong tendon, into the fore and upper part of the trochanter major.

Uſe. To aſſiſt the former in pulling the thigh outwards and backwards, and in rolling it.

Gluteus minor, Albinus.

2. PYRIFORMIS,

Ariſes, within the pelvis, by three tendinous and fleſhy origins, from the ſecond, third, and fourth pieces of the os ſacrum; from thence growing gradually narrower, it paſſes out of the pelvis along with the poſterior crural nerve, below the niche in the poſterior part of the os ilium, where it receives a few fleſhy fibres.

Inſerted, by a roundiſh tendon, into the upper part of the cavity at the inner ſide of the root of the trochanter major.

Uſe. To move the thigh a little upwards, and roll it outwards.

Pyriformis, ſeu *iliacus externus*, Douglas.

3. GEMINI,

Ariſes by two diſtinct origins; the ſuperior from the ſpi-

nous proceſs, and the inferior from the tuberoſity of the os iſchium; alſo, from the poſterior ſacro-iſchiatic ligament. They are both united by a tendinous and fleſhy membrane, and form a purſe for the tendon of the obturator internus muſcle, which was formerly deſcribed.

Inſerted, tendinous and fleſhy, into the cavity at the inner ſide of the root of the trochanter major, on each ſide of the tendon of the obturator internus, to which they firmly adhere.

Uſe. To roll the thigh outwards, and to preſerve the tendon of the obturator internus from being hurt by the hardneſs of that part of the iſchium over which it paſſes; alſo, to hinder it from ſtarting out of its place, while the muſcle is in action.

Gemelli, Winſlow.

4. Quadratus Femoris,

Ariſes, tendinous and fleſhy, from the outſide of the tuberoſity of the os iſchium; and, running tranſverſely, is

Inſerted, fleſhy, into a rough ridge, continued from the root of the large trochanter to the root of the ſmall one.

Uſe. To roll the thigh outwards.

CHAP. XXX.

Muscles *ſituated on the* Thigh.

THESE are called *muſcles* of the *leg*; and conſiſt of *one*, on the *outſide*; *two*, on the *inſide*; *four*, *before*; and *four*, *behind*.

Previous

Previous to the deſcription of the muſcles that are ſituated on the thigh and leg, it is neceſſary to take notice of a broad tendinous faſcia or ſheath, which is ſent off from the back and from the tendons of the glutei and adjacent muſcles.

It is a ſtrong thick membrane on the outſide of the thigh and leg; but, towards the inſide of both, it gradually turns thinner, and has rather the appearance of cellular ſubſtance, than a tendinous membrane. A little below the trochanter major, it is firmly fixed to the linea aſpera; and, farther down, to that part of the head of the tibia that is next the fibula; where it ſends off the tendinous expanſion along the outſide of the leg.

It ſerves to ſtrengthen the action of the muſcles, by keeping them firm in their proper places while in action, particularly the tendons that paſs over the joints, where this membrane is thickeſt; and it gives origin to a number of the fleſhy fibres of the muſcles.

On the outſide is,

Tensor Vaginæ Femoris,

Ariſes, by a narrow tendinous, and fleſhy beginning, from the external part of the anterior ſuperior ſpinous proceſs of the os ilium.

Inſerted, a little below the trochanter major, into the inner ſide of the membranous faſcia which covers the outſide of the thigh.

Uſe. To ſtretch the membranous faſcia, to aſſiſt in the abduction of the thigh, and ſomewhat in its rotation inwards.

Muſculus faſciae latae, Winſlow.

On

On the inside are,

1. Sartorius,

Arises, tendinous, from the anterior superior spinous process of the os ilium, soon grows fleshy, runs down for some space upon the rectus, and going obliquely inwards, it passes over the vastus internus, and about the middle of the os femoris, over part of the triceps, it runs down farther between the tendon of the abductor magnus and that of the gracilis muscle.

Inserted, by a broad and thin tendon, into the inner side of the tibia, near the inferior part of its tubercle.

Use. To bend the leg obliquely inwards, or to bring one leg across the other.

2. Gracilis,

Arises, by a thin tendon, from the os pubis near the symphysis of these two bones: soon grows fleshy; and, descending by the inside of the thigh, is

Inserted, tendinous, into the tibia under the sartorius.

Use. To assist the sartorius.

Gracilis internus, five *Rectus internus*, Winslow.

Before are,

1. Rectus,

Arises, fleshy, from the inferior anterior spinous process of the os ilium, and tendinous from the dorsum of the ilium, a little above the acetabulum; runs down over the anterior part of the cervix of the os femoris, the fibres not being straight, but running down like the plumage of a feather obliquely outwards and inwards, from a tendon in the middle.

Inserted,

Inſerted, tendinous, into the upper part of the patella, from which a thin tendon runs down, on the fore-part of this bone, to terminate in a thick ſtrong ligament, which is ſent off from the inferior part of the patella, and inſerted into the tubercle of the tibia.

Uſe. To extend the leg, and in a powerful manner, by the intervention of the patella, like a pully.

Rectus, ſive *Gracilis anterior*, Winſlow.

2. Vastus Externus,

Ariſes, broad, tendinous, and fleſhy, from the root of the trochanter major, and upper part of the linea aſpera, its origin being continued from near the inſertion of the gluteus minimus, the whole length of the linea aſpera, by fleſhy fibres which run obliquely forwards to a middle tendon, where they terminate.

Inſerted into a large ſhare of the upper part of the patella; and part of it ends in an aponeuroſis, which is continued down to the leg, and in its paſſage is firmly fixed to the head of the tibia.

Uſe. To extend the leg.

3. Vastus Internus,

Ariſes, tendinous and fleſhy, from between the fore-part of the os femoris and root of the trochanter minor, and from almoſt all the inſide of the linea aſpera, by fibres running obliquely forwards and downwards.

Inſerted, tendinous, into the upper and inſide of the patella, continuing fleſhy lower than the vaſtus externus. Part of it likewiſe ends in an aponeuroſis continued down to the leg, and fixed in its paſſage to the upper part of the tibia.

Uſe. To extend the leg.

4. Cru-

4. CRURALIS,

Ariſes, fleſhy, from between the two trochanters of the os femoris, but nearer the minor, and firmly adhering to moſt of the fore-part of the os femoris, and connected to both vaſti muſcles.

Inſerted, tendinous, into the upper part of the patella, behind the rectus.

Uſe. To aſſiſt in the extenſion of the leg.

Cruralis, Albinus.

N. B. Theſe four muſcles *before*, being inſerted into the patella, have the ſame effect upon the leg, as if they were immediately inſerted into it, by means of the ſtrong tendon, or rather ligament, which is ſent off from the inferior part of the patella to the tibia.

Behind are,

1. SEMITENDINOSUS,

Ariſes, tendinous and fleſhy, in common with the long head of the biceps, from the poſterior part of the tuberoſity of the os iſchium; and ſending down a long roundiſh tendon, which ends flat, is

Inſerted into the inſide of the ridge of the tibia, a little below its tubercle.

Uſe. To bend the leg backwards and a little inwards.

Seminervoſus, Winſlow and Douglas.

2. SEMIMEMBRANOSUS,

Ariſes, tendinous, from the upper and poſterior part of the tuberoſity of the os iſchium, ſends down a broad flat tendon, which ends in a fleſhy belly, and, in its deſcent, runs at firſt on the fore-part of the biceps, and, lower, between it and the ſemitendinoſus.

Inſerted,

Inserted, tendinous, into the inner and back-part of the head of the tibia.

Use. To bend the leg, and bring it directly backwards.

N. B. The two last form what is called the *inner hamstring*.

3. Biceps Flexor Cruris,

Arises by two distinct heads. The first, called *longus*, arises, in common with the semitendinosus, from the upper and posterior part of the tuberosity of the os ischium. The second, called *brevis*, arises from the linea aspera, a little below the termination of the gluteus maximus, by a fleshy acute beginning, which soon grows broader as it descends to join with the first head, a little above the external condyle of the os femoris.

Inserted, by a strong tendon, into the upper part of the head of the fibula.

Use. To bend the leg.

Biceps cruris, Albinus.

Biceps, Winslow and Douglas.

N. B. This muscle forms what is called the *outer hamstring*; and between it and the inner, the nervus popliteus, and arteria and vena poplitea, are situated.

4. Popliteus,

Arises, by a round tendon, from the lower and back part of the external condyle of the os femoris; then runs over the ligament that involves the joint, firmly adhering to it, and part of the semilunar cartilage. As it runs over the joint, it becomes fleshy, and the fibres run obliquely inwards, being covered with a thin tendinous membrane.

Inserted, broad, thin, and fleshy, into a ridge at the upper and internal edge of the tibia, a little below its head.

Use. To assist in bending the leg, and to prevent the capsular

ſular ligament from being pinched. After the leg is bent, this muſcle ſerves to roll it inwards.

CHAP. XXXI.

Muscles *ſituated on the* Leg.

THESE are called *Muſcles* of the *Foot*; and may be divided into *two claſſes*, viz. *Extenſors* and *Flexors* of the *Foot*. 2. *Common Extenſors* and *Flexors* of the *Toes*.

Firſt Claſs.

Extenſors.

Theſe conſiſt of three:

1. Gastrocnemius Externus, *ſeu* Gemellus,

Ariſes by two diſtinct heads. The firſt head ariſes from the upper and back part of the internal condyle of the os femoris, and from that bone, a little above its condyle, by two diſtinct tendinous origins. The ſecond head ariſes tendinous from the upper and back part of the external condyle of the os femoris. A little below the joint, their fleſhy bellies unite in a middle tendon; and, below the middle of the tibia, it ſends off a broad thin tendon, which joins a little above the extremity of the tibia with the tendon of the following.

2. Soleus, *ſeu* Gastrocnemius Internus,

Ariſes by two origins. The firſt is from the upper and back part of the head of the fibula, continuing to receive many

many of its fleſhy fibres from the poſterior part of that bone for ſome ſpace below its head. The other origin begins from the poſterior and upper part of the middle of the tibia; and runs inwards along the inferior edge of the popliteus towards the inner part of the tibia, from which it receives fleſhy fibres for ſome way down. The fleſh of this muſcle, covered by the tendon of the gemellus, runs down near as far as the extremity of the tibia; a little above which the tendons of both gaſtrocnemii unite, and form a ſtrong round chord, which is called *tendo Achillis*.

Inſerted, into the upper and poſterior part of the os calcis; by the projection of which the tendo Achillis is at a conſiderable diſtance from the tibia.

Uſe. To extend the foot, by bringing it backwards and downwards.

Gemellus and *Soleus*, Albinus.

Gaſtrocnemii and *Soleus*, Winſlow.

Extenſor tarſi ſuralis, vel *Extenſor magnus*, Douglas.

3. Plantaris,

Ariſes, thin and fleſhy, from the upper and back part of the root of the external condyle of the os femoris, near the inferior extremity of that bone, adhering to the ligament that involves the joint in its deſcent. It paſſes along the ſecond origin of the ſoleus, and under the gemellus, where it ſends off a long, ſlender, thin tendon, which comes from between the great extenſors, where they join tendons; then runs down by the inſide of the tendo Achillis.

Inſerted into the inſide of the poſterior part of the os calcis, below the tendo Achillis.

Uſe. To aſſiſt the former, and to pull the capſular ligament of the knee from between the bones. It ſeems likewiſe to aſſiſt in rolling the foot inwards.

Tibialis gracilis, vulgo *Plantaris*, Winſlow.

Extenſor tarſi minor, vulgo *Plantaris*, Douglas.

N. B. This muſcle has been ſometimes found wanting on both ſides.

Flexors:

Theſe conſiſt of four; two that belong to the tibia, and two to the fibula.

1. Tibialis Anticus,

Ariſes, tendinous and fleſhy, from the middle of that proceſs of the tibia, to which the fibula is connected above; then it runs down fleſhy on the outſide of the tibia; from which, and the upper part of the interoſſeous ligament, it receives a number of diſtinct fleſhy fibres; near the extremity of the tibia, it ſends off a ſtrong round tendon, which paſſes under part of the ligamentum tarſi annulare near the malleolus internus.

Inſerted, tendinous, into the inſide of the os cuneiforme internum, and poſterior end of the metatarſal bone that ſuſtains the great toe.

Uſe. To bend the foot, by drawing it upwards, and, at the ſame time, to turn the toes inwards.

2. Tibialis Posticus,

Ariſes, by a narrow fleſhy beginning, from the fore and upper part of the tibia, juſt under the proceſs which joins it to the fibula; then paſſing through a perforation in the upper part of the interoſſeous ligament, it continues its origin from the back-part of the fibula next the tibia, and from near one half of the upper-part of the laſt named bone; as alſo, from the interoſſeous ligament, the fibres running towards a middle tendon, which ſends off a round one that paſſes in a groove behind the malleolus internus.

Inſerted,

Inserted, tendinous, into the upper and inner part of the os naviculare, being further continued to the os cuneiforme internum and medium; besides, it gives some tendinous filaments to the os calcis, os cuboides, and to the root of the metatarsal bone that sustains the middle toe.

Use. To extend the foot, and to turn the toes inwards.

3. Peronæus Longus,

Arises, tendinous and fleshy, from the fore-part of the head of the perone, or fibula, the fibres running straight down; also from the upper and external part of the fibula, where it begins to rise into a round edge; as also, from the hollow between that and its anterior edge, as far down as to reach within a hand's breadth of the ankle, by a number of fleshy fibres, which run outwards towards a tendon, that sends off a long round one, which passes through a channel at the outer ankle, in the back-part of the inferior extremity of the fibula; then, being reflected to the sinuosity of the os calcis, it runs along a groove in the os cuboides, above the muscles in the sole of the foot.

Inserted, tendinous, into the outside of the root of the metatarsal bone that sustains the great toe, and by some tendinous fibres in the os cuneiforme internum.

Use. To move the foot outwards, and to extend it a little.

Peroneus maximus, vulgo *Peroneus posterior*, Winslow.

Peroneus primus, seu *Posticus*, Douglas.

4. Peroneus Brevis,

Arises, by an acute fleshy beginning, from above the middle of the external part of the fibula; from the outer side of the anterior spine of this bone; as also, from its round edge externally, the fibres running obliquely outwards towards a tendon on its external side; it sends off a round

round tendon which paſſes through the groove at the outer ankle, being there included under the ſame ligament with that of the preceding muſcle; and a little farther, it runs through a particular one of its own.

Inſerted, tendinous, into the root and external part of the metatarſal bone that ſuſtains the little toe.

Uſe. To aſſiſt the former in pulling the foot outwards, and extending it a little.

Peroneus medius, vulgo *Peroneus anticus*, Winſlow.

Peroneus ſecundus, ſeu *Anticus*, Douglas.

Second Claſs.

Common extenſors.

Theſe conſiſt of two.

1. Extensor Longus Digitorum Pedis,

Ariſes, tendinous and fleſhy, from the upper and outer part of the head of the tibia, and from the head of the fibula where it joins with the tibia, and from the interoſſeous ligament; alſo from the tendinous faſcia, which covers the upper and outſide of the leg by a number of fleſhy fibres; and tendinous and fleſhy, from the anterior ſpine of the fibula, almoſt its whole length, where it is inſeparable from the *peroneus tertius*. It ſplits into four round tendons, under the ligamentum tarſi annulare.

Inſerted, by a flat tendon, into the root of the firſt joint of each of the four ſmall toes; and is expanded over the upper ſide of the toes, as far as the root of the laſt joint.

Uſe. To extend all the joints of the four ſmall toes.

Extenſor longus, Douglas.

N. B. A portion of this muſcle, which

Ariſes,

Ariſes, from the middle of the fibula, continues down to near its inferior extremity, and ſends its fleſhy fibres forwards to a tendon, which paſſes under the annular ligament, and is

Inſerted into the root of the metatarſal bone that ſuſtains the little toe: it is called by Albinus *Peroneus tertius*; and by others, the *Nonus Veſalii*.

Uſe. To aſſiſt in bending the foot.

2. Extensor Brevis Digitorum Pedis,

Ariſes, fleſhy and tendinous, from the fore and upper part of the os calcis; and ſoon forms a fleſhy belly, diviſible into four portions, which ſends off an equal number of tendons that paſs over the upper part of the foot, under the tendons of the former.

Inſertèd, by four ſlender tendons, into the tendinous expanſion from the extenſor longus, which covers the ſmall toes, except the little one; alſo into the tendinous expanſion from the extenſor pollicis that covers the upper part of the great toe.

Uſe. To extend the toes.

Extenſor brevis, Douglas.

Flexors.

Theſe may be reckoned three.

1. Flexor Brevis Digitorum Pedis, Sublimis Perforatus,

Ariſes, by a narrow fleſhy beginning, from the inferior and poſterior part of a protuberance of the os calcis, between the abductors of the great and little toes; ſoon forms a thick fleſhy belly, which ſends off four tendons that ſplit for the paſſage of the flexor longus.

Inſerted

Inserted into the second phalanx of the four lesser toes. The tendon of the little toe is often wanting.

Use. To bend the second joint of the toes.

Perforatus, seu *Sublimis*, Douglas.

2. Flexor Longus Digitorum Pedis, Profundus, Perforans,

Arises, by an acute tendon, which soon becomes fleshy, from the back part of the tibia, some way below its head, near the entry of the medullary artery; which beginning, is continued down the inner edge of this bone by short fleshy fibres, ending in its tendon; also by tendinous and fleshy fibres, from the outer edge of the tibia; and between this double order of fibres the tibialis posticus muscle lies inclosed. Having passed under two annular ligaments, it then passes through a sinuosity at the inside of the os calcis; and, about the middle of the sole of the foot, divides into four tendons, which pass through the slits of the perforatus; and, just before its division, it receives a considerable tendon from that of the flexor pollicis longus.

Inserted into the extremity of the last joint of the four lesser toes.

Use. To bend the last joint of the toes.

This muscle is assisted by the

Flexor Digitorum Accessorius, seu Massa Carnea Jacobii Sylvii,

Arises, by a thin fleshy origin, from most part of the sinuosity at the inside of the os calcis, which is continued forwards, for some space on the same bone; also by a thin tendinous beginning, from before the tuberosity of the os calcis externally; and, soon becoming all fleshy, is

Inserted

Inserted into the tendon of the flexor longus, just at its division into four tendons.

Use. To assist the flexor longus.

3. Lumbricales Pedis,

Arises, by four tendinous and fleshy beginnings, from the tendon of the flexor profundus, just before its division, near the insertion of the massa carnea.

Inserted, by four slender tendons, into the inside of the first joint of the four lesser toes, and are lost in the tendinous expansion that is sent from the extensors to cover the upper part of the toes.

Use. To increase the flexion of the toes, and to draw them inwards.

CHAP. XXXII.

Muscles *which are chiefly situated on the* Foot.

THESE may be divided into the *muscles* of the *great toe*, of the *little toe*, and of the *metatarsal bones*.

Muscles of the great toe.

These are five;

1. Extensor Proprius Pollicis Pedis,

Arises, by an acute, tendinous, and fleshy beginning, somewhat below the head and anterior part of the fibula, along which it runs to near its lower extremity, connected to it by

a number of fleſhy fibres, which deſcend obliquely towards a tendon.

Inſerted, tendinous, into the poſterior part of the firſt and laſt joint of the great toe.

Uſe. To extend the great toe.

Extenſor longus, Douglas.

2. Flexor Longus Pollicis Pedis,

Ariſes, by an acute, tendinous, and fleſhy beginning, from the poſterior part of the fibula, ſome way below its head, being continued down the ſame bone, almoſt to its inferior extremity, by a double order of oblique fleſhy fibres; its tendon paſſes under an annular ligament at the inner ankle.

Inſerted into the laſt joint of the great toe, and generally ſends a ſmall tendon to the os calcis.

Uſe. To bend the laſt joint of this toe.

Flexor longus, Douglas.

3. Flexor Brevis Pollicis Pedis,

Ariſes, tendinous, from the under and fore-part of the os calcis, where it joins with the os cuboides, from the os cuneiforme externum, and is inſeparably united with the abductor and adductor pollicis.

Inſerted into the external os ſeſamoideum and root of the firſt joint of the great toe.

Uſe. To bend the firſt joint.

4. Abductor Pollicis Pedis,

Ariſes, fleſhy, from the inſide of the root of the protuberance of the os calcis, where it forms the heel; and tendinous from the ſame bone, where it joins with the os naviculare.

Inſerted,

Inserted, tendinous, into the internal os sesamoideum, and root of the first joint of the great toe.

Use. To pull the great toe from the rest.

Thenar, Winslow.

5. Adductor Pollicis Pedis,

Arises, by a long thin tendon, from the os calcis, from the os cuboides, from the os cuneiforme externum, and from the root of the metatarsal bone of the second toe.

Inserted into the external os sesamoideum, and root of the metatarsal bone of the great toe.

Use. To bring this toe nearer the rest.

Antithenar, Winslow.

Muscles of the little toe.

These, besides the common extensors and flexors, are two, viz.

1. Abductor Minimi Digiti Pedis,

Arises, tendinous and fleshy, from the semicircular edge of a cavity on the inferior part of the protuberance of the os calcis, and from the root of the metatarsal bone of the little toe.

Inserted into the root of the first joint of the little toe externally.

Use. To draw the little toe outwards from the rest.

Parathenar major, and *Metatarsus*, Winslow.

2. Flexor Brevis Minimi Digiti Pedis,

Arises, tendinous, from the os cuboides, near the sulcus or furrow for lodging the tendon of the peroneus longus; fleshy from the outside of the metatarsal bone that sustains this toe, below its protuberant part.

Inserted into the anterior extremity of the metatarsal bone, and root of the first joint of this toe.

Use. To bend this toe.

Parathenar minor, Winslow.

Muscles from the metatarsal bones.

These are four external and three internal interossei, and one muscle which is common to all the metatarsal bones.

Interossei Pedis externi, Bicipites.

1. Abductor Indicis Pedis,

Arises, tendinous and fleshy, by two origins, from the root of the inside of the metatarsal bone of the fore toe, from the outside of the root of the metatarsal bone of the great toe, and from the os cuneiforme internum.

Inserted, tendinous, into the inside of the root of the first joint of the fore toe.

Use. To pull the fore toe inwards from the rest of the small toes.

2. Adductor Indicis Pedis,

Arises, tendinous and fleshy, from the roots of the metatarsal bones of the fore and second toe.

Inserted, tendinous, into the outside of the root of the first joint of the fore toe.

Use. To pull the fore toe outwards towards the rest.

3. Adductor Medii Digiti Pedis,

Arises, tendinous and fleshy, from the roots of the metatarsal bones of the second and third toes.

Inserted,

Inſerted, tendinous, into the outſide of the root of the firſt joint of the ſecond toe.

Uſe. To pull the ſecond toe outwards.

4. Adductor Tertii Digiti Pedis,

Ariſes, tendinous and fleſhy, from the roots of the metatarſal bones of the third and little toe

Inſerted, tendinous, into the outſide of the root of the firſt joint of the third toe.

Uſe. To pull the third toe outwards.

Interoſſei Pedis interni.

1. Abductor Medii Digiti Pedis,

Ariſes, tendinous and fleſhy, from the inſide of the root of the metatarſal bone of the middle toe internally.

Inſerted, tendinous, into the inſide of the root of the firſt joint of the middle toe.

Uſe. To pull the middle toe inwards.

2. Abductor Tertii Digiti Pedis,

Ariſes, tendinous and fleſhy, from the inſide and inferior part of the root of the metatarſal bone of the third toe.

Inſerted, tendinous, into the inſide of the root of the firſt joint of the third toe inwards.

Uſe. To pull the third toe inwards.

3. Abductor Minimi Digiti Pedis,

Ariſes, tendinous and fleſhy, from the inſide of the root of the metatarſal bone of the little toe.

Inſerted, tendinous, into the inſide of the root of the firſt joint of the little toe.

Uſe. To pull the little toe inwards.

The

The muſcle which brings the extremities of the metatarſal bones towards each other, is named

TRANSVERSALIS PEDIS,

Ariſes, tendinous, from the under part of the anterior extremity of the metatarſal bone of the great toe, and from the internal os ſeſamoideum of the firſt joint, adhering to the adductor pollicis.

Inſerted, tendinous, into the under and outer part of the anterior extremity of the metatarſal bone of the little toe, and ligament of the next toe.

Uſe. To contract the foot, by bringing the great toe and the two outermoſt toes nearer each other.

N. B. The muſcles ſituated on the ſole of the foot are covered by a ſtrong tendinous aponeuroſis, which is extended from the os calcis to the firſt joints of all the toes, and ſerves to preſerve the ſubjacent parts from being compreſſed in ſtanding and walking.

OF

MUSCULAR MOTION.

MOTION in the human body is not performed by one individual organ. In every animal and vegetable fibre, even in hair, feathers, membranes, the cellular texture, and in the humid muſcular fibre, in ſhort, in animal and vegetable gluten, there is a contractile or elaſtic power, which preſerves the fibre in its natural ſtate, preventing it from being too much extended, and reſtoring it again to its proper dimenſions, when the ſtretching power is removed. This power never ceaſes endeavouring to bring the elementary particles into the cloſeſt contact that the mechaniſm of the part can admit. It continues even many days after death; ſo that fibres of a divided muſcle contract towards each extremity, leaving a wide gap in the middle. An artery, when cut, likewiſe contracts itſelf in length.

This force may be called *vis mortua*, becauſe it continues to be efficacious after death, and is different from the powers of life. In the living animal, indeed, it is ſomewhat briſker; for, both from cold and fear, the ſkin trembles, grows harder, and has its papillae erected; and along with this hardneſs it contracts itſelf in length. Again, the cellular fibres are animated with this perpetual niſus to ſhorten, always endeavouring to contract themſelves. Hence, when the ſkin

of

or any other membrane is extended, as ſoon as the cauſe of extenſion is taken off, it returns by a gentle effort to its former ſhortneſs. But it even ſomewhat reſiſts the attempt to perpetual diſtenſion; and by a gentle, but continual, approach of its elements to one another, it propels the contained fat, or water, or other bodies accidentally introduced into it. The ſame power alſo ſeems to limit the excretion of vapour; for the fibres and plates of the cellular texture being preternaturally relaxed, an immenſe quantity, either of fat or of watery humour, is depoſited in that texture. And this debility ſeems to be the principal cauſe of a dropſy. The ſame cauſe being always efficacious, and at work in the heart, joints, and every where throughout the body of the embyro, brings into nearer contact the arteries, auricles, and verticles; produces flexures; and contracts the heart into the form of a cone. The ſame cauſe, by an unknown or hidden power, ſeems to form the ſhape of moſt parts of the human body; and while it expels the gluten received into the cells, it brings the terreſtrial particles nearer to one another, and gives the proper ſolidity, curvature, and ſituation, to the different parts.

It is the nature of this power to act continually by gentle but uninterrupted efforts. It is common for it alſo to be excited by poiſons, in every membrane, fibre, and in the cellular texture; but never by cutting or puncturing with a ſharp inſtrument. Theſe are the known properties of the red muſcular fibre. The ſtructure of this fibre, then, it is now neceſſary for us to conſider.

By the name of *muſcular* or *moving fibres* in the human body, we call bundles of reddiſh coloured threads, which perform all the motions viſible in the human body. When many of theſe fibres are collected together, and appear more evidently red, they are called *a muſcle*. The extreme ſimplicity of the texture in theſe parts has been the cauſe of the obſcurity

obſcurity in underſtanding how a ſmall ſoft portion of fleſh, can produce ſuch ſtrong and ample motions as we ſee in man, and more eſpecially in the cruſtaceous inſects.

In every muſcle we meet with long ſoft threads of *fibres*, ſomewhat elaſtic, or extenſible, and almoſt conſtantly diſpoſed parallel with each other; and theſe, being ſurrounded with a good deal of cellular ſubſtance, are by that faſtened together into little bundles, called *lacertuli*; which are again tied together into larger bundles, by a more looſe cellular net-work, that contains ſome fat; and between theſe we conſtantly perceive membranous partitions and ſtripes of the cellular ſubſtance removing them farther from each other, till at laſt a number of them combined together, either parallel or inclined, are ſurrounded with a more thin and denſe cellular membrane, continuous with that of their partitions; and this being again ſurrounded by a thicker plate of the cellular ſubſtance, parts the whole bundle from the adjacent fleſh, and gives it the denomination of a *ſingle* or *entire muſcle*. In every one of theſe threads there appears a leſſer ſeries of filaments, which, by oblique extremities, are cemented to others of the ſame kind, forming together a large fibre *.

The generality of the muſcles but more eſpecially thoſe which are inſerted into the bones, have other fibres fixed to them; but theſe are condenſed into a more ſlender, hard, and ſhining ſubſtance, of a ſilver colour, which has the name of *tendon*. When the tendinous fibres expand into a broad flat ſurface, they form what is called an *aponeuroſis*.

* Swammerdam, Lyonet, and Roeſel, have elegantly painted the appearance of the muſcular fibres in frogs, and other animals, which Leeuwenhoeck, Cowper, and Muyſs, have ſhewn in man. But their ſtructure is more particularly deſcribed and delineated in a ſmall work by Proſchaſka, *De Carne muſculari*, publiſhed at Vienna 1778.

neurosis. The cellular texture which covers the whole tendon, is called its *vagina* or *sheath*, and resembles the coat of a muscle.

That fleshy fibres truly change into such as are tendinous, says Dr Haller, is evident from comparing a foetus, in which there are few tendons, with a child of some years growth, in which there are many more; and both with an adult, in which there are the greatest number. But Dr Wrisberg observes, in opposition to this, that many tendons are found in a foetus, which could not assume this nature by muscular action; as the tendo Achillis, aponeurosis plantaris, centrum diaphragmatis, &c. Besides, in various parts of the body there are tendons found without corresponding muscles.

Muscles which are not inserted into any of the bones, have commonly no tendons, as the sphincters and muscular membranes of the viscera and vessels. But those commonly end in long tendons, which are required to pass round the joints and heads of the bones, to be inserted in those extremities which are more moveable. In a foetus the muscles are evidently inserted into the periosteum only; but in adults, where the periosteum is more closely joined with the bone itself, the tendons, being confused with the periosteum, pass together with that even into the foveoli of the bone.

The tendinous fibres, indeed, often lie in a straight line with the fleshy ones, and are as it were a continuation of them. But in many parts of the body the fleshy fibres are obliquely inclined to the tendon, and adhere to it; and the tendon itself grows thicker in its progress by continually receiving new fibres. This is called a *tendinous muscle*. Other tendons lie in the middle between two plates of fibres, which are inserted on each side of the tendon at angles obtuse downward; and this is called a *pennated muscle*,

muſcle. There are inſtances of numerous tendons, pennated in different places, formed into one muſcle. There are alſo other methods by which the tendinous fibres are joined with the fleſhy ones.

Within the cellular tunic that ſurrounds the fibres, the arteries and veins, running in company with each other, are ſubdivided into net-works. From the ſmaller of theſe veſſels a vapour is exhaled into the thinner, and the fat is transfuſed into the thicker, cellular ſubſtance; from whence again they are both abſorbed by the lymphatic veſſels; which can be diſtinctly ſeen both on the ſurface and in the ſubſtance of the muſcles.

The nerves of the muſcles are ſtill more evident. They are commonly very large; and enter by ſo many branches, that ſome anatomiſts have conſidered the muſcular fibres as formed by them. Several arguments are now offered againſt this opinion: One of the ſtrongeſt is, that muſcles in the limbs of animals do not ſhrink, although the nerves entering theſe muſcles have been cut through for a conſiderable length of time. (See *Monro's Obſervations on the Nervous Syſtem*). The nerves enter the muſcles in a way ſomewhat ſimilar with that of the arteries and veins; but it is impoſſible to trace them a great way among the fleſhy fibres, for they at length depoſite their harder covering, and become ſoft, and diſappear before they can be traced to their terminations.

The fabric of the leaſt fibres, which are ſuppoſed to be the elementary particles of a muſcle, being inveſtigated by the microſcope in man and other animals, is found to be ſimilar to that of the larger fibres, and they are all joined together by the intermediate cellular ſubſtance. The ſurface of theſe fibres, however, puts on a curious waved or zigzag appearance, as was formerly mentioned. This undulated appearance Proſchaſka thinks is nothing elſe than

impreſſions made by the veſſels and cellular ſubſtance and perhaps by the nerves: But Dr *Monro*, in the work above quoted, has deſcribed and delineated a ſimilar appearance in the tendons and nerves; and is of opinion, that they are to be conſidered as folds or joints, ſerving to accommodate the parts to the different ſtates of flexion and extenſion. In proof of this, he finds, that thoſe parts which appear thus in their relaxed ſtate, loſe it when they are much ſtretched.

With regard to the nature of the ultimate moving fibres, there have been many diſputes. Some anatomiſts think them ſolid; others, hollow, formed of a ſeries of veſſels or rhomboidal chains communicating with each other: Some, again, have been of opinion, that they are full of a kind of down or woolly ſubſtance, &c.

It may be aſked, Whether they are hollow? whether they are continued from the arteries? or whether the difference between muſcular and tendinous fibres, lies in the latter being rendered more denſe, and beat cloſſer together by an expulſion of the fluids? That theſe are not probable, appears from the minuteneſs of the fibres, which are found leſs than the red-blood globules, and from the whiteneſs of a muſcle after the blood is waſhed out of it; to which add the phyſiological reaſons, after mentioned, (p. 385:) And, in general, more ſtrength may be expected from a ſolid fibre.

The ſtructure of the tendons and aponeuroſes agrees in ſome reſpects with, but differs in others from, that of the muſcles. We obſerve their fibres regularly diſpoſed, and ſeparated by cellular ſubſtance and blood-veſſels; and without doubt they have lymphatics and nerves: But the tendinous fibres are cloſſer together than thoſe of the muſcles, the cellular ſubſtance which ſeparates them is finer, their red veſſels are fewer in number, and the nerves cannot be traced, without difficulty, into their ſubſtance.

It

It has been doubted whether the fibres of tendons are a continuation of the moving fibres, or of a different nature. Many, both ancients and moderns, have embraced the first opinion, others the second; but if we consider that the tendinous fibres are not irritable, have no contraction, that they differ little from those which constitute the ligaments, and that they degenerate sometimes into a substance truly cellular, we would be inclined to adopt the latter opinion.

According to Dr Haller, a muscle therefore is endowed at least with a threefold force. First the *vis mortua*, in common to it with other animal fibres. Another, which he has called the *vis insita*, and which has different phenomena from the former *. It is more peculiar to life; and though it may continue for a few hours after death, yet it disappears much sooner than the former. Again, in most cases, it acts by alternate oscillations; so that, being driven hither and thither, it sometimes contracts the muscle towards the middle; sometimes again it extends the muscle from the middle towards the extremities, and sometimes also it has a reiterated motion. Moreover, it is manifestly quicker, and performs the greatest motions *. It is excited both by the pricking with a sharp instrument, and in the hollow muscles by inflated air, by water, and every kind of acrimony, but most powerfully of all by a torrent of electrical matter. Lastly, it is peculiar to the muscular fibre, and is found in no other part of the human body with the qualities above mentioned. But we must give a more particular explication of its phenomena.

It is natural to every muscle to shorten itself, by drawing the extremities towards its belly or middle. But to discover the moving power of a muscle from the fabric which we have described, it will be of use to consider the appearances observable in the muscular contraction. Every muscle then becomes

* See observations on this subject in the subsequent pages of this Chapter.

becomes ſhorter and broader in its action. But this contraction of its length is various; in ſome more, in others leſs; and is very conſiderable in ſome of the ſphincters, inſomuch that they appear to be contracted more than one third of their extent.

The inteſtines are exceedingly tenacious of their vis inſita, as they continue to contract themſelves long after they are taken out of the body, and even after they are cold; and the heart is even more tenacious of this principle than the inteſtines, as is evident in a young chicken, and in cold animals. Different muſcles are beſt excited into action by different ſtimuli; as the bladder by urine, the heart by the blood, and the inteſtines by air. Though the nerves are removed, or the connection, made by them between the muſcles and the brain, cut off, the muſcles loſe but little of their irritable nature. It appears alſo from the example of polypi and other inſects, that the ſame irritable diſpoſition extends very widely through the animal fibres. In many of theſe creatures, neither brain nor nerves can be traced, yet they are exceedingly affected by ſtimulus. Laſtly, we perceive this vis inſita in vegetables, many of which expand and contract their flowers and leaves according to the different ſtates of the atmoſphere; and ſome plants exert ſudden and violent motions from ſtimuli externally applied, appearing little inferior in this reſpect to animals. This vis inſita does not depend upon weight, attraction, or elaſticity; for it is ſeated in a ſoft fibre, and vaniſhes as the fibre grows hard.

That the cauſe of motion is conveyed through the nerves into the muſcles, is certain from the experiments made on the brain and nerves. For the nerve alone has feeling; this alone carries the commands of the ſoul; and of theſe commands there is neither intimation nor perception in that part, whoſe nerve is either tied or cut, or which has no nerve.

nerve. Moreover, on irritating the nerve or ſpinal marrow, even in an animal that has been dead for ſome time, the muſcle or muſcles which have nervous branches from thoſe parts are moſt vehemently convulſed. When the nerve of any muſcle is cut or tied, or the baſis of the ſpinal marrow compreſſed, or that part of the brain from whence the nerve has its origin, the muſcle becomes paralytic, and languiſhes, and cannot by any force be recalled into an action ſimilar to the vital one. But if the ligature be taken off from the nerve, the force by which the muſcle is put into action is again recovered. The nerve being irritated below the place where it is cut, the muſcle to which that nerve goes is contracted. This appears from numerous experiments, eſpecially thoſe made on the phrenic and recurrent nerves.

This force, called *vis nervea*, is not the ſame with the vis inſita. The former comes to the muſcle from without: whereas the other reſides conſtantly in the muſcle itſelf. The nervous power ceaſes when life is deſtroyed; the other appears, from certain experiments, to remain for ſome time after death: it is alſo ſuppreſſed by tying a ligature upon the nerve, by hurting the brain, or by taking opium. The *vis inſita* ſuffers nothing from all theſe circumſtances; it remains after the nerve is tied: it continues in the inteſtines though they be taken out of the body and cut in pieces: it appears with great ſtrength in ſuch animals as are deſtitute of brain: that part of the body is moved which has no feeling; and the parts of the body feel which are without motion. The will excites and removes the nervous power, but has no power over the *vis inſita*.

In the motion of the muſcles, whether owing to the *vis inſita* or the *vis nervea*, the fibres are contracted towards the middle of their belly, and recede from one another outwards; they are alſo diverſified with various tranſverſe wrinkles:

wrinkles: the whole muſcle alſo becomes ſhorter, and draws its extremities towards the middle; hence it draws towards each other thoſe parts with which it was connected: the muſcle alſo ſwells by its contraction, becoming hard at the ſame time, and as it were increaſes its circumference on every ſide. Haller has never obſerved it to turn pale. Whether on the whole it is increaſed in bulk, or acquires more in breadth than it loſes in length, is difficult to be known. It draws after it the tendons, which are obſequious to its motions, though of themſelves neither moveable nor irritable. The whole muſcle is alſo capable of being moved at once, or only a part of it; if one extremity is fixed to an immoveable part, that only is moved which is capable of yielding.

If it be demanded, Whether the arteries conduce any thing to muſcular motion? and whether the palſy of the lower limbs after a ligature upon the aorta, be not an argument of this opinion? we anſwer, No farther than as they conduce to the ſound ſtate of the muſcle, or to the conſervation of the mutual ſtructure and habit of the parts, which they ſupply with vapour, fat, &c.: for the irritation of an artery does not affect or convulſe the muſcle to which it belongs; nor does a ligature on it cauſe a palſy, unleſs after a conſiderable time, when the muſcles begin to be deſtroyed by a gangrene. Moreover, it is impracticable to explain the motion of peculiar muſcles from a cauſe derived with an equal force from the heart to all parts of the body. Laſtly, the influence of the will is confined to the nerves, without reſiding in the arteries or other ſolid parts of the body.

But the direct manner by which the nerves excite motion in the muſcles, is ſo obſcure, that we may almoſt for ever deſpair of its diſcovery. The opinion, that the nervous veſicles ſwell by a quicker flux of the nervous ſpirits, is inconſiſtent

conſiſtent with anatomical truth, which demonſtrates the leaſt viſible fibres to be cylindrical, and in no part veſicular; it is likewiſe repugnant to the celerity with which muſcular motion is performed; and it ſeems to be completely refuted by the well known fact, that the bulk of a muſcle is rather diminiſhed than increaſed during its action. Again, the inflation of rhomboidal chains in the fibres is equally repugnant to the celerity of muſcular motion, and to anatomy; they would alſo occaſion an immenſe waſte of ſtrength, and after all render the muſcle but little ſhorter. The nerves want that irritable nature which is obſerved in the muſcular fibre. Finally, it is by no means demonſtrable, that the fibres, from ſo few nerves, can be ſo numerous, or diſtributed in ſo many different tranſverſe directions, with reſpect to the muſcular threads, as thoſe hypotheſes require to be allowed. A complication of the nerves round the extremities of fibres, ſo as to contract them by their elaſticity, is founded upon a falſe ſtructure of the muſcular fibre, ſuppoſing the nerves to be diſtributed, where filaments of the cellular ſubſtance only can be demonſtrated. Moreover, the phenomena of animals, which, having neither brain nor nerves, are yet very apt for motion, apparently demonſtrate the intrinſic fabric of the muſcles to be ſufficient for their motion, without other aſſiſtance from the nerves. Other explanations, derived from ſpherules of air in the blood, are founded on a falſe hypotheſis, for the blood contains no elaſtic air. (See article *Reſpiration*). The animal ſpirits are not of the nature of an electric torrent.

If we may add any thing to theſe phenomena, we may ſuppoſe the nervous fluid to be of a ſtimulating nature, by which means it forces the elementary particles of the muſcular fibre to approach nearer to each other. The motive cauſe which occaſions the influx of the animal ſpirits into the

the muſcle ſo as to excite it into action, ſeems not to be the ſoul, but a law derived immediately from God. For animals newly born, or newly transformed, without any attempt, or exerciſe, know how to execute compound motions, very difficultly to be defined by calculation; whereas the ſoul knows not how to perform any actions until ſhe has learnt, by making experiments, the methods of performing them. A muſcle then is contracted when in a given time it receives more of the nervous fluid, whether that be occaſioned by the will, or by ſome irritating cauſe ariſing in the brain, or applied to the nerve.

But though we may ſuppoſe the ſoul to be the cauſe of the nervous motion, we cannot do the ſame with regard to that ariſing from the vis inſita. The heart and inteſtines, and alſo the organs of generation, are governed by a vis inſita, and by ſtimuli. Theſe powers do not ariſe from the will; nor are they leſſened, or excited, or ſuppreſſed, or changed by it. No cuſtom nor art can make theſe organs ſubject to the will, which have their motions from a vis inſita; nor can they be made to obey the commands of the ſoul, like attendants on voluntary motion. It is ſo certain that motion is produced by the body alone, that we cannot even ſuſpect any motion to ariſe from a ſpiritual cauſe, beſides that which we ſee is occaſioned by the will; and, even in that motion which is occaſioned by the will, a ſtimulus will occaſion the greateſt exertions, when the mind is very unwilling.

There ſeems to be this difference between the muſcles obeying the will, and thoſe which are governed by a vis inſita; namely, that the latter, being more irritable, are very eaſily excited into motion by a gentle ſtimulus; as for inſtance, the heart and inteſtines; which organs are moſt manifeſtly, and greatly, and conſtantly, irritable. On the other hand, the muſcles, which obey the will, are neither endowed

endowed with ſo great nor ſo durable a power of this kind. Hence they either ſtand in need of the power of the will, or of a ſtronger ſtimulus; when they are excited by a ſtimulus, they are animated to motion againſt the will. Thus it happens, that in apoplexies the muſcles which obey the will languiſh, and become paralytic, as being deſtitute of all influx from the brain; while the vital muſcles, having no occaſion for the operation of the brain, continue to be excited into contraction by their ſtimuli, independent of the will, as the heart by the blood, and the inteſtines by the air and aliments.

The ſtrength of the muſcles is very conſiderable in all perſons, but more eſpecially phrenetic and robuſt men; ſince frequently, with the uſe of a few muſcles only, they will raiſe a weight equal to, or much greater than, that of the whole human body. In a healthy man, very ſlender muſcles ſuffice to lift 200 or 300 pounds. The muſcles of the back will even ſuſtain 3000. Notwithſtanding this, we ſee, that much the greater part of the force or power exerted by a muſcle is always loſt without producing any viſible effect. For all the muſcles are inſerted nearer the point or centre of motion, than the weights they are applied to; and therefore their action is weaker, in the ſame proportion as they move a ſhorter part of the lever than that to which the weight is applied. Moreover, in moſt of the bones, eſpecially thoſe of the limbs, the muſcles are inſerted at very acute angles; whence again the effect which a muſcle exerts in action is proportionally leſs as the ſine of the angle intercepted between the bone and the muſcle is leſs than the whole ſine. Again, the half of all muſcular force is loſt, becauſe a muſcle, like an extended cord, exerts as much force at its fixed as at its moving extremity. Beſides, many of the muſcles are ſeated in the

angle of two bones; and when the joint is bent, the muſcle becomes alſo bent; conſequently a quantity of force muſt be firſt applied to ſtretch the muſcle, before the muſcle can act on the bone in which it is inſerted. Many of them paſs over ſeveral joints, each of which they bend in ſome degree, whereby a leſs part of their remaining force goes to bend the joint to which they are particularly deſtined. The fleſhy fibres themſelves of the muſcles frequently intercept angles with the tendon in which they terminate; whence their force is diminiſhed in proportion to the ſine of the angle of their inſertion. Finally, the muſcles move their oppoſed weights with the greateſt velocity and expedition, ſo as not only to overcome the equilibrium, but likewiſe to add a conſiderable celerity to the weight.

All theſe loſſes of power being computed, make it evident, that the force exerted by muſcles in their contraction, is exceeding great, and almoſt beyond any mechanical computation. The effect is ſcarce one-ſixtieth of the whole force exerted by the muſcle, and yet only a ſmall number of theſe muſcles, weighing but a few pounds, are able not only to raiſe ſome thouſands of pounds, but alſo with a conſiderable celerity. Nor is this to be reputed any defect of wiſdom in the Creator: For all thoſe loſſes of power were neceſſary for a juſt ſymmetry or proportion of the parts, and for various motions and celerities in different directions. But we may, however, conclude from hence, that the action of the nervous or animal fluid is very powerful, ſince, in an engine ſo ſmall, it can exert a force equal to ſome thouſand pounds for a conſiderable time, even for many days together; nor does this ſeem to be otherwiſe explicable, than by the incredible celerity by which the influx of this fluid obeys the command of the will. But how or whence it acquires ſuch a velocity, is not in our power to ſay; it is ſufficient that we know the laws of its motion are ſuch, that a given action of

the will produces a new and determinate celerity in the nervous fluid.

There are various means by which the motions of the muſcles are rendered more ſafe, certain, and eaſy. The large long muſcles, by which the greater motions of flexure are performed, being included in tendinous capſules or caſes, drawn and tightened by other muſcles, are thus ſecured and ſtrengthened; and thus the muſcle, in a ſtate of contraction, remains preſſed againſt the bone all the time that the limb is bent, and conſequently avoids a conſiderable loſs of its power. The long tendons, which are incurvated or extended over joints in their motion, are received and confined by peculiar ligaments, which retain them within their ſlippery channels; and in theſe ſheaths a particular liquor is ſeparated for the lubricating of the tendons. Nature makes a ſimilar proviſion for thoſe muſcles which perforate others in their courſe. In other parts, the tendons are either carried round certain eminences of the bone, in order that they may be inſerted at greater angles into the bone which they move; or elſe they are inſerted into another bone, from whence a different tendon deſcends under a larger angle into the bone to be moved. In other parts, the muſcles which are derived from convenient ſituations, have their tendons carried round in a contrary direction by nature, ſo that they paſs into the part to be moved as it were round a pulley. Nature has likewiſe ſurrounded the muſcles on all ſides with a conſiderable quantity of fat, which is ſpread alſo between their bundles of fibres, and alſo between the ſmall fibres themſelves: Which fat, being preſſed out by the turgeſcence of the muſcles and fibres, renders them ſoft, flexible, ſlippery, and fit for motion.

Moreover, the power and action of one muſcle is determined by the co operations or oppoſitions of others, which

ſerve

ſerve either to hold firm ſome part from whence the muſcle ariſes, or to bend it together with the muſcle, or elſe to change its action from the perpendicular to the diagonal, by concurring to aſſiſt its force at the ſame time. Remote muſcles often aſſiſt each other, the ſuperior muſcles keeping the bone, whence the inferior acting muſcles ariſe, firm and ſteady. Therefore, the action of no one muſcle can be underſtood from conſidering it alone; but all the others, which are either inſerted into the muſcle itſelf, or into any of the parts to which the muſcle adheres, muſt likewiſe be brought into the conſideration.

By theſe muſcles, variouſly conſpiring and oppoſing each other, are performed walking, ſtanding, flexion, extenſion, deglutition, and all the other geſtures and offices of the ſeveral parts in the living body. But the muſcles have likewiſe ſome common actions, by which they are of uſe to the whole animal. They haſten the return of the venous blood, by preſſing it out both from the veins of the muſcles themſelves, and from the veins which lie between them; for the blood in theſe veſſels, diſtributed between the turgid bundles of a contracted muſcle, is, by the valves, determined towards the heart: They likewiſe return the fat to the blood; and ſhake, grind, or condenſe the arterial blood, and return it quicker to the lungs; they contribute in a great degree to the ſecretions and excretions: Again, in the liver, meſentery, womb, &c. they promote the courſe of the contained blood, bile, and other juices, ſo as to leſſen the danger of their ſtagnation: They ſerve alſo to increaſe the ſtrength of the ſtomach, by adding their own ſtrength to it, whereby digeſtion is promoted; inſomuch that all ſedentary and inactive courſes of life are contrary to nature, and pave the way to diſeaſes from a ſtagnation of the humours, or from a corruption or crudity of the aliments. The large muſcles, which are

generally

generally placed round any of the cavities of the body, propel the blood contained in that cavity, and preſs it towards the heart. By too much exerciſe or action, the muſcles themſelves grow hard and tendinous on all ſides; they render the parts upon which they are incumbent cartilaginous, or elſe change thoſe which are membranous into a bony nature; they increaſe the roughneſs, protuberances, and proceſſes, of the bones which lie next to them, and excavate their flat parts; they dilate the cells of the diploe, and incurvate the bones.

The muſcle which the ſtimulus has ceaſed to irritate, or for the action of which the mind has no occaſion, is relaxed, and grows ſoft; its wrinkles grow plain; its fibres are rendered longer, receding from the middle towards the fixed extremities: and its ſwelling falls. Whatever is the cauſe of additional contraction is then taken away; but the vis inſita remains. It may be aſked, What becomes of the ſpirit that is ſent from the brain? A part of it perhaps exhales; a part of it may be ſuſpected to adhere to the fibre; and thus it happens that the muſcles grow ſtrong with exerciſe, and their brawny parts become thicker.

"The three noted claſſes of animal powers, *elaſticity*, *irritability*, and *ſenſibility*, Dr Wriſberg obſerves, have been and are yet too much confounded, although it is no difficult taſk to diſtinguiſh theſe affections from one another. The elaſticity and cohetion of the fibres, which in different degrees, are found in every part of the body, was fully known to Bellini, Bagliui, Stahl, Pacchioni, Juncker, &c. That power, known to Stahl's followers under the appellation of *tone*, has no ſimilarity to irritability, ſenſibility, and what is called vital power. It either alone performs the various actions of the animal and vegetable body, or adds ſtrength and vigour to them: The former is manifeſt in the motion of the ribs

and

and cartilages; and the latter in the conſtruction of the uterus, veſſels, and membranes. It by no means depends on the vital powers, but may endure long after death; it is not completely deſtroyed even by putrefaction. During life it is diminiſhed by various cauſes, and again reſtored by ſeveral remedies. Irritability, which Haller thought exiſted in the fibres of the muſcles alone, and which indeed was known by name, but not in reality to Gliſſon, is a kind of animal power, unknown to earlier anatomiſts, and is different from that power which Hippocrates calls ενορμουν. It is proved, from the experiments of Lups, Haller, Fontana, Hoffman, and ſeveral others, to differ from elaſticity in its riſe, duration, ſeat, cauſes, effects, and phenomena. We ſhall add a few remarks:

1. It is moſt powerful in the muſcular fibres of the whole body, but not equally diſperſed through all; more powerful in the heart, muſcles of reſpiration, and inteſtines, but becomes gradually weaker among the voluntary muſcles; and it exiſts perhaps, in a ſmall degree, in the veſſels and membranes, as appears from the doubts offered by Whyte, De Haen, Van Doeveren, &c. which Haller and the learned Cigna have anſwered.

2. The phenomena of irritability, and the irritations themſelves by which theſe phenomena are produced, are not always the ſame: In ſome parts we conſtantly perceive a manifeſt irritability produced by every irritation, as is the caſe in almoſt all the muſcles. In many other parts you ſee the greateſt inconſtancy and very irregular effects, varying differently, at different times, being ſometimes increaſed and ſometimes diminiſhed, at one time yielding to, and at another reſiſting the irritations: All which is evident in the ſkin, viſcera, veſſels, and iris.

3. The learned pathologiſts Eller, Tiſſot, and Gerhard, have long ago acknowledged the great uſe of the doctrine of

of irritability. It would be of much importance to know the remedies, which particularly conduce to excite irritability, when it is languid, or to diminish it when it is too great. Opium, and the other narcotics, camphor, cantharides, acrid poisons, bark, the electric shock shew a clear influx of animal spirits in the production of irritability.

4. That it is different from the sentient faculty, and therefore by no means depends upon the nerves, appears partly from other reasons, and partly from the irritability of vegetables. Though I even wish to take into account some phenomena of the dionaea muscipula, according to Ellis's observations, or of the hedysarum or the antherae of certain other plants, I would be averse to compare this contractile power of some parts of vegetables with irritability; for the internal tremor of the constituent parts, which makes the particular character of irritability, is wanting in all vegetables; we see contraction and motion alone, which are also observable in other elastic bodies, where we suppose no irritability to exist. The sentient faculty, depending solely upon the nerves, although it has been regarded as one and the same thing with irritability, has been more strongly opposed by Haller's opponents, De Haen, Whyte, Le Cat, Gerhard, &c. than irritability itself. The sensibility of the parts is to be referred both to the various quantity of the nerves, their situation and state, according to Haller's and Castell's experiments, and to the various violence of irritation, and the nature of the irritating or offending body; for some parts are frequently much affected at one time and less at another, and Haller thinks that some of them may sometimes be altogether insensible. I shall not repeat what has been often offered on the opinion, that a greater pain having preceded, absorbs a less pain following; thus we do not feel the taste of a drop of wine if we have taken a very small quantity of rectified alcohol upon the tongue a little before. It cannot however be denied, that in

inflammatory

inflammatory diſeaſes, affections of the mind, and other cauſes, it may happen, that hurt parts may feel, which, under any other condition, ſeem inſenſible. The *vital power* of certain learned men of later times, as Vanden Bos, Bikker, Gaubius, Albinus, &c. ſeems rather compounded of all the animal powers comprehended together; which opinion, except in ſome minutiae, the great Boerhaave and Simpſon have more exactly adopted."

As the doctrine delivered above, concerning the exiſtence of a *vis inſita* different from the *vis nervea*, has been the cauſe of conſiderable debate, and is at preſent called in queſtion by ſeveral anatomiſts, particularly by Dr Monro, we think it neceſſary to give a few objections as ſtated in his Obſervations on the Nervous Syſtem. The chief experiment, ſays the Doctor, which ſeems to have led Dr Haller to this opinion, is the well known one, that the heart and other muſcles, after being detached from the brain, continue to act ſpontaneouſly, or by ſtimuli may be rouſed into action for a conſiderable length of time; and when it cannot be alleged, ſays Dr Haller, that the nervous fluid is by the mind or otherwiſe impelled into the muſcle.

That in this inſtance, we cannot comprehend by what power the nervous fluid or energy can be put in motion, muſt, perhaps, be granted; But has Dr Haller given a better explanation of the manner in which his ſuppoſed vis inſita becomes active?

If it be as difficult to point out the cauſe of the action of the vis inſita as that of the action of the vis nervea, the admiſſion of that new power, inſtead of relieving, would add to our perplexity.

We ſhould then have admitted, that two cauſes, of a different nature, were capable of producing exactly the ſame effect;

effect; which is not in general agreeable to the laws of nature.

We ſhould find other conſequences ariſe from ſuch an hypotheſis, which tend to weaken the credibility of it. For inſtance, if in a ſound animal the vis nervea alone produces the contraction of the muſcles, we will aſk, what purpoſe the vis inſita ſerves? If both operate, are we to ſuppoſe that the vis nervea, impelled by the mind or living principle, gives the order, which the vis inſita executes; and that the nerves are the internuntii; and ſo admit two wiſe agents employed in every the moſt ſimple action? But inſtead of ſpeculating farther, let us learn the effects of experiments, and endeavour from theſe to draw plain concluſions.

1. When I poured a ſolution of opium in water under the ſkin of the leg of a frog, the muſcles, to the ſurface of which it was applied, were very ſoon deprived of the power of contraction. In like manner, when I poured this ſolution into the cavity of the heart, by opening the vena cava, the heart was almoſt inſtantly deprived of its power of motion, whether the experiment was performed on it fixed in its place, or cut out of the body.

2. I opened the thorax of a living frog, and then tied or cut its aorta, ſo as to put a ſtop to the circulation of its blood. I then opened the vena cava, and poured the ſolution of opium into the heart; and found not only that this organ was inſtantly deprived of its powers of action, but that in a few minutes the moſt diſtant muſcles of the limbs were extremely weakened. Yet this weakneſs was not owing to the want of circulation; for the frog could jump about for more than an hour after the heart was cut out.

In the firſt of theſe two experiments, we obſerve the ſuppoſed vis inſita deſtroyed by the opium; in the latter, the vis nervea; for it is evident that the limbs were affected by the

ſympathy of the brain, and of the nervous ſyſtem in general, with the nerves of the heart.

3. When the nerve of any muſcle is firſt divided by a tranſverſe ſection, and then burnt with a hot iron, or punctured with a needle, the muſcle in which it terminates contracts violently, exactly in the ſame manner as when the irritation is applied to the fibres of the muſcle. But when the hot iron, or needle, is confined to the nerve, Dr Haller himſelf muſt have admitted, that the vis nervea, and not the vis inſita, was excited. But here I would aſk two queſtions.

Firſt, Whether we do not as well underſtand how the vis nervea is excited when irritation is applied to the muſcle as when it is applied to the trunk of the nerve, the impelling power of the mind ſeeming to be equally wanting in both caſes?

Secondly, If it appears that irritation applied to the trunk of a nerve excites the vis nervea, why ſhould we doubt that it can equally well excite it when applied to the ſmall and very ſenſible branches and terminations of the nerve in the muſcle?

As, therefore, it appears that the ſuppoſed vis inſita is deſtroyed or excited by the ſame means as the vis nervea; nay, that when, by the application of opium to the heart of a frog, after the aorta is cut, and the circulation interrupted, we have deſtroyed the vis inſita, the vis nervea is ſo much extinguiſhed, that the animal cannot act with the diſtant muſcles of the limbs; and that theſe afterward grow very torpid, or loſe much of their ſuppoſed vis inſita; it ſeems clearly to follow, that there is no juſt ground for ſuppoſing that any other principle produces the contraction of a muſcle.

A

SYSTEM OF ANATOMY,

WITH THE

PHYSIOLOGY.

PART III.

CONTAINING

EIGHT ANATOMICAL TABLES

OF THE

HUMAN BODY;

Exhibiting the Principal Parts of the

SKELETON AND MUSCLES

REPRESENTED IN THE

LARGE TABLES OF ALBINUS.

TO WHICH ARE ADDED

CONCISE EXPLANATIONS.

By JOHN INNES.

WITH THE ADDITION OF

TWO TABLES AND EXPLANATIONS,

EXHIBITING

The differences between the Male and Female Bones of the Pelvis.

TO THE

MEDICAL SOCIETY.

GENTLEMEN,

I EMBRACE this opportunity of expreſſing my gratitude for the many favours I have received from you, both in your collective and individual capacity; but I am unwilling to offend either by encomiums on the utility of your inſtitution, or by exhibiting the private friendſhips with which I have been honoured by many of your number.

IN compliance with your ſolicitations, I lately publiſhed a ſhort Deſcription of the Human Muſcles, which has been fortunate enough to receive your approbation. It was, however, your opinion, that a ſet of Tables would render the deſcriptions ſtill more perfect and uſeful. I have, therefore, cauſed the following plates to be engraved; and I hope they will not altogether diſappoint your expectations.

THE Tables of Albinus, though accurate and complete, are not, perhaps, on account of their ſize, the prolixity of the deſcriptions, and the number of references, ſo fully ſuited to the purpoſes of private diſſection as could be wiſhed.

To remedy this inconvenience, I have copied eight of Albinus's tables on a ſmaller ſcale. Two of them contain a fore and back view of the ſkeleton; and the other ſix are repreſentations of the muſcles.

To each plate I have prefixed an explanation; and I have avoided, as much as was conſiſtent with perſpicuity, loading the figures with an unneceſſary quantity of referring letters. For this purpoſe, I have all along made uſe of one alphabet only.

To prevent the defacing of the muſcles, and to enable the learner to diſtinguiſh them without the aſſiſtance of references, the figures are only lettered on one ſide.

IF this little work be of uſe in facilitating your anatomical ſtudies, it will give the greateſt pleaſure to,

GENTLEMEN,

Your very much obliged humble ſervant, and Fellow-member,

JOHN INNES.

TAB. I.

D Lazare Sculp.

EXPLANATIONS

OF THE

SKELETON AND MUSCLES,

AS REPRESENTED IN THE

TABLES.

TABLE I.

THIS figure repreſents a front view of the human ſkeleton, with ſome of the ligaments and cartilages, which connect the bones to each other.

HEAD AND NECK.

a, Os frontis.

b, Os parietale.

Between *a* and *b*, part of the coronal ſuture.

c, The pars ſquamoſa of the temporal bone.

Between *b* and *c*, the ſquamous ſuture.

Below the pars ſquamoſa, the zygoma; and, lower down above *f*, the maſtoid proceſs.

Between the pars ſquamoſa and the cavity, which contains the eye-ball, called *orbit*, the temporal proceſs of the ſphenoid bone is ſeen.

d, Os

d, Os malae.

Above *d*, a portion of the tranſverſe ſuture.

e, Os maxillare ſuperius, with the eight teeth of the right ſide.

The naſal proceſs of the ſuperior maxillary bone has the os naſi joined, by the lateral naſal ſuture, to its inſide; and at the outſide, within the orbit, the os unguis.

The oſſa naſi joined to each other before, by the anterior naſal ſuture.

f, Os maxillare inferius with ſixteen teeth; the four anterior named *inciſores*, the two corner ones *canini*, and the five poſterior on each ſide, *molares*.

Oppoſite to *f*, the angle of the lower jaw; above *f*, the condyloid proceſs, by which the jaw is connected to the temporal bone, at the root of the zygoma; and behind the os malae, the coronoid proceſs.

g, The ſeven cervical vertebrae, with their intermediate cartilages.

Oppoſite to *g*, their tranſverſe proceſſes.

TRUNK.

a, Sternum.

a, Its middle piece, to which one half of the cartilage that connects the ſecond rib, the whole of the cartilages of the third, fourth, fifth, ſixth, and one half of the ſeventh, are fixed.

Above *a*, the firſt or upper triangular piece, to which the clavicle and one half of the cartilage that connects the ſecond rib are fixed.

Below *a*, the extremity, or third piece of the ſternum, named *cartilago enſiformis*, to which one half of the cartilage that connects the ſeventh rib is fixed.

b, The ſeventh or laſt true rib.

c, The

c, The twelfth, or laſt of the five falſe ribs.

d, The five lumbar vertebrae, with their intermediate cartilages.

Oppoſite to *d*, their tranſverſe proceſſes.

e, The os ſacrum, with its five diviſions.

f, Os innominatum, divided into

g, Os ilium,

h, Os pubis,

i, Os iſchium.

Oppoſite to *i*, the foramen thyroideum.

SUPERIOR EXTREMITY.

a, The clavicle fixed before, to the firſt piece of the ſternum, and outwards to the acromion of the ſcapula.

b, The ſcapula.

Above *b*, the cervix of the ſcapula.

Oppoſite to it, the inferior coſta; and below the outward extremity of the clavicle, the ſuperior coſta, and coracoid proceſs are ſeen.

c, The os humeri.

The upper end of it, which is connected to the cavity of the ſcapula, named *glenoid*, below the acromion, is named its *head* or *ball*; on each ſide of which is ſeen a tubercle, named *external* and *internal*; and between theſe, a groove for lodging the long head of the biceps flexor cubiti.

d, The internal condyle.

e, The external condyle.

Between *d* and *e*, the trochlea, upon which the ulna moves.

f, The radius.

The upper end, which moves on the external condyle of the os humeri, is named its *head*; below that, the tubercle for

the

the insertion of the biceps flexor cubiti, and between these the cervix.

The inferior end of it is connected to the carpus.

g, Ulna.

The upper end of it forms the coronoid process for the insertion of the brachialis muscle.

The inferior end has a process named *styloid*, which is connected to the carpus by a ligament.

h, The carpus, formed of eight bones.

i, Metacarpal bone of the thumb.

k, The metacarpal bones of the four fingers.

l, The two joints of the thumb.

m, The three joints or phalanges of the fore-finger; and the same are seen in each of the other three.

INFERIOR EXTREMITY.

a, Os femoris.

The upper end of it is named its *head* or *ball*, which is lodged in a deep socket of the os innominatum, named *acetabulum*.

Between the head and trochanter major, the cervix.

b, Trochanter major.

c, Trochanter minor.

d, Internal condyle.

e, External condyle.

f, Patella.

The place where it moves upon the os femoris, is named *trochlea*.

g, Tibia.

Between the tibia and the condyles of the os femoris, the semilunar cartilages are seen; and below the joint, the tubercle of the tibia.

h, Fibula.

i, Malleolus internus.

k, Malleolus

TAB. II.

D.Lizare Sculp.

k, Malleolus externus.

l, Os calcis.

Between *l* and *m*, the other ſix bones of the tarſus.

m, Metatarſal bones of the four toes.

n, The three joints, or phalanges of the four toes.

o, Metatarſal bone of the great toe.

p, The two joints of the great toe.

TABLE II.

THIS Table repreſents the firſt Layer of Muſcles ſituated on the anterior part of the whole body, immediately under the common Integuments and tendinous Faſciae.

MUSCLES *ſituated on the* HEAD *and* NECK.

a, The anterior fleſhy belly of the occipito-frontalis ſituated on the os frontis.

Above *a*, the tendinous oponeuroſis of the occipito-frontalis, covering the upper part of the parietal bones.

b, Attollens aurem.

Under it, the tendinous aponeuroſis covering the temporal muſcle.

Anterior auris between *c* and the ear.

c, Orbicularis palbebrarum.

Its tendon is ſeen at the inner canthus, fixed to the naſal proceſs of the ſuperior maxillary bone.

Levator labii ſuperioris alaeque naſi.

Seen divided into two portions running down along the ſide of the noſe; and on the outſide of it, the levator anguli oris.

Next this, the

Zygomaticus minor.

Farther outwards,

Zygomaticus major.

On the ala and tip of the nose, the

Compressor naris.

d, Depressor anguli oris.

And beneath it, a portion of the depressor labii inferioris.

e, Orbicularis oris.

f, Platysma-myoides.

Behind *f*, the sterno-cleido-mastoidaeus is seen through the platysma-myoides.

TRUNK.

a, Pectoralis major.

The upper part of it is covered by the origin of the platysma-myoides.

b, Serratus magnus.

The other portions resemble this.

c, Latissimus dorsi.

d, Obliquus externus descendens.

e, Linea semilunaris.

f, Linea alba.

Below *f*, umbilicus.

Between *e* and *f*, the rectus abdominis; and, at the inferior part of the linea alba, opposite to *g*, the pyramidales appear through the tendons of the oblique muscles.

g, Ring of the external oblique muscle; with the spermatic chord, passing through it, and covered by the cremaster muscle.

SUPERIOR

SUPERIOR EXTREMITY.

a, Deltoides.

Above the clavicle, a portion of the trapezius is ſeen.

b, Biceps flexor cubiti.

At the bending of the arm are ſeen its tendon going towards the radius; and the part, from which the tendinous aponeuroſis that covers the fore-arm, is cut off.

On the inſide of the biceps, part of the triceps extenſor cubiti; and on the outſide, part of the brachialis internus.

c, Supinator radii longus.

d, Pronator teres.

e, Palmaris longus.

f, Palmaris brevis.

On the palm of the hand, the aponeuroſis palmaris is ſeen extended from the annular ligament at the wriſt, to the roots of the metacarpal bones of the four fingers.

g, Flexor carpi radialis.

h, Part of the flexor ſublimis perforatus.

i, Inſertion of the flexor carpi ulnaris.

k, Abductor pollicis.

INFERIOR EXTREMITY.

a, Tenſor vaginae femoris, the vagina or tendinous faſcia being cut off.

On the outſide of it a portion of the glutaeus maximus.

b, Part of the iliacus internus.

On the inſide of it, between *b* and *c*, part of the pſoas magnus.

c, Pectinalis.

d, Triceps longus.

e, Gracilis.

f, Sartorius.

g, Rectus

g, Rectus cruris.
Its tendon is seen inserted into the patella, from which a strong tendon is sent to be fixed to the tubercle of the tibia.

h. Vastus externus.

i, Vastus internus.

k, Tibialis anticus.

l, Peronaeus longus.
On the outside of it, a portion of the solaeus.

m, Extensor longus digitorum pedis, with the peronaeus tertius, and extensor proprius pollicis pedis.

n, Gastrocnemius externus, seu Gemellus.

o, Solaeus.

p, Ligamentum tarsi annulare.

q, Abductor pollicis pedis.

TABLE III.

THIS Table represents the second Layer of Muscles on the anterior Part of the whole Body.

MUSCLES *situated on the* HEAD *and* NECK.

a, Corrugator supercilii.

b, Temporalis.

c, Masseter.

d, Levator anguli oris.

e, Buccinator.

f, Orbicularis oris.
Opposite to the right ala nasi, the portion of this muscle which Albinus names
Nasalis labii superioris.

g, Depressor

TAB. III.

D.Lazars Sculp.

g, Depreſſor labii inferioris.

h, Sterno-cleido maſtoideus, which is

Seen below, ariſing from the ſternum and clavicle, by two heads.

i, Sterno-hyoidaeus.

On the outſide of it, the

Omo-hyoidaeus.

Further out, a portion of the

Hyo-thyroidaeus.

k, Levator ſcapulae.

TRUNK.

a, Subclavius.

b, Pectoralis minor.

c, Serratus magnus.

d, Rectus abdominis, divided into ſeveral fleſhy portions by its tendinous interſections.

e, Pyramidalis.

f, Obliquus aſcendens internus.

g, Spermatic cord, with the origin of the cremaſter muſcle.

SUPERIOR EXTREMITY.

a, Biceps flexor cubiti.

b, Short head of the biceps.

Beneath the upper part of it, a portion of the coraco-brachialis.

Beneath the under part, a portion of the brachialis internus.

c, Long head of the biceps.

At the bending of the arm, the tendon of the biceps, and the place where the tendinous aponeuroſis was cut from it, are ſeen.

d, Extenſor carpi radialis longior.

Beneath

Beneath it a portion of the
Extenſor carpi radialis brevior.

e, Flexor ſublimis perforatus.

f, Inſertion of the extenſor carpi ulnaris.

g, Extenſors of the thumb.

h, Opponens pollicis.
On the inſide of it a portion of the
Flexor pollicis brevis.

i, Tendon of the flexor longus pollicis manus, after paſſing through the flexor brevis pollicis manus.

k, Abductor minimi digiti manus.

l, Flexor parvus minimi digiti manus.

m, Ligamentum carpi annulare.

INFERIOR EXTREMITY.

a, Iliacus internus.
Between *a* and *b*, part of the pſoas magnus.

b, Pectinalis.

c, Triceps longus.

d, Gracilis.

e, Rectus cruris cut off near its origin.

f, Tendon of the rectus cruris cut off above the patella, from which a ſtrong tendon is ſent to be inſerted into a tubercle of the tibia.

g, Portion of the glutaeus medius.
On the inſide of it, part of the glutaeus minimus.

h, Vaſtus internus.

i, Vaſtus externus.

k, Cruraeus.

l, Inſertion of the biceps flexor cruris into the fibula.

m, Tendons of the gracilis and ſemitendinoſus inſerted into the tibia.

n, Solaeus.

o, Peronaeus longus.

p, Extenſor

TAB. IV.

FIG. 3.



FIG. 2.

FIG. 1.

FIG. 4.

FIG. 6.

FIG. 5.

D. Lizars Sculp.

p, Extenſor longus digitorum, with the peronaeus tertius on the outſide, and extenſor pollicis proprius on the inſide.

q, Solaeus.

r, Flexor longus digitorum.

ſ, Tendons of the tibialis poſticus and flexor longus digitorum pedis.

t, Flexor brevis digitorum pedis.

TABLE IV.

Fig. 1.

THIS Figure repreſents the right Eye-ball, the ſix Muſcles which move it, taken out of the orbit, with the Optic Nerve.

a, The eye-ball.

b, Optic nerve.

c, Muſculus trochlearis, ſeu obliquus ſuperior.

d, The trochlea, or pully, with a piece of the os frontis through which the tendon of the muſcle paſſes towards the eye-ball.

e, Obliquus inferior, with a piece of the ſuperior maxillary bone, from whence it ariſes.

f, Levator oculi.

g, Depreſſor oculi.

h, Adductor oculi.

i, Abductor oculi.

Fig. 2.

Repreſents the anterior Part of the Cartilage of the right Ear, with its proper Muſcles.

a, Helicis major.

b, Helicis minor.

c, Tragicus.

d, Antitragicus.

Tranſverſus auris, *vid.* Tab. viii. fig. 3.

Fig. 3.

Repreſents the third Layer of Muſcles, with ſome of the Ligaments, Cartilages, and naked bones on the anterior Part of the whole Body.

a, Depreſſor labii ſuperioris alaeque naſi.

b, Orbicularis oris, after moſt of the muſcles, which are fixed to it, and aſſiſt to form it, have been taken away.

c, Buccinator.

Above *c*, part of the pterygoidaeus externus is ſeen paſſing behind the coronoid proceſs of the lower jaw.

d, Levator labii inferioris.

e, Sterno-thyroidaeus.

Immediately above, and ſeemingly the continuation of it, the

Hyo thyroidaeus.

f, Scalenus anticus.

Contiguous to it, on the inſide, the

Scalenus medius.

Above it, a portion of the

Trachelo-maſtoidaeus.

Between the ſcalenus anticus, and ſterno-thyroideus, and and hyo-thyroidaeus, the

Rectus

Rectus capitis anterius major, and
Longus colli.

TRUNK.

a, Third row of external intercoſtal muſcles.
The reſt appear in the ſame manner between the other ribs.

b, Third row of internal intercoſtal muſcles.
The reſt appear between the other ribs.

c, Tranſverſalis abdominis.

d, The place from which the inferior part of the tendon of the tranſverſalis, that paſſes before the rectus and pyramidalis muſcles, is cut off.
Between theſe portions of each ſide, the peritonaeum is laid bare, and the ligaments of the bladder which were formerly the umbilical arteries and urachus.
Between this portion and the os pubis, the ſpermatic cord is ſeen cut.

e, The inferior edge of the upper part of the tendon of the tranſverſalis, which paſſes behind the rectus, and immediately adheres to the peritonaeum.

f, The anterior lamella of the internal oblique, which joined the tendon of the external to paſs over the rectus.
Between *f* and *g*, the poſterior lamella of the internal oblique, joining with the tendon of the tranſverſalis, to paſs behind the rectus.

g, The place at the linea alba, from which the tendon of the external oblique, and anterior lamella of the internal, were cut off.
At *g*, Umbilicus.

SUPERIOR EXTREMITY.

a, Subſcapularis.

b, Teres minor.

c, Coraco-brachialis.

The part from which the ſhort head of the biceps flexor cubiti was cut off from it, is ſeen at its upper end.

d, Brachialis internus.

e, Brachialis externus, or third head of the triceps.

f, Extenſor carpi radialis longior, and with it the extenſor carpi radialis brevior.

Both theſe are diſtinctly ſeen in the right hand.

Between the tendon of the brachialis internus and extenſor radialis, the

Supinator radii brevis is ſeen.

g, Flexor longus pollicis manus, with the fleſhy portion of it which ariſes from the internal condyle of the os humeri.

h, Flexor profundus perforans, which ſplits into four tendons, which paſs under the ligamentum carpi annulare.

i, Pronator quadratus.

k, Adductor minimi digiti manus.

l, One of the lumbricales.

The other three appear in the ſame manner, along the tendons of the flexor profundus.

Behind theſe, the internal interoſſei are ſeen.

INFERIOR EXTREMITY.

a, Glutaeus minimus.

b, Iliacus internus.

On the inſide of it, between *b* and *c*, the pſoas magnus.

c, Obturator externus.

d, Adductor brevis femoris.

e, Adductor magnus femoris.

f, Gracilis which is

Seen inſerted into the inſide of the head of the tibia.

g, The ſhort head of the biceps flexor cruris.

h, Peronaeus

b, Peronaeus longus.

i, Peronaeus brevis.

Between these two peronaei and tibia, the tibialis posticus is seen.

k, Tendon of the tibialis posticus, covering the tendon of the flexor longus digitorum pedis.

l, Extensor brevis digitorum pedis.

Fig. 4.

Represents the Lavatores Ani cut off from the Bones before, with their connection to the Extremity of the Rectum, and Bulb of the Urethra.

a, The urethra and its corpus cavernosum cut off.

b, Bulb of the urethra.

c, The circular fibres which surround the verge of the anus; by some named *sphincter internus*.

d, The anterior edge of the levator ani, cut off from the os pubis, and side of the pelvis.

Fig. 5.

Represents the anterior part of the sphincter Ani, and Bulb of the Urethra.

a, The urethra, and its corpus cavernosum cut.

b, Bulb of the urethra.

c, Left half of the sphincter ani, running obliquely upwards, to join with the right half.

d, The acute portion, which is inserted into the perinaeum.

Fig.

Fig. 6.

Repreſents the Corpora Cavernoſa of the Penis, Corpus Cavernoſum Urethrae, Acceleratores Urinae Tranſverſales and Erectores Penis.

a, Corpus cavernoſum penis cut.

b, Urethra, and its corpus cavernoſum cut.

c, Erector penis covering the crus.

d, Tranſverſalis penis.

e, Accelerator urinae.

f, Tranſverſalis penis alter, running along the inſide of the left erector.

TABLE V.

REPRESENTS a Back View of the Human Skeleton, with ſome of the Ligaments and Cartilages which connect the Bones.

HEAD AND NECK.

a, Os parietale, joined to its fellow by the ſagittal ſuture.

b, The os occipitis, joined to the parietal bones by the lambdoid ſuture, which is between *a* and *b*.

c, Os malae.

d, Maxilla inferior, with a view of the teeth of both jaws from behind.

e, The ſeven cervical vertebrae.

TRUNK.

a, The ſeventh or laſt true rib.

b, The twelfth or laſt rib.

c, The

TAB. V.

D.Lizars Sculp.

c, The five lumbar vertebrae.
d, Os ſacrum.
e, Os coccygis.
f, Os innominatum, divided into
g, Os ilium.
h, Os pubis.
i, Os iſchium.

SUPERIOR EXTREMITY.

a, The clavicle, joined outwards to the acromion of the ſcapula.
b, The ſcapula.
c, Os humeri.
d, Internal condyle.
e, External condyle.
f, Radius.
g, Ulna, its upper end, named *olecranon*; and near the wriſt, its ſtyloid proceſs.
h, The eight bones of the carpus.
i, The metatarſal bone of the thumb.
k, The metatarſal bone of the four fingers.
l, The two joints of the thumb.
m, The three joints or phalanges of the four fingers.

INFERIOR EXTREMITY.

a, Os femoris.
b, Trochanter major, and at the inſide of it the cervix.
c, Trochanter minor.
d, Internal condyle.
e, External condyle.
f, Tibia.
g, Fibula.
h, Malleolus internus.
i, Malleolus externus.

k, The

k, The ſeven bones of the tarſus.

l, The metatarſus.

m, The joints or phalanges of the toes.

TABLE VI.

REPRESENTS a Back View of the Muſcles which are immediately ſituated below the common Integuments.

HEAD AND NECK.

a, Part of the occipito-frontalis muſcle, with its aponeuroſis.

b, Attollens aurem.

c, Anterior auris.

d, Retrahentes auris.

TRUNK.

a, Trapezius, ſeu cucularis.

b, Its tendinous edge joining with its fellow in the nape of the neck, which is called *ligamentum nuchae* ſeu *colli*.

c, The fleſhy belly of the latiſſimus dorſi.

d, The tendon of the latiſſimus dorſi, which ariſes in common with the ſerratus poſticus inferior.

e, Part of the obliquus externus abdominis.

SUPERIOR EXTREMITY.

a, Deltoides.

b, Infraſpinatus, with a portion of the teres minor and major below it.

c, Triceps

TAB. VI.

D.Lazara Sculp.

c, Triceps extenſor cubiti.

Its tendon is ſeen inſerted into the head of the ulna, called *olecranon*; and, on the inſide of it, the *anconaeus*.

d, Extenſor carpi radialis longior, covered by a portion of the ſupinator radii longus; and, under it, a portion of the extenſor carpi radialis brevior.

e, Extenſor digitorum communis manus, which ſplits into four tendons, and paſs with the indicator, under the ligamentum carpi annulare externum, at the extremities of the metacarpal bone, under ligaments proper to themſelves; and are loſt in a broad tendon, which covers the back of the four fingers.

f, Extenſor oſſis metacarpi pollicis manus.

g, Extenſor primi internodii pollicis manus.

h, Extenſor ſecundi internodii pollicis manus.

i, Extenſor carpi ulnaris.

k, Part of the flexor carpi ulnaris.

Under it, part of the

Flexor profundus perforatus.

And on the inſide, part of the

Flexor ſublimus perforatus, which are more diſtinctly ſeen on the right fore-arm. Likewiſe, on the right hand, are ſeen part of the abductor pollicis manus, abductor minimi digiti manus, and the aponeuroſis palmaris.

INFERIOR EXTREMITY.

a, Glutaeus maximus.

b, Part of the glutaeus medius.

c, Part of the tenſor vaginae femoris.

d, Vaſtus externus.

e, The long head of the biceps flexor cruris;

And beneath it,

f, Part of the ſhort head.

g Semitendinoſus;

g, Semitendinoſus:
And beneath it, on each ſide,
A portion of the ſemimembranoſus is ſeen.

h, Gracilis.
On the outſide of it,
A portion of the adductor magnus is ſeen.

i, A ſmall part of the vaſtus internus.

k, Gaſtrocnemius externus, ſeu gemellus;
And within its outer head,
A Portion of the plantaris.

l, Solæus ſeu gaſtrocnemius internus.

m, Tendo Achillis, with the plantaris.

n, Peronaeus longus.

o, Peronaeus brevis; between it and the tendo Achillis, a portion of the flexor longus digitorum pedis.

p, Tendons of the extenſor longus digitorum pedis, with the peronaeus tertius paſſing under the ligamentum tarſi annulare; and the flexor brevis digitorum pedis is ſeen beneath them.

q, Abductor minimi digiti pedis; and above it the tendons of the peronaeus longus and brevis, paſſing under proper ligaments of their own.

TABLE VII.

REPRESENTS the ſecond Layer of the Muſcles on the Back-part of the Body.

HEAD AND NECK.

a, Temporalis; its tendon is ſeen paſſing below the zygoma.
b, Maſſeter.

TAB. VII.

D.Lucas Sculp.

b, Maſſeter.

c, Splenius capitis et colli.

d, Portion of the complexus.

e, Levator ſcapulae, ſeu muſculus patientiae.

TRUNK.

a, Rhomboides major.

b, Rhomboides minor.

And immediately above it, the upper edge of the ſerratus poſticus ſuperior is ſeen.

c, The ſerratus poſticus ſuperior on the right ſide.

d, Serratus poſticus inferior.

e, Part of the ſpinalis dorſi.

f, Part of the longiſſimus dorſi.

g, Part of the ſacro lumbalis.

h, Serratus magnus.

i, The broad tendon, by which the latiſſimus dorſi begins, and from which the tendon of the ſerratus poſticus inferior is inſeparable.

k, Part of the obliquus internus aſcendens abdominis.

l, The ſphincter ani, fixed to the point of the os coccygis; at the ſide of which the coccygaeus, and a portion of the levator ani, are ſeen; and lower down, oppoſite to *l*, part of the tranſverſalis penis.

SUPERIOR EXTREMITY.

a, Supra-ſpinatus.

b, Infra-ſpinatus.

c, Teres minor.

d, Teres major.

e, Triceps extenſor cubiti.

f, Its head called *longus*.

g, The brevis: And,

h, A ſmall portion of the third head, named *brachialis externus*.

i, The tendon of the triceps, inſerted into the olecranon.

k, Part of the brachialis internus.

l, Anconaeus, which ſeems to be continued from that part of the brachialis externus immediately above it.

m, Extenſor carpi radialis longior; and beneath it, the brevior; both are ſeen at the wriſt, inſerted into the metacarpal bones of the fore and middle fingers.

n, Flexor carpi ulnaris.

o, Part of the ſupinator radii brevis.

p, Extenſor oſſis metacarpi pollicis manus.

q, Extenſor primi internodii pollicis manus.

r, Extenſor ſecundi internodii pollicis manus.

ſ, Indicator, inſerted into the root of the firſt joint of the fore-finger.

t, One of the three external interoſſei manus. The other two are diſtinctly ſeen without letters.

u, One of the tendons of the extenſors of the fingers cut; and the ſame is ſeen in each of the other three fingers, joining with the tendons and aponeuroſes of the interoſſei and lumbricales, and ſpread upon the back of the fingers.

N. B. On the right hand, part of the flexors of the fingers, the abductor pollicis and minimi digiti, are ſeen.

INFERIOR EXTREMITY.

a, Glutaeus medius.

b, Pyriformis.

c, The two muſcles called *gemini*, between which the tendon and fleſhy belly of the obturator internus paſſes over the tuberoſity of the os iſchium, are ſeen within the pelvis, partly covered by the coccygaeus and levator ani.

d, Quadratus

d, Quadratus femoris.

e, Vaſtus externus.

f, *f*, Parts of the triceps magnus.

g, Long head of the triceps flexor cruris, and beneath it part of the ſhort head is ſeen.

h, Semitendinoſus, and beneath it parts of the ſemi-membranoſus are ſeen on each ſide of it.

i, Gracilis.

k, A ſmall portion of the vaſtus internus.

l, Poplitæus.

m, The fleſhy belly of the plantaris; and its long ſlender tendon is ſeen paſſing over the inſide of the ſolæus.

n, Solæus.

o, The place where the tendon of the gemellus was cut off; but the fleſh of the ſolæus runs farther down.

p, Tendon Achillis, with the plantaris.

q, Peronæus longus, paſſing at the outer ankle to the ſole of the foot; beneath it, the peronæus brevis to the root of the metatarſal bone of the little toe; and, between it and the tendo Achillis, a portion of the flexor longus digitorum pedis.

r, Tendons of the extenſor longus digitorum pedis, with the peronaeus tertius; and beneath theſe, the extenſor brevis digitorum pedis.

ſ, Flexor brevis minimi digiti pedis.

TABLE

TABLE VIII.

Fig. 1.

REPRESENTS the third Layer of Muſcles on the poſterior part of the Body, with ſome of the Ligaments and naked Bones.

MUSCLES *on the* HEAD *and* NECK.

a, Part of the buccinator.

b, Complexus.

c. Trachelo-maſtoidaeus; on the outſide of it, the tranſverſalis colli.

d, Scalenus medius.

e, Scalenus poſticus.

TRUNK.

a, Spinalis dorſi; and beneath it, the multifidus ſpinae.

b, Longiſſimus dorſi, which ſends off a fleſhy ſlip to the trachelo-maſtoidaeus.

c, Sacro lumbalis, with the cervicalis deſcendens ſent off from it along the ſide of the neck, and outſide of the tranſverſalis colli.

d, Semiſpinalis dorſi.

e, Tranſverſalis abdominis.

N. B. The ſpaces between the ſpinous proceſſes of the vertebrae have muſcular faſciculi between them, particularly thoſe of the neck; and are named *interſpinales colli*, *dorſi*, and *lumborum*; but thoſe of the back ſeem to be tendinous and ligamentous.

SUPERIOR

TAB. VIII.

D.Lizars Sculp.

SUPERIOR EXTREMITY.

a, Teres major.

b, Part of the coraco-brachialis.

c, Part of the brachialis internus.

d, The third head of the triceps extenſor cubiti, called *brachialis externus*, after the longus and brevis have been cut off.

e, Extenſor radialis longior.

f, Extenſor radialis brevior.

g, Part of the flexor profundus perforans.

h, Supinator radii brevis.

i, Part of the adductor pollicis manus.

k, One of the three external interoſſei; the other two may be eaſily diſtinguiſhed without letters.

l, Tendons of the extenſors of the fingers, joining with thoſe of the lumbricales and interoſſei, which form a tendinous expanſion on the back of the four fingers.

N. B. On the right hand, part of the flexors of the fingers and thumb, part of the adductor pollicis, and the whole of the adductor minimi digiti, are ſeen.

INFERIOR EXTREMITY.

a, Glutaeus minimus.

b, Obturator internus; its fleſhy belly is ſeen within the pelvis. Beneath *b*, the tendon of the obturator externus.

c, Semimembranoſus.

d, The ſhort head of the biceps flexor cruris.

e, Triceps magnus.

f, Gracilis.

In the ham, the origins of the two heads of the gaſtrocnemius externus and plantaris are ſeen.

g, Poplitaeus.

h, Tibialis poſticus.

i, Flexor

i, Flexor longus digitorum pedis.

k, Flexor pollicis longus.

l, Peronaeus longus, running down to be inserted into the metatarsal bone of the little toe.

Beneath it, the peronaeus brevis, passing to the sole of the foot.

m, Extensor brevis digitorum pedis.

n, Part of the flexor longus digitorum pedis.

Fig. 2.

Represents the fourth Layer of Muscles on the posterior part of the Neck.

a, Rectus capitis posticus major.

b, Rectus capitis posticus minor.

c, Obliquus capitis superior.

d, Obliquus capitis inferior.

e, Scalenus medius.

f, Part of the multifidus spinae, covered by the semispinalis colli.

Between the spinous processes of the vertebrae, the interspinales colli are seen double; because these processes are bifurcated.

Between the transverse processes, the posterior row of the intertransversales colli is seen.

Fig. 3.

Represents a portion of the posterior part of the cartilage of the ear, to shew,

a, Transversus auris.

Situated on the parts opposite to the antihelix and scapha.

Fig.

Fig. 4.

Repreſents an outline of the Baſe of the Os Occipitis, and inſide of the Lower Jaw; with a view of the Muſcles which ſurround the back Part of the Larynx and Pharynx.

a, The upper part of the membrane of the pharynx.

b, The trachea, cut.

c, The oeſophagus, cut.

d, The inner tranſverſe fibres of the oeſophagus, laid bare.

e, The outer fibres deſcending obliquely backwards.

f, Conſtrictor inferior pharyngis.

g, Conſtrictor medius pharyngis.

h, The cornu of the os hyoideus.

i, Conſtrictor ſuperior pharyngis.

k, The part of it which joins with the buccinator.

l, Stylo-pharyngaeus.

Fig. 5.

Repreſents an outline of the Inſide of the Os Pubis, Os Iſchium, and Back of the Os Coccygis, after the Os Sacrum and Ligaments have been taken away; with a poſterior view of the Levatores Ani, and Extremity of the Rectum, reſembling the ſhape of a funnel.

a, The anterior portion of the levator ani, viewed on its inſide within the pelvis, ariſing from the os pubis and upper part of the foramen thyroidaeum.

b, Its origin from the ſpinous proceſs of the os iſchium.

c, The poſterior part ſeen on its outer ſide.

d, Its

d, Its insertion into the os coccygis, below which the fleshy fibres are continued with its fellow.

l, The circular fleshy fibres surrounding the extremity of the rectum, which authors name *sphincter internus ani*.

m, The anus.

Fig. 6.

Represents an outline of the Back of the Os Coccygis and posterior part of the sphincter Ani.

a, The insertion of the sphincter ani into the extremity of the os coccygis; to which, from

b, The anus,

The fibres meet from each side, in angles; which are more acute as they point upwards.

Fig. 7.

Represents the first Layer of Muscles on the Sole of the Foot, after the Aponeurosis Plantaris has been taken away.

a, Abductor pollicis pedis.

bb, Abductor minimi digiti pedis.

c, Flexor brevis digitorum pedis, which splits into four tendons, that are perforated by the tendons of the flexor longus digitorum pedis.

Between these tendons the lumbricales are seen.

d, Tendon of the flexor pollicis longus.

And beneath it, the flexor pollicis brevis.

e, Transversalis pedis.

Fig.

TAB IX



D. Lizars Sculpt

Fig. 8.

Repreſents the ſecond Layer of Muſcles on the Sole of the Foot, after the Abductor Pollicis Pedis, and Abductor Minimi Digiti Pedis, and the Flexor Brevis Digitorum Pedis, have been taken away.

a, Tendon of the flexor longus digitorum pedis.

bb, Flexor acceſſorius, ſeu maſſa carnea Jacobi Sylvii.

cccc, The four lumbricales ariſing from the tendons of the flexor longus digitorum.

dd, Tendon of the flexor pollicis longus; which, in its progreſs, joins the tendon of the flexor longus digitorum pedis.

e, Tendon of the tibialis anticus.

f, Tendon of the peronaeus longus.

g, Tendon of the peronaeus brevis.

h, Flexor pollicis brevis, with a portion of the abductor pollicis, on the inſide of the great toe.

On the outſide of *i*, is the abductor pollicis.

k, Flexor brevis minimi digiti pedis.

l, A ligament which ſupports the bones of the tarſus.

Before *l*, two of the interoſſei are ſeen.

m, Tranſverſalis pedis.

TABLE IX.

REPRESENTS a front view of the Bones of the Pelvis in a Female Skeleton.

A, The five vertebrae of the loins.

B, The os ſacrum.

C, The os coccygis.

DD, The ossa ilium.

EE, The ossa ischium.

F, The ossa pubis.

GG, The foramina magna.

HH, The acetabula.

IIII, The brim of the pelvis, or that circumference of its cavity which is described, at the sides by the inferior parts of the ossa ilium, and at the fore and back parts by the superior parts of the ossa pubis and sacrum.

In this table, besides the general structure and figure of the several bones, the dimensions of the brim of the pelvis, and the distance between the under parts of the ossa ischium, are particularly to be attended to; from which it will appear, that the cavity of the brim is wider from side to side than from the back to the fore part; but that the sides below are in the contrary proportion. In general, the brim of the female pelvis measures about five inches and a quarter from side to side, and four inches and a quarter from the back to the fore part, there being likewise the same distance between the inferior parts of the ossa ischium. All these measures, however, must be understood to be taken from the skeleton; for, in the living subject, the cavity of the pelvis is considerably diminished by the integuments and contents.

TABLE

D.Lizars Sculp!

TABLE X.

REPRESENTS a lateral and internal View of the Pelvis, divided longitudinally.

A, The three lower vertebrae of the loins.

B, The os ſacrum.

C, The os coccygis.

D, The left os ilium.

E, The left os iſchium.

F, The left os pubis.

G, The acute proceſs of the os iſchium.

H, The foramen magnum.

III, The brim of the pelvis.

This plate ſhews the diſtance from the ſuperior part of the os ſacrum to the oſſa pubis, as well as from the laſt mentioned bones to the os coccygis, which in each amounts to about four inches and a quarter. The depths of the poſterior, lateral, and anterior parts of the pelvis are likewiſe ſhewn, not in the line of the body, but in the line of the pelvis, from its brim downwards, which is generally three times deeper on the poſterior than the anterior part, and twice the depth of this laſt on the ſides.

In this figure appears alſo the angle made by the projection of the laſt vertebra of the loins, and firſt of the ſacrum; and likewiſe the concavity or hollow ſpace in the poſterior internal part of the pelvis, ariſing from the curvature of the ſacrum and coccyx: Finally, the diſtance from the coccyx to the poſterior parts of the oſſa iſchium are alſo here expreſſed.

A

SYSTEM OF ANATOMY.

PART IV.

OF THE DIFFERENT CARTILAGES, LIGAMENTS, &c. OF THE FRESH BONES*.

OF THE HEAD.

THE condyloid proceſſes of the os occipitis, the glenoid cavities or articular foſſulae of the oſſa temporum, the eminences next theſe cavities, and the condyloid proceſſes of the lower jaw, are all covered over with very white and ſmooth cartilages; and there is likewiſe an interarticular or moveable cartilage in each articulation of the lower jaw with the temporal bones.

This cartilage is thick near the circumference; very thin and tranſparent, and ſometimes perforated in the middle. The lower ſide is uniformly concave, anſwering to the oblong convexity of the maxillary condyle; but the upper ſide is partly concave, and partly convex, ſuited to the foſſula and eminence of the temporal bone. It is fixed by its circumference to the inner ſide of the capſular ligament.

The

* The anatomy of the Freſh Bones in general, has been conſidered in the firſt part of this Work.

The remaining cartilages of the bones of the head, viz. the cartilaginous ſeptum, and other cartilages of the noſe, the ſmall cartilaginous pulley in each orbit, the cartilages of the outer ear, and thoſe which are joined to the os hyoides, muſt be referred to the deſcription of the viſcera.

The ligaments of the bones of the head are, Thoſe of the articulation of the lower jaw with the temporal bones; thoſe between the occipital bone and vertebrae of the neck; thoſe by which the os hyoïdes is connected to the ſtyloid proceſs.

LIGAMENTS BY WHICH THE LOWER JAW IS FIXED TO THE TEMPORAL BONES.

Ligamentum capſulare, compoſed of firm and ſtrong fibres, fixed by one extremity round the glenoid or articular foſſula and eminence of each temporal bone; by their middle round the interarticular cartilage, and by their other extremity round each condyle of the lower jaw, in ſuch a manner as to allow the intermediate cartilage to follow the motions of the condyles, and to change their ſituation from the glenoid cavities to the tubercles of the zygomatic proceſſes, and to return again, as was mentioned in the deſcription of the bones.

Ligamentum laterale maxillae inferioris, which ariſes from the inner ſurface of the angle of the lower jaw, near the paſſage where the veſſels and nerves go into the bone, is fixed to the root of the ſtyloid proceſs, and to the poſterior margin of the articular cavity of the temporal bone. It aſſiſts in keeping the jaw in its proper place.

LIGAMENTS BETWEEN THE OCCIPITAL BONE AND VERTEBRAE OF THE NECK.

Ligamenta capſularia, ariſe from the edges of the condyloid

loid proceſs of the os occipitis, and are fixed to the edges of the ſuperior oblique proceſſes of the firſt vertebra.

Ligamentum latum anterius, ariſes from the fore part of the foramen magnum occipitis, and runs down to be fixed to the anterior arch of the firſt vertebra of the neck.

Ligamentum latum poſterius, ariſes from the poſterior margin of the foramen magnum occipitis, and is inſerted into the upper part of the poſterior arch of the firſt vertebra.

Ligamentum proceſſus odontoidis ſeu perpendiculare, ariſes from the fore-part of the foramen magnum, and runs down to be fixed to the proceſſus dentatus of the ſecond vertebra. This ligament is ſhort, but remarkably ſtrong; it aſſiſts in fixing the head to the ſpine, but is twiſted in the rotation.

Ligamenta lateralia, are two ſhort, but very ſtrong ligaments, which run over from each ſide of the proceſſus dentatus to be fixed to the inner ſide of the firſt vertebra, and to the edge of the foramen magnum. See deſcription of the Vertebrae.

Ligamentum cervicale ſeu nuchae, ariſes from the ſpinous proceſs of the os occipitis, runs down upon the back-part of the neck, adhering to the ſpinous proceſſes of the cervical vertebrae, and giving origin to part of the trapezius muſcle.

The bones of the head, as well as all the other bones of the body, are covered by a particular membrane; but that which covers the cranium is termed *pericranium*, while that ſurrounding the other bones is called ſimply *perioſteum*.

The internal ſtructure of the bones of the head being for the moſt part cellular, they contain alſo diſtinct portions

tions of marrow included in membranous cells lying in the diploe.

The ſynovial glands of the maxillary and occipital articulations have nothing particular to them; they are proportioned to the joints to which they belong, and lie between the capſular ligaments and circumference of the cartilages.

OF THE VERTEBRAE.

The cartilages of the vertebrae in general are of two kinds; one proper to each vertebra, the other common to two vertebrae that lie next each other. The firſt may be termed *cartilages of articulation;* the others, *cartilages of ſymphyſis.*

The proper articular cartilages of each vertebra of the ſpine, are thoſe four which cover the ſurfaces of the oblique proceſſes. In the natural ſtate, they are very white and ſmooth, and much thicker than in dry bones. Their circumference is the ſame with that of the articulated ſides of the proceſſes, except in thoſe places where there are ſmall ſuperficial notches. In the firſt vertebra of the neck, and vertebrae of the loins, theſe cartilages are thicker than in the reſt.

The cartilages of the two inferior oblique proceſſes of the firſt vertebra, and of the two ſuperior oblique of the ſecond, ſeem to be diſproportionate, though not ſo much as in dry bones; and in ſome ſubjects we find moveable or interarticular cartilages between the proceſſes of theſe two vertebrae.

The firſt vertebra of the neck has a cartilaginous covering on the back-part of the anterior arch, correſponding with another on the fore ſide of the proceſſus dentatus of the next vertebra; ſo that each of thoſe two vertebrae has

five

five articular cartilages, beſides the interarticular ones already mentioned.

The vertebrae of the back, beſides the four cartilages of their oblique proceſſes, have others which do not belong to their articulations with each other, viz. thoſe that cover the lateral foſſula in the bodies of theſe vertebrae, and the foſſulae of their tranſverſe proceſſes, by both of which they are articulated with the ribs.

The cartilages of ſymphyſis lie between the bodies of the vertebrae, uniting them cloſely together; their breadth and circumference anſwering exactly to that of the ſurfaces to which they are connected; but their height or thickneſs is different in each claſs of the vertebrae. Between the vertebrae of the loins, they are a third or fourth part of an inch in thickneſs, according to the ſize of the ſubject: In thoſe of the neck, they are not ſo thick; and the thinneſt of all are between the vertebrae of the back.

Theſe cartilages are not of an equal thickneſs in all their parts. Thoſe on the neck and loins appear to be thickeſt on the anterior ſide, and thoſe of the back rather thickeſt on the poſterior ſide, but theſe differences are moſt remarkable in the vertebrae that lie near the middle of each claſs.

The internal ſtructure of theſe cartilages is different from that of all the other cartilages of the body; and indeed they reſemble the reſt in nothing but in whiteneſs and elaſticity. When we view their circumferences only they ſeem to be one uniform maſs, as the others generally are; but when they are divided by an inciſion parallel to that ſurface of the vertebrae to which they are joined, we ſee they are compoſed of many cartilaginous concentrical lamellae contained within each other. The moſt external lamellae are fibrous, thickeſt, and firmeſt, and ſeparated by conſiderable intervals: the internal approach nearer and nearer together,

becoming

becoming gradually thinner, and of a ſofter conſiſtence, till at laſt they are almoſt in the form of a glairy liquor in the centre.

Theſe rings do not form an entire circumference; being turned inwards on the back-part, anſwering to the fore-part of the paſſage for the ſpinal marrow. They lie horizontally between the vertebrae. The interſtices of the rings are filled with a mucilaginous ſubſtance, leſs fluid than that of the joints. Each lamina taken ſeparately is very pliable, according to its length; but taken together, they are not ſo eaſily bent, partly becauſe of their circular figure, and partly becauſe of their vicinity and multiplicity. They yield, however, in the inflections of the ſpine; and their external ſurface, which in the ordinary ſituation of the ſpine is even with the ſurface of the vertebrae, becomes prominent or juts out on that ſide towards which the inflection is made, the cartilages being then compreſſed by the vertebrae.

They likewiſe yield on all ſides, without any inflection of the ſpine, to the weight of the upper part of the body: but this is done by very ſmall and imperceptible degrees, and eſpecially at the under part of the true vertebrae, and when the body is loaded with an additional weight.

They reſtore themſelves afterwards merely by being freed from compreſſion: So that a man is really taller after lying, than after he has walked or carried a burden for a length of time; the moſt natural and ſimple reaſon that can be given for the different height of the ſame perſon at different times, firſt obſerved in England, and afterwards confirmed by Mr Morand, a member of the Royal Academy of Sciences, being the different ſtate of the intervertebral cartilages. According to Sabatier, &c. the ſame perſon is ſometimes more than four or five lines, or twelfths of an inch, higher in the morning than in the evening. The interverte-

bral cartilages of the neck, lying for the moſt part between the convex ſide of one vertebra, and the concave ſide of another, are of a greater extent in proportion to the ſize of theſe vertebrae, than thoſe of the back and loins. Without this convexity and hollowneſs in theſe vertebrae, which are the leaſt of all, the cartilages could not have been made large enough to reſiſt ſtrains and great exertions.

Ligaments *of the* Vertebrae.

The vertebrae are ſtrongly connected to each other by different kinds of ligaments; ſome of which are proper to a certain number of them, others are common to the whole.

Ligamentum tranſverſum vertebrae colli primae, ariſes from a rough protuberance on the inner ſide of the firſt vertebra, and goes acroſs to the other ſide behind the proceſſus dentatus, which it prevents from wounding the ſpinal marrow in the rotation of the head. About the middle of the fore-ſide, its texture is very cloſe where the proceſſus dentatus plays upon it.

The ligaments of the proceſſus dentatus of the ſecond vertebra have been already deſcribed.

Ligamentum anticum commune vertebrarum. One of the moſt remarkable is a ſtrong ligamentous band, which embraces their convex ſurface from the upper to the under end of the ſpine. It begins at the ſecond vertebra of the neck, and paſſes down as low as the os ſacrum, becoming gradually larger and ſtronger in its deſcent. The fibres of this ligament have a longitudinal direction; but it is much thicker in its middle that at its ſides. After it has arrived at the laſt lumbar vertebra, it ſpreads over the anterior ſurface of the os ſacrum, where it becomes thinner, and by degrees vaniſhes near the under end of this bone. Through its whole courſe it ſends off many ſmall proceſſes to be fixed

ed to the bodies of the vertebrae, by which their connection is made more ſecure.

Ligamenta intervertebralia. Behind the former ligament each vertebra is connected to that above and below it by numerous, ſhort, but ſtrong ligaments, which croſs each other obliquely, and are fixed round the edges of the body of each vertebra. Theſe crucial ligaments cover the circumference of the intervertebral cartilages, and adhere cloſely to them. They ſeem to be looſer in the cervical and lumbar vertebrae than in thoſe of the back; and by that means yield to the cartilages in the different flexions of the ſpine already mentioned.

Ligamentum poſticum commune vertebrarum. The ſpinal canal is lined with a ligament ſomewhat ſimilar to that which covers the anterior convex ſurface of the vertebrae. It begins at the ſecond vertebra of the neck; and after having ſent a conſiderable proceſs, which paſſes behind the tranſverſe ligament of the firſt vertebra, to be fixed to the anterior part of the foramen magnum, it deſcends on all the other vertebrae, to end at the lower part of the os ſacrum. The real ligamentous fibres occupy little more than the middle of the bodies of the vertebrae. Thoſe which are ſtretched over the lateral parts are very thin, and properly ſpeaking purely membranous. Winſlow deſcribes this as a complete tube, while Weitbrecht denies its exiſtence at the back part of the canal; but admits of an additional membrane there adhering firmly to the dura mater. It is only attached to the ſuperior and inferior edges of the vertebrae, leaving at their middle a ſpace occupied by a kind of tranſverſe ſinus, which communicates with others ſituated longitudinally upon the ſides of the poſterior part of the whole canal.

Ligamenta interſpinoſa, are ſhort and firm ligaments, which run from the whole upper edge of the bony bridge and ſpinous

ous proceſſes of one vertebra, to the correſponding parts of the vertebra next; and thereby joining the different vertebrae together, and dividing the muſcles on the right from thoſe on the left ſide of the ſpine.

Ligamenta intertranſverſalia, are ſhort ſmall ligaments placed between the extremities of all the tranſverſe proceſſes.

Ligamenta capſularia, are formed of numerous ſhort, ſtrong, ligamentous faſciculi, ariſing from and ſurrounding the oblique proceſſes of all the vertebrae. The two oblique proceſſes of the os ſacrum are joined to the inferior oblique proceſſes of the laſt vertebra of the loins, in the ſame manner as thoſe of the lumbar vertebrae.

Of the STERNUM and RIBS.

The ſternum of an adult has commonly ſixteen cartilages; fourteen of which are articular, the other two ſymphyſes. Of the articular cartilages, two belong to the articulations of the clavicles; and twelve to thoſe of the true ribs, from the ſecond to the ſeventh incluſive. The two ſymphyſes are thoſe between the ſternum and the firſt rib on each ſide.

There is likewiſe another ſymphyſes, by which the upper portion of the ſternum is connected to the lower; the cartilage of which is often obliterated in an advanced age. But at an early period of life this cartilage can be diſtinctly ſeen; and it allows a conſiderable degree of motion between theſe two bones.

The cartilago enſiformis is often bony towards the ſternum, and more or leſs cartilaginous at the other end. In very aged perſons it has been found entirely oſſified, and ſometimes wholly cartilaginous even in adults.

All the ribs have cartilaginous portions, which differ from each other in length, breadth, incurvation, adheſion, and

and in their extremities; all which were explained in the deſcription of the ſkeleton. We have only to obſerve here, that theſe cartilages are whiter, more poliſhed, broader, and thicker in the natural ſtate than when they are dried. The cartilages of the falſe ribs are naturally more ſlender and pliable than thoſe of the true ribs: The middle or inner ſubſtance acquires the conſiſtence of bones in old age; and their extremities ſometimes oſſify, and are immoveably fixed to the ſternum. The poſterior extremities of the ribs are likewiſe tipped with cartilage where they are joined to the vertebrae.

Ligaments *proper to the* Sternum.

Membrana ſterni propria, is a firm expanſion, compoſed of many tendinous fibres, running in different directions, but chiefly longitudinally, covering both the external and internal ſurface of the ſternum. On the fore-part of the ſternum the external fibres begin at the articulation of the cartilages of the ribs, and run acroſs in a radiated manner to their fellows on the oppoſite ſide, while the internal fibres have a longitudinal direction.

Ligamentum cartilaginis enſiformis, is compoſed of tendinous fibres ſimilar to the former, ariſing from the cartilaginous extremity of the ſeventh rib and correſponding part of the ſternum; and which, after deſcending obliquely, are fixed to the cartilago enſiformis. The fibres of this are intermixed with thoſe of the membrana ſterni.

Ligaments *between the* Sternum *and* Ribs.

Ligamenta capſularia cartilaginum coſtarum verarum, ariſe round the cartilages of the ſeven true ribs, to be fixed to the articular pits in the ſides of the ſternum. On the upper and under ſide of each articulation, theſe ligaments are very ſhort; but on the anterior ſide many fibres are produced, which

run

run in a radiated manner on the fore-part of the ſternum to the cartilages on the oppoſite ſide.

Ligaments *proper to the* Ribs.

Ligamenta coſtarum ipſarum propria, are ligaments by which the ribs are joined to each other. They deſcend ſomewhat in a perpendicular direction from the cartilage of each rib to that of the next; but the ligaments between the three laſt ribs are longer and looſer than thoſe of the reſt. Hence the two laſt ribs are leſs ſteady in their motions.

Ligaments *between the* Ribs *and* Vertebræ.

Ligamenta capſularia capitulorum majorum coſtarum, are ſhort ligamentous faſciculi which ariſe round the cartilaginous ſurface of the head of each rib, and are fixed to the circumference of the ſmall pits in the ſides of the bodies of the vertebrae and intervertebral cartilages.

Ligamenta capſularia capitulorum minorum, ariſe from the tubercles of the ten uppermoſt ribs, and are fixed round the articular pit on the point of the tranſverſe proceſſes of the vertebrae of the back; much in the ſame manner with thoſe between the heads of the ribs and vertebrae.

Ligamenta interna colli coſtarum, ariſe from the upper part of the neck of the ribs, and are fixed to the inferior ſurface of the tranſverſe proceſſes.

Ligamenta externa colli coſtarum, ariſe from the outer ſurface of the ſuperior margin of the neck of all the ribs. They aſcend obliquely, to be fixed near the inferior oblique proceſſes of all the vertebrae excepting the firſt.

Ligamenta duo ſpecialia, ariſe by a broad origin from the inferior margin of the laſt rib, and are fixed to the tranſverſe proceſs of the firſt and ſecond lumbar vertebrae.

The ligamentous expanſions of the vertebrae are in place

of

of a periosteum: at least they are blended together both on the inner and outer side of the spine. The sternum and bony portions of the ribs have a periosteum like the other bones.

The cartilaginous portions of the ribs are covered by a membrane of the same kind, termed *perichondrium*. As the internal structure of these bones is cellular or spongy, they contain only small separate portions of marrow, or a red medullary juice, like that in the vertebrae.

The synovial glands of all these articulations are very small; but are accompanied by many other fatty moleculae lying round each joint. The inner surface of the ligamentous substance which lines the bony canal of the spine, is lubricated by an oily, or adipose substance, which shall be mentioned hereafter.

Of the SUPERIOR EXTREMITIES.

The scapula in many subjects has a small cartilaginous border along its whole basis; which in children is tolerably distinct, but in adults it disappears.

The glenoid cavity of this bone is covered with a cartilage, which is thicker towards the circumference than in the middle, and a little raised above the edge of the bone. This thickness of the cartilaginous circumference makes the cavity greater than it appears in the skeleton; and sometimes, in its place, there is an additional border, which is thickest at the circumference of the cavity, thin toward the bottom, and very narrow. It is of a pliable slippery substance, yet something different from that of a cartilage: resembling, in some measure, the border of the acetabulum of the os innominatum.

The small cartilaginous surface of the acromion, mentioned in the treatise of the dry bones, is thicker in the natural state, and very little convex, and the triangular surface

face at the extremity of the ſpine of the ſcapula, near the baſis, is covered with a very thin cartilaginous lamina; but, being tranſparent, it does not appear very white. There are no other cartilages commonly found in the ſcapula: though we ſometimes obſerve in dry bones ſeveral places which ſeem to have been cartilaginous; but this is owing to the dried remains of ligaments and tendons.

The ſternal extremity of the clavicle is cruſted over with a cartilage which is a little convex, and covers its whole triangular ſurface; beſides which it has another moveable interarticular cartilage, reſembling that at the articulation of the lower jaw, and in ſome meaſure ſerving the ſame purpoſe. The ſmall cartilaginous ſurface of the ſcapular extremity of the clavicle, anſwering to that of the acromion, is much thicker in freſh than in dry bones, and appears, like that of the acromion, to be a little convex.

Between theſe two cartilages of the clavicle and acromion, there is in ſome ſubjects a thin interarticular cartilage very ſmooth on both ſides.

Ligaments *of the* Clavicle *and* Scapula.

Ligamentum interclaviculare, is a long narrow ſtrong ligament which goes behind the furca of the ſternum, from the internal angle of one clavicle to that of the other.

Ligamenta capſularia antica claviculae, are ſhort and ſtrong ligaments ariſing round the ſternal extremity of the clavicle, near the edge of the triangular ſurface; and from thence paſſing over the interarticular cartilage, to which they adhere, are inſerted round the clavicular cavity of the ſternum.

Ligamentum rhomboideum, ariſes from the inferior rough ſurface at the anterior extremity of the clavicle, and running obliquely, is fixed to the cartilage of the firſt rib.

Ligamenta capſularia poſtica. The articulation of the ſcapulary

ſcapulary end of the clavicle with the acromion, is ſtrengthened quite round by thick ſtrong ligaments which go from one bone to the other.

Ligamentum trapezoideum ſcapulae, ariſes from the internal ſurface of the coracoid proceſs of the ſcapula, and ends at the poſterior extremity of the clavicle.

Ligamentum conoideum ſcapulae, ariſes from the root of the coracoid proceſs, is inſerted into the rough protuberance of the poſterior extremity of the ſcapula.

Ligamentum proprium ſcapulae anticum, ariſes from the external ſurface of the coracoid proceſs, and is fixed to the poſterior margin of the acromion.

Ligamentum proprium ſcapulae poſticum, ariſes from the middle of the ſuperior margin, and terminates at the root of the coracoid proceſs. Under this ligament the veſſels and nerves paſs to the ſhoulder.

The cartilage which covers the head of the os humeri is thick in its middle; but becomes gradually thinner towards its edges.

The four ſurfaces of the tuberoſities which appear cartilaginous in dry bones, ſerve only for the inſertion of the tendons of the four muſcles which move the os humeri on the ſcapula.

The channel through which the tendon of the long head of the biceps muſcle runs between the tuberoſities, is covered partly by a thin cruſt, which appears rather ligamentous than cartilaginous; and partly by a tendinous ſtratum.

Ligaments *between the* Scapula *and* Os Humeri.

Ligamentum capſulare humeri, ariſes from the whole margin of the glenoid cavity of the ſcapula, and is fixed round the under end of the neck of the os humeri, looſely incloſing the head of this bone. The upper part of the liga-

ment ſends down a ſheath between the two tuberoſities of the humerus, over the tendon of the long head of the biceps muſcle, which it accompanies as far as the fleſhy part and prevents it from ſliding out of the groove in which it is placed. The capſular ligament is ſtrengthened by other ligamentous bands which adhere firmly to its anterior ſurface; but what ſeems to give moſt force to the capſule are the tendons of the neighbouring muſcles, which increaſe its thickneſs conſiderably.

The trochlea, and ſmall head of the lower extremity of the os humeri, are covered by a common cartilage, in which the ſame proportion of thickneſs is obſervable as in that of the upper extremity. This is generally the caſe in all the convex articular cartilages. The foſſulae near the pulley are likewiſe covered with a kind of cartilaginous varniſh. The two ſigmoid cavities in the upper extremity of the ulna are covered by a cartilage common to both; which is a little interrupted, about the middle of the edges of the cavities, by the tranſverſe notches mentioned in the deſcription of the bones. This cartilaginous cruſt ſeems to be thicker at the edges than in the middle. The cartilage which covers the head of the radius is likewiſe turned over its cylindrical border; and a lateral portion of the muſcular tuberoſity, immediately below the neck, is alſo covered with a thin ſhining cartilage.

Ligaments *of the joint of the* Elbow.

Ligamentum capſulare, ariſes from the lower end of the os humeri, above the edge of the cartilaginous ſurface, and is fixed to the top of the ulna round the edge of the great ſigmoid cavity, including both the apex of the olecranon and that of the coronoid proceſs. It likewiſe runs over the head of the radius, and is fixed to the coronary ligament quite round. Thus it completely ſurrounds the articulation

culation of theſe three bones; and ſerves to contain the mucilaginous liquor in the cavity of the joint. It appears to be ſtrengthened by a ligamentous web; the fibres of which croſs each other in different directions: Beſides this, there are ſome tendinous fibres of muſcles to which the capſular ligament adheres very cloſely.

Near the under end of the body of the os humeri, there are two particular intermuſcular ligaments, which are long, narrow, and thin; but ſtrong, fixed by one end along the two lower thirds of the bone, and reaching to both condyles. They increaſe the ſurface for the origin of muſcles, and thereby ſupply the place of bones.

The lower extremity of the os humeri is alſo joined to the bones of the fore-arm by the two following faſciculi of ligamentous fibres.

Ligamentum brachio-cubitale ſeu laterale internum, ariſes from the fore-part of the internal condyle of the os humeri; and running down over the capſula, to which it cloſely adheres, is ſpread out in a radiated manner to be fixed to the inſide of the coronoid proceſs of the ulna. It is covered on the outſide by ſeveral tendons, which are connected cloſely to it, and ſeem to ſtrengthen it.

Ligamentum brachio-radiale ſeu laterale externum, is diſpoſed much after the ſame manner; but is of a greater extent. It is expanded from the external condyle of the os humeri, as from a centre, and is inſerted round the coronary ligament, and from thence down to the neck of the radius; and alſo in the neighbouring parts of the ulna. Through all this paſſage, it covers the capſular ligament, and is covered by ſeveral tendons, adhering cloſely to both.

Ligaments *joining the* Head *of the* Radius *to that of the* Ulna.

Ligamentum coronarium ſeu orbiculare. The head of the radius is joined to that of the ulna, and the following ligament ſurrounds the head of the radius, reaching from one ſide of the ſmall lateral ſigmoid cavity of the ulna to the other, in an arch which is about three quarters of a circle. It is very ſtrong, and comes near the ſolidity of a cartilage. The ſide next the radius is very ſmooth; and though it connects that bone cloſely to the ulna, yet it leaves it room enough to turn in the motions of pronation and ſupination.

Ligamentum obliquum, ariſes from the tubercle of the ulna, which gives riſe to the bracheus internus muſcle, and is inſerted into the tubercle of the radius.

Ligaments *between the* Bodies *of the* Radius *and* Ulna.

Ligamentum interoſſeum fills up the ſpace between the two bones of the fore-arm. It is fixed by one edge along the ſharp angle of the radius, and by the other along that of the ulna. The greater number of the fibres which compoſe it deſcend from the radius to the ulna. Some, however, aſcend and croſs the former obliquely, ſo as to make it appear as if compoſed of two planes. Small ſpaces are left in different parts of it for the paſſage of blood veſſels; and a large opening is left above, which is occupied by the oblique ligament, &c. The interoſſeous ligament ties the two bones together, and gives inſertion to muſcles. In the ſupination of the hand, it is ſtretched; and in the pronation, it is relaxed.

All the concave ſide of the baſis of the radius is cartilaginous, and often divided by a ſmall cartilaginous prominent line.

line. The lateral notch of the baſis is alſo covered by a continuation of the ſame cartilage.

At this end of the radius, there is likewiſe a particular additional cartilage, or triangular production, longer than it is broad, very thin, and rather flat than concave on both its ſmooth ſides. It is fixed by its baſis, or ſhorteſt ſide, to the lateral ſigmoid notch of the baſis of the radius, in ſuch a manner, that one ſide of it is on a level with the large cartilaginous ſurface of the baſis of the bone, and its apex directly oppoſite to the ſtyloid proceſs. The other ſide touches the flat extremity of the ſmall head of the ulna, but is not fixed to it. This cartilage may be termed the *interarticular cartilage* of the joint of the wriſt. It is tied to the radius by very ſhort ligaments; and, playing the ſmall head of the ulna, it follows all the motions of the radius. It is therefore a ſort of particular production of the lower ſide of the baſis of the radius, and fills, in the natural ſtate, the void ſpace which, in the ſkeleton, appears between the end of the ulna and the neighbouring bone of the carpus. The inferior extremity, or ſmall head of the ulna, is cruſted over by a cartilage round its cylindrical border, in the notch near the ſtyloid proceſs, and for ſome ſpace on the proceſs itſelf.

Ligaments *of the inferior extremity of the* Bones *of the* Fore-Arm.

Ligamentum capſulare, ariſes round the edge of the glenoid cavity of the lower end of the radius and head of the ulna. It is fixed to the cartilaginous edges of the three firſt bones of the carpus.

Ligamentum capſulare ſeu facciforme, ariſes from the edges of the ſemilunar cavity, at the under end of the radius, and ſurrounds the head of the ulna.

Ligamenta duo tranſverſa. One of theſe ariſes from the ſtyloid

ſtyloid proceſs at the under end of the radius, and is inſerted into the os naviculare. The other ariſes from the ſtyloid proceſs at the lower end of the ulna, and is fixed to the os cuneiforme and unciforme.

All the bones of the carpus, metacarpus, and fingers, are cruſted over with cartilages at the places which play upon each other; but in freſh ſubjects, they are thicker, ſofter, and whiter, than in the ſkeleton. In adults, their figure remains the ſame in both; but it changes in the dry bones of younger ſubjects; and in thoſe of children it is quite different. The impreſſions and notches in which the bodies called *mucilaginous glands* are lodged, are moſt ſenſible in the cartilages of freſh bones, becauſe of their thickneſs.

Ligaments *of the* Carpus.

Ligamenta oſſium carpi brevia, are ſmall ſhort ligaments, running in various directions, and joining the carpal bones; firſt of the ſame row, then of the two rows together. They have their names from their figure and the direction of their fibres; as *obliqua*, *tranſverſa*, *capſularia*, and *propria oſſium carpi*.

Ligamentum oſſium carpi commune capſulare, ariſes from the cartilaginous edges of the firſt row of carpal bones, and is inſerted into thoſe of the ſecond row.

Ligaments *between the* Carpus *and* Metacarpus.

Ligamenta articularia, ſhort firm ligaments, by which the ſecond ſeries of carpal bones are joined to the poſterior extremities of the metacarpal bones. On account of the variety in ſituation, and diverſity of the direction of their fibres, they have got the name of *Ligamenta dorſi manus*, *lateralia*, *recta*, *perpendicularia*.

Ligamenta interoſſea metacarpi, are ſmall ligaments which join

join the posterior and anterior extremity of the metacarpal bones together.

Ligaments *of the* Bones *of the* Fingers.

Ligamenta capsularia phalangum digitorum, arises from the anterior extremities of the metacarpal bones with the posterior extremities of the first phalanx of the fingers.

Ligamenta lateralia phalangum digitorum, are strong ligaments, which lie between the bones of the first phalanx of the fingers. They are fixed at each end to the capsular ligaments.

Ligamentum capsulare pollicis, arises from the posterior extremity of the first bone of the thumb, and is fixed round the os trapezium of the carpus.

Ligaments *retaining the* Tendons *of the* Muscles *of the* Hand *and* Fingers *in situ*.

Ligamentum carpi transversale externum, arises from the styloid process of the ulna and os pisiforme of the carpus, and, running transversely on the back of the wrist, it spreads broad to end in the styloid process of the radius. Between this ligament and the bones, the tendons of the extensor muscles of the carpus and fingers pass.

Ligamenta vaginalia, adhere to the former ligament and bones, and serve as a kind of sheaths to the tendons.

Ligamenta tendinum extensorum transversa, are short tendinous ligaments, running transversely on the back of the hand behind the roots of the fingers, and serving to join the tendons of the musculus extensor digitorum communis together.

Ligamenta palmaria transversa, are fixed to the anterior extremities of the metacarpal bones, from which they run transversely. In their passage, they cover the musculi lumbricales,

bricales, and are inſerted into the metacarpal bones and ſheaths of the tendons of the flexor muſcles.

Ligamenta vaginalia tendinum flexorum, ariſe from the internal tranſverſe ligament of the wriſt, and as ſheaths embrace the tendons of the flexor muſcles of the fingers; they terminate at laſt with the tendons of the muſculus perforans.

Ligamenta vaginalia ſeu cruciata phalangum, run in a circular and crucial direction over the former vaginae and tendons, and are fixed to the ridges on the concave ſide of the bones of the fingers. They ſerve as fraena to the tendons while their muſcles are in action.

Ligamentum tendinum flexorum acceſſoria, are ſmall but firm tendinous ſubſtances, which ariſe from the firſt and ſecond phalanx of the fingers; they are covered by the vaginal ligaments of the tendons, and terminate in the tendons of the two flexor muſcles of the fingers.

All the bones of the ſuperior extremities are covered with their perioſteum, and the quantity of marrow correſponds with the ſhape of the bone. All the joints have likewiſe ſynovial ſubſtances; but they are ſmall when compared with thoſe in the inferior extremities. The moſt conſiderable are placed in the cavities at the under end of the os humeri, for lodging the coronoid proceſs and olecranon of the ulna, in the flexion and extenſion of the fore-arm.

Of the PELVIS and INFERIOR EXTREMITIES.

The cartilages of the oſſa innominata are not ſo numerous as one might imagine on examining the ſkeleton. We are apt to think we ſee the dried remains of cartilages on the ſpine of the os ilium, on the tuberoſity of the os iſchium,

iſchium, and on the grooves and notches which give paſſage to the tendons of muſcles: but none of theſe incruſtations are true cartilages, being for the moſt part tendinous, aponeurotic, or ligamentous; which being dried, look more like cartilages than the true cartilages themſelves.

The cruſt which covers the ſpine of the os ilium is chiefly tendinous in adult bodies; but in children, and in very aged perſons, it appears cartilaginous. In children, the parts which are not completely offified are eaſily taken for true cartilages; and in old age, the tendons are often hardened to ſo great a degree, as to have the very ſame appearance. The ſubſtance which covers the tuberoſity of the os iſchium is almoſt entirely tendinous; and that which lines the grooves and notches of the tendons is chiefly ligamentous.

The true cartilages of the offa innominata are five in number; three common, and two proper. The firſt and principal common cartilage is that which makes the ſymphyſis of the offa pubis. It reaches from the interval between the ſpines of theſe two bones to the angle formed by the crura where they begin to ſeparate. It is ſomething thicker or broader at its upper part than for a conſiderable ſpace lower down; but the inferior part is by much the broadeſt. It fills the angle already mentioned, and forms a kind of arch, which is more conſiderable in women than in men.

The two other common cartilages join the offa ilium to the os ſacrum, but are thinner than that of the offa pubis.

The proper cartilages are thoſe that line the acetabula. Concerning theſe, it has been already obſerved in the deſcription of the Skeleton, that in the edge of each there is a notch or opening between the anterior and inferior parts; and that, in the cavity itſelf, there is a broad unequal ſhallow de-

preſſion for the ſynovial gland, reaching from the notch beyond the middle of the cavity. All the reſt of the ſurface is covered with a very white ſhining ſmooth cartilage, which terminates preciſely at the edge of the cavity.

The circumference of the acetabulum has, beſides, a border of a particular kind; the ſubſtance of which is neither wholly cartilaginous, nor wholly ligamentous; but it may be rather placed among the ligaments. The os ſacrum has no cartilage, excepting that between its upper end and the laſt vertebra of the loins, and thoſe by which it is connected with the oſſa innominata. The intervertebral cartilages of this bone are, for the moſt part, entirely obliterated in the adult. The cartilages which join the different portions of the os coccygis, are preſerved in ſome ſubjects to a very great age; in others they ſoon become entirely oſſified.

No part of the os femoris is covered with cartilage, excepting the uniform convexity of its head; and here the cartilage runs as far as the union between the head and neck of the bone. The trochanters have no true cartilage; what looks like it being only the remains of tendinous inſertions, as was obſerved of the ſpine of the os ilium. The cartilaginous ſubſtance which, to a certain age, unites the epiphyſes to the body of the bone, does not belong to this place, becauſe it is only found in the time of youth, and in adults is converted into bone. The cartilaginous matter by which the head of the os femoris is cemented, deſerves, however, to be obſerved; becauſe that epiphyſis has been ſeparated by violent falls.

Ligaments *proper to the* Bones *of the* Pelvis.

Ligamenta ileo-ſacra, are ſtrong ligaments ariſing from the poſterior part of the ſpine of the os ilium, which is oppoſite to the ſide of the os ſacrum, and deſcending obliquely,

liquely, are fixed to the firſt, third, and fourth ſpurious tranſverſe proceſſes of the os ſacrum.

Ligamenta pelvis tranſverſalia ſuperius et inferius, are two ligaments ariſing from the poſterior ſpinous proceſs of the os ilium; the ſuperior is fixed to the tranſverſe proceſs of the laſt vertebra of the loins; the inferior is fixed to the firſt tranſverſe proceſs of the os ſacrum.

Ligamenta ſacro-iſchiatica. Between the os iſchium and os ſacrum, we find two very ſtrong ligaments called *Sacro-ſciatic*; one broad and external, the other ſmall and internal. The external ariſes from the anterior and external edge of the falſe tranſverſe proceſſes of the os ſacrum. From thence diminiſhing in breadth, it deſcends obliquely towards the tuberoſity of the os iſchium, and is inſerted immediately below the ſinus, which lies between the tuberoſity and ſpine of that bone. This inſertion is afterwards continued over the whole internal labium of the inferior portion of the os iſchium, and of the crus of that bone, and the inferior portion of the crus of the neighbouring os pubis. When it arrives at the os iſchium, it produces a kind of falx; one edge of which is fixed to the bones, the other lies looſe; and by this inſertion of the falx, it forms, together with the bones, a kind of deep channel or groove.

The internal ſacro-ſciatic ligament adheres cloſely to the inſide of the poſterior portion of the former. It ariſes internally from the edge of the inferior part of the fourth falſe tranſverſe proceſs, and from the whole ſide of the os ſacrum, and from the baſis of the upper part of the os coccygis. From this it runs up a little obliquely to the ſpine of the os iſchium; to the ſharp point and upper part of which it is fixed.

By theſe two ligaments two openings are formed; a large one, with the ſuperior ſciatic ſinus, through which the pyriform muſcle, the poſterior crural veſſels, and the ſciatic nerve,

nerve, pafs out of the pelvis; and a fmall one for the paſſage of the internal obturator muſcle.

Ligamentum obturans foraminis ovalis. The obturator ligament fills up all the foramen thyroideum, except the oblique notch at its upper part for the paſſage of the obturator veſſels and nerves. It is fixed to the edge of that hole from the anterior part of the oblique notch, as far as the ſymphyſis between the os pubis and os iſchium. From thence to the poſterior part of the inferior notch, it is fixed to the internal labium of the edge of the circumference, forming a kind of ſmall channel with the external labium; and afterwards it is fixed to the common edge of the foramen ovale and cotyloid notch. This ligament not only aſſiſts in ſupporting the parts contained in the pelvis, but alſo gives origin to the two obturator muſcles.

On the inſide of the upper and anterior part of the os pubis, there is a tranſverſe ligament fixed by its upper part to the os pubis, from the oblique notch of the foramen ovale, all the way to the lower part of the ſymphyſis, at a ſmall diſtance from the circumference of the laſt-mentioned hole. This ligament is about half an inch broad in an adult body, and, poſteriorly, below the ſuperior oblique notch of the foramen ovale, it joins the obturator ligament by means of a particular fold; and by parting from it afterwards, a deep narrow groove is formed between them; the tranſverſe ligament being at this place ſupported by ligamentous fraena of different ſizes.

Ligamentum inguinale, ſeu Poupertii, ſeu Fallopii, the inguinal ligament, is chiefly the under end of the tendon of the external oblique muſcle of the abdomen. It is fixed by one end to the anterior ſuperior ſpinous proceſs of the os ilium, and is ſtretched over to be fixed by its other end to the ſpine of the os pubis. The middle portion of it is very narrow, but expands conſiderably towards both extremities.

Under

Under this ligament the femoral veſſels and anterior crural nerve go out of the pelvis.

Ligamentum capſulare coccygis, ariſes from the upper end of the os coccygis, and is inſerted round the under end of the os ſacrum.

Ligamenta longitudinalia coccygis, ſmall ligaments ariſing from the inner ſurface of the os coccygis, and terminating in the os ſacrum. They fix the two bones firmly together.

Ligaments *between the* Pelvis *and Head of the* Os Femoris.

The capſular ligament is the moſt conſiderable, largeſt, and ſtrongeſt, of all the articular ligaments of the human body. It ariſes quite round the outer edge of a thick ſtrong cartilago ligamentous border, on the brim of the acetabulum, and from thence largely ſurrounds the whole head and ſuperior portion of the neck of the os femoris, and is cloſely inſerted to the lower portion of the neck that is between its baſis and middle narrow part. This ligament is made up of ſeveral ſorts of fibres, the chief of which are longitudinal and oblique; and it is much thicker and ſtronger in ſome parts than in others. It is very thick on its fore-part, on account of two ligamentous bands which run downwards and outwards from the inferior anterior ſpinous proceſs of the os ilium to the further extremity of the neck of the thigh bone. It is ſomewhat thinner on its outer and back-part, and thinneſt of all at the inner and back-part.

Ligamentum teres ſeu rotundum. This ligament is not round as the name expreſſes; it reſembles a flat cord, being compoſed of a bundle of fibres cloſely interwoven; one end of it is in a manner divided into two flat bands, which are fixed to the inner corners of the notch of the acetabulum,

lum, and alſo to the edge of the rough impreſſion at the bottom of the acetabulum. From the inſertion it runs obliquely backwards and a little upwards, between the ſynovial gland within the acetabulum and the cartilaginous convexity at the head of the os femoris, and ends in the upper part of the ſmall ſemilunar notch. This inſertion is oblique, a little rounded on the upper part, and flat on the lower; and in ſome ſubjects there is a ſort of depreſſion in the head of the bone for the paſſage of the ligament.

The perioſteum of the bones of the pelvis agrees with that on other flat bones.

The rough unequal depreſſion at the bottom of the acetabulum is filled with a broad ſynovial gland, bordered with a fatty ſubſtance, and covered by a fine membrane, through which a mucilaginous liquor paſſes to moiſten the joint and facilitate its motions. This membrane riſes above the gland, and gives a ſort of covering to the ligament contained in the joint.

The blood-veſſels of this gland paſs between the bottom of the acetabulum and the ligament at the inner edge of that cavity.

The blood-veſſels paſs chiefly through the ſmall holes in both convex and concave ſurfaces of theſe bones; and ramifying upon the bony cells, they end in a great number of ſmall capillary tubes, which make the medullary juice appear red.

The cartilage which covers the lower extremity of the femur is exactly fitted to the ſemi-oval convexity of the inferior ſurface of each condyle, and to the pulley formed by their union.

The two cartilages which cover the two ſuperior ſurfaces of the head of the tibia are gently hollow; but the internal is more depreſſed than the external: The back-part of the

the latter is ſenſibly depreſſed, by which a ſort of convexity is formed. The cartilage is abſent where the crucial ligament of the patella has a thick cartilage on its poſterior ſide, divided by a ſuperficial longitudinal riſing, proportioned to the two portions of the pulley of the os femoris; at the outer and under part of the head of the tibia another cartilage appears, where the head of the fibula is articulated. The cartilage at the head of the fibula ſeems to be thicker than that at the lower extremity.

The ſemilunar cartilages, which get their name on account of their figure, are thick at their outer edge where they are tied to the capſular ligament; thin at their inner edge, particularly at the middle; concave ſuperiorly next the condyles of the femur; flatted below next the tibia; and tied together by a ſmall ligament. Theſe cartilages are in ſhape of a creſcent, or Roman C. Their convexity or greateſt curvature is very thick; their concavity or ſmalleſt curvature very thin; ſomething like the edge of a ſickle. They lie on the two upper ſurfaces of the head of the tibia; their thickeſt part or convexity correſponding with the edges of the head, and their thin ſharp edges to the middle of each ſurface; their extremities or cornua being turned toward each other. Each cartilage covers about two thirds of the ſurface of the tibia on which it lies, leaving one third bare in the middle. Their under ſides are flat, the upper ſides hollow; and, together with the middle portions of the ſurfaces of the head of the tibia, form cavities proportional to the convexities of the condyles of the os femoris.

Ligaments *within the* Joint.

Ligamenta cruciata. One of theſe, called *poſterius*, is fixed to the internal ſuperficial impreſſion in the notch between the condyles of the femur; and running almoſt ſtraight

ſtraight down, is fixed by its other end to the notch in the head of the tibia, behind the cartilaginous tubercle which lies between the two ſuperficial ſurfaces for receiving the condyles of the femur.

The other, called *anterius*, is fixed by one end to the external impreſſion in the notch of the os femoris; and running obliquely downwards and forwards, croſſing the former, is fixed by the other end to the head of the tibia a little before the other ligament. Theſe two ligaments croſs each other when we turn the point of the foot inwards, and they ſeparate from each other when the foot is turned outwards. They prevent the leg from being bent forwards on the thigh, and from rolling too much inwards.

Ligamentum alare majus et minus: are two broad ligaments ariſing from the inner ſides of the capſular ligament, and are fixed to the ſides of the patella, and to the fatty ſubſtance placed there.

Ligaments *at the* Joint *of the* Knee.

The internal is fixed to the femur under the tuberoſity, which is near its internal condyle; the fibres ſpread out in deſcending, and terminate at the upper and inner part of the tibia; along which they ſlide from behind forwards, till they have got more than two inches and a half under the head of this bone.

The external is narrower and ſhorter. It is fixed above to the external tuberoſity of the femur, and deſcends to embrace the anterior part of the head of the fibula, where it enlarges a little, though its fibres are not radiated like thoſe of the external ligament. Its length is about two inches, and it is ſomewhat looſe. Theſe two ligaments are not placed in the middle of the articulation; they are a little farther back, ſo as to allow the ready flexion and extenſion

tenſion of the limb. They are quite looſe in the flexion, and put upon the ſtretch in the extenſion of the limb.

Ligamentum poſticum, has an irregular form. It deſcends from the poſterior, inferior, and external part of the outer condyle of the femur; and having croſſed the poſterior part of the articulation, it terminates in the poſterior, ſuperior, and internal part of the tibia. Some fibres go by the ſide of the former, to end in the upper and back part of the tibia. The three ligaments above deſcribed are firmly glued to the capſular ligament.

The capſular ligament is fixed quite round the inferior extremity of the os femoris, at a ſmall diſtance above the anterior, lateral, and poſterior parts of the cartilage, and above the poſterior part of the great notch; from this it runs down to be fixed round the edge of the head of the tibia and in the edge of the patella, in ſuch a way that the patella itſelf forms a part of the capſula of the joint. It is ſtrengthened by the tendinous aponeuroſis and tendons of the muſcles which ſurround the joint, and likewiſe by the ligaments already mentioned; internally, it adheres to the ſemilunar cartilages, and ſends off a very fine vagina over the ligaments, &c. within the joint.

Ligamenta cartilaginea, are two ſmall ligaments which join the ſemilunar cartilages to each other, and likewiſe to the os femoris and tibia.

Ligamentum patellae, is a very ſtrong ligament which ariſes from the point of the patella, and is fixed to the upper and fore-part of the tibia.

The marrow of the os femoris lies in a large maſs in the middle cavity of the bone, and in ſmall diſtinct cluſters in the cells of each extremity. The firſt is penetrated at different diſtances by the bony filaments or ramifications of the reticular texture, and thereby ſuſtained in the violent motions of the body.

The ſynovial ſubſtances of the knee, which lie near the edges of the patella, are the moſt conſiderable of any in the body; being diſpoſed in form of fringes, and ſupported by a great quantity of fatty matter.

This common maſs is contained within the capſular ligament; and on the ſide of the joint is covered by a very fine membrane which likewiſe lines the inner ſurface of the ligament.

The ſuperior portion of this fat is as it were ſupported by a ſmall ligament fixed in the anterior part of the great notch between the condyles of the femur, and which runs to the upper part of the patella.

There are other fatty ſubſtances both above and below the edges of the ſemilunar cartilages, and likewiſe in the ham; ſome of which ſerve for the joint, the reſt for the crucial ligaments. Theſe laſt lie in folds formed by the internal membrane of the capſular ligament, which give particular coverings to the crucial ligaments, and to the other bundles of ligamentous fibres near them.

Ligaments *of the* Fibula.

Ligamentum capſulare extremitatis ſuperioris, is a very ſtrong ligament, which runs from the head of the fibula to be fixed to the external condyle of the tibia.

Ligamentum interoſſeum, fills up the ſpace between the tibia and fibula. One ſide of it is fixed to the poſterior external angle of the tibia, the other to the internal angle of the fibula. It is compoſed of ſtrong tendinous fibres, which croſs each other obliquely, and in various parts leave ſmall ſpaces for the paſſage of veſſels and nerves. At its upper part there is a large opening, where the veſſels and nerves paſs to the fore-part of the leg, and where the muſcles on oppoſite ſides are contiguous. It ſerves chiefly as a ligamentous

mentous ſeptum for the origin of muſcles; in which reſpect it ſupplies the place of bones.

Ligamentum extremitatis inferioris, conſiſts of four ſhort ſtrong ligaments, two of which are anterior and two poſterior; they ariſe from the malleolus externus of the fibula, and are inſerted into the under and outer part of the tibia.

The marrow of theſe bones lies in large maſſes in the great cavities, and in diſtinct moleculae in the ſpongy parts, as in other bones of the ſame ſhape. The ſynovial glands lie in the ſmall ſpaces, depreſſions, and ſuperficial notches; near the edges of the cartilages of each joint they are covered by the capſular ligaments, and more or leſs mixed with a fatty ſubſtance.

Ligaments *of the* Inferior Extremity *of the* Bones *of the* Leg.

Ligamentum fibulae anticum, ariſes from the fore-part of the malleolus externus of the fibula, and is fixed to the upper and outer part of the aſtragalus.

Ligamentum fibulae medium, ariſes from the point of the malleolus externus, and runs ſtraight down to be fixed to the outſide of the os calcis.

Ligamentum fibulae poſticum, ariſes from the under and back-part of the malleolus externus, and is bent obliquely backwards to be fixed to the outer and back-part of the aſtragalus.

Ligamentum tibiae deltoideum, ariſes from the malleolus internus, and is fixed to the aſtragalus and os naviculare.

Ligamentum capſulare, ariſes from the whole edge of the articular cavity of the tibia, and is fixed entirely round the aſtragalus.

Ligaments *of the* Bases *of the* Metatarsal Bones.

Ligamentum capsulare, joins the metatarsal bone of the great toe to the os unciforme internum.

Ligamenta articulatoria, join the posterior extremities of the metatarsal to the anterior edges of the tarsal bones. On account of their different situations, they have the name of *ligamenta plantaria, pedis dorsalia, lateralia.*

Ligamenta transversa dorsi pedis, are three in number, and are situated on the upper part of the foot, and join the bases of all the metatarsal bones together, excepting that which belongs to the great toe.

Ligamenta transversa plantae are likewise three, but placed in the sole; and lie partly in the interstices of the bones.

Ligamenta interossea metatarsi, like the former, are also three in number, but run immediately from the side of one bone to that of the bone next it, filling up the interstitial spaces.

Ligaments *of the* Bones *of the* Toes.

Ligamenta capsularia, arises from the posterior extremities of the first phalanx of the bones of the toes, and are fixed to the anterior extremities of the metatarsal bones.

The second and third phalanges of all the toes are joined by capsular ligaments, much in the same manner with those already mentioned.

Ligamenta lateralia, lie at the sides of the second and third joint of each of the toes; they arise from the sides of the bases, and are fixed to the sides of the heads of the bones of each of these joints.

All these ligaments, in the same manner as in the hand, are covered and strengthened on the dorsum pedis by an expansion of the tendons of the extensor muscles, and in the sole by the sheaths of the tendons of the flexor muscles.

LIGAMENTS *retaining the* TENDONS *of the* MUSCLES *of the* FOOT *and* TOES *in ſitu.*

Ligamentum vaginale tibiae. This ſtrong ligament is part of the tendinous ſheath that covers the muſcles on the fore-part of the leg; one ſide of it is fixed to the anterior angle at the lower part of the tibia, the other is fixed to the outer part of the fibula.

Ligamentum tranſverſum tarſi ſeu cruciatum. This is a double ligament; one part ariſes in the outer part of the foot above the malleolus externus of the fibula, the other from the proceſs of the os calcis. They run over, the one to be fixed to the malleolus internus, the other to the inner ſide of the os naviculare; they ſerve to keep the tendons of the muſcles in their places.

Ligamentum tendinum peroneorum, ariſes from the fore-part of the os calcis, and is fixed to the outer ſide of an eminence of this bone.

Ligamentum laciniatum, ariſes from the edge of the malleolus internus, and runs down in a radiated manner to be loſt in the fat ſituated there, then in the membrana propria of the abductor muſcle of the great toe, and at laſt in the inner ſide of the os calcis; it covers the tendons, veſſels, and nerves, running in the hollow of the os calcis.

Ligamentum vaginale extenſoris pollicis, ſurrounds the tendon of this muſcle.

Ligamentum vaginale flexoris proprii pollicis, ſurrounds the tendon of the flexor longus pollicis in the hollow of the os calcis; it is fixed to the under and inner part of this bone.

Ligamenta vaginalia tendinum flexorum. Theſe are ligamentous ſheaths ſurrounding the tendons of the flexor muſcles of the toes, and are ſimilar to thoſe of the flexors of the fingers.

Ligamenta

Ligamenta acceſſoria tendinum flexorum, are ſhort, but ſtrong ligaments, like thoſe on the fingers, ariſing from the phalanges of the toes; and being included in the ſheaths of the tendons, they terminate in the tendons.

Ligamentum tendinum extenſorum tranſverſa, are ſmall ligaments running between the tendons of theſe muſcles, and ſerving to bind them together, and to keep them in their places.

The perioſteum of all theſe bones is of the ſame kind with that of the bones of the leg.

The marrow is ſuitable to their internal ſtructure; that is, in moleculae in the cavernous portions, and in maſſes in thoſe which have large cavities. Thus the marrow of all the tarſal bones is diſperſed in moleculae, becauſe their internal ſtructure is ſpongy. In the metatarſal bones and firſt phalanges of the toes, it is diſpoſed in the ſame manner as in the tibia and fibula; that is, it lies in moleculae in the extremities, the ſtructure of which is cavernous; but in the middle portions of them it lies in maſſes greater or leſs, according to the ſize of the cavities. In the other phalanges, which are entirely ſpongy, it is accordingly diſpoſed in moleculae.

The ſynovial ſubſtances anſwer in number and figure to the depreſſions between the cartilaginous edges and ligaments.

A

SYSTEM OF ANATOMY.

PART V.

Containing a DESCRIPTION of the

COMMON INTEGUMENTS.

ALL the parts of the human body are inveſted by ſeveral common and univerſal coverings, to which anatomiſt give the name of *integuments*.

There have been many diſputes about the number of theſe integuments. The ancients reckoned up five, viz. the *epidermis*, *ſkin*, *membrana adipoſa*, *panniculus carnoſus*, and *membrana muſculorum communis*.

The three firſt of theſe coverings are truly common or univerſal, that is, extended over all parts of the body.

The two other coverings are not univerſal, but confined to particular parts of the body.

The moderns divide the integuments into *cuticula*, *rete mucoſum*, *cutis vera*, and *corpus adipoſum*.

CUTI-

Cuticula.

The outside of the skin is covered by a thin transparent insensible pellicle, closely joined to it, which is called *epidermis*, *cuticula*, or *scarf-skin*.

The substance of the cuticle appears to be very uniform on the side next the skin, and to be composed on the other side of a great number of very fine small squamous laminae, without any appearance of fibrous or vascular texture, unless we take into account the numerous small vessels by which it is connected to the parts under it.

This substance is very solid and compact, but yet capable of being extended and thickened, as we see by steeping it in water, and by the blisters raised on the skin by vesicatories or any other means; and from thence it would seem that it is of a spongy texture. It yields very much in swellings, but not so much as the skin without breaking or cracking. It is not readily destroyed by putrefaction.

Hard and reiterated frictions loosen it insensibly, and presently afterwards a new stratum arises, which thrusts the first outward, and may itself be loosened and thrust outward by other strata.

It is nearly in this manner that callosities are formed on the feet, hands, and knees; and the several laminae or strata observable at the same time on many other parts of the body, are owing to the same cause, though many anatomists have thought them to be natural. But it must be acknowledged, that, on the palms of the hands and soles of the feet, the cuticle is commonly thicker than on any other part.

The cuticle adheres very closely to the cutaneous papillae under it; but it may be separated by boiling, or steeping for a long time in cold water. It is not impossible to separate

parate it with the knife; but this management teaches us nothing of its ſtructure.

It adheres ſtill cloſer to the corpus mucoſum, which is eaſily raiſed along with it; and they ſeem to be true portions or continuations of each other.

The colour of the cuticle is naturally white; and its apparent colour is owing to that of the corpus mucoſum. For, if we examine the European and African, we find the cuticle to be nearly of the ſame colour in both, whereas their corpus mucoſum is very different.

The cuticle covers the ſkin through its whole extent, excepting at the nails. It is marked with the ſame furrows and lozenges as the ſkin, and has the ſame openings and pores, and may be ſaid to cover not only the whole external part of the body, but to line many of the large paſſages, as the alimentary canal, the lungs, vagina, urethra, &c. In theſe paſſages, however, it is ſomewhat different. The cuticle on the external parts of the body gives paſſages to the ducts of the ſebaceous glands, while that lining the cavities already mentioned, is pierced with the ducts of the mucous follicles.

When we examine narrowly the ſmall paſſages through which the ſebaceous matter of the ſkin paſſes, the cuticle ſeems to enter theſe, in order to complete the ſecretory tubes. The foſſulae of the hairs have likewiſe the ſame productions of the cuticle, and it ſeems to give a kind of covering to the hairs themſelves. Laſtly, the almoſt imperceptible ducts of the cutaneous pores are lined by it.

If the ſkin be macerated for a long while in water, the cuticle, with its elongations, may be ſeparated from it. By this obſervation we may explain how bliſters may remain for a long time on the ſkin without giving paſſage through theſe holes to the matter which they contain;

which holes one would think ought to be increaſed by this dilatation and tenſion of the cuticle.

But when the cuticle is ſeparated from the ſkin, it carries along with it part of theſe cutaneous fibres; which being compreſſed by the matter contained in the bliſter, ſhut the pores of the ſeparated cuticle like ſo many valves; and it is probable, that ſome of theſe ſmall portions being detached, are the ſubſtances which have been taken for valves of the cutaneous tubes.

With reſpect to its origin, ſome authors have ſuppoſed it to be formed by a moiſture exhaled from the whole ſurface of the body, which gradually hardens when it is expoſed to the air; but the foetus in utero, where no air is admitted, is a proof againſt this opinion; and it grows readily under plaſters applied to any part of the body. Leeuwenhoeck ſuppoſed its formation to be owing to the expanſion of the extremities of the excretory veſſels, which are found every where upon the ſurface of the true ſkin. Ruyſch attributed its origin to the nervous papillae of the ſkin; and Heiſter thinks it probable that it may be owing both to the papillae and the excretory veſſels. Morgagni, on the other hand, contends, that it is nothing more than the ſurface of the cutis, hardened and rendered inſenſible by the liquor amnii in the uterus, or by the preſſure of the air. In fact, we know little about its origin; but the regeneration of it is very evident, ſudden, and ſurpriſing; for, let it be deſtroyed ever ſo often, it ſtill grows again.

Corpus Mucosum.

Under the cuticle, we meet with a ſubſtance of a greyiſh colour, which has been thought to repreſent a net-work; hence it has got the name of *corpus reticulare*, or *mucoſum*. It is of a ſoft, mucilaginous, and viſcid nature; and fills up the interſtices of the fibres running between the cutis

vera

vera and cuticula. After raiſing the cuticle in a negro, where it is thickeſt and moſt diſtinct, this ſubſtance appears of a black colour, and is compoſed of two layers. It is this that chiefly gives the colour to the ſkin; for it is black in the African; white, brown, or yellowiſh, in the European.

The origin of this mucous ſubſtance has not hitherto been ſufficiently explained; nor has it been fully determined what purpoſes it ſerves in the hman body. Haller thinks it probable, that it is compoſed of a humour tranſuding from the ſurface of the cutis vera. The reaſon why it is black in the negro has been ſuppoſed to be for ſerving as a defence againſt the external heat, by preventing the rays of the ſun from penetrating his body; but the matter ſtill lies hid in obſcurity.

Cutis Vera.

The cutis vera, or *ſkin* properly ſo called, is a ſubſtance of very large extent, made up of ſeveral kinds of fibres, cloſely connected together, and running in various directions, being compoſed of the extremities of numerous veſſels and nerves.

This texture is what we commonly call *leather;* and it makes, as it were, the body of the ſkin. It is not eaſily torn; may be elongated in all directions, and afterwards recovers itſelf, as we ſee in fat perſons, in women with child, and in ſwellings: and it is thicker and more compact in ſome places than in others.

Its thickneſs and compactneſs are not, however, always proportional: For on the poſterior parts of the body it is thicker and more lax than on the fore-parts; and on the palms of the hands, and ſoles of the feet, it is both very thick and very ſolid. It is generally more difficult to be pierced by pointed inſtruments in the belly than in the back.

The

The outer ſurface of this ſubſtance is furniſhed with ſmall eminences which anatomiſts have thought fit to call *papillae*. They appear through its whole ſurface like ſmall granulations; and ſeem to be calculated to receive the impreſſions of touch, being the moſt eaſily obſerved where the ſenſe of feeling is the moſt delicate, as in the points of the fingers and palms of the hands; and are ſuppoſed by many to be the capillary filaments of the cutaneous nerves, which terminate by ſmall radiated pencils: But they muſt be allowed to be formed like the other parts of the cutis; only the nervous fibrillae will be found to be more numerous in them than in other parts.

Theſe papillae differ very much in figure and diſpoſition in the different parts of the body; and they may be diſtinguiſhed into ſeveral kinds.

The greateſt part of them is flat, of different breadths; and ſeparated by ſulci, which form a kind of irregular lozenges. The pyramidal figure aſcribed to them is not natural; and appears only when they are contracted by cold, by diſeaſes, by boiling, or by ſome other artificial preparation, which alters their ordinary ſtructure.

The papillae of the palm of the hand, of the ſole of the foot, and of the fingers and toes, are higher than on the other parts of the body; but they are likewiſe ſmaller, cloſely united together, and placed as it were end-wiſe with reſpect to each other, in particular rows, which repreſent on the ſkin all kinds of lines, ſtraight, crooked, waving, ſpiral, &c. Theſe ſeveral lines are often diſtinctly viſible in thoſe parts of the palm of the hand which are next the firſt phalanges of the fingers.

The red part of the lips is made up of papillae, repreſenting very fine hairs or villi cloſely united together.

There is another particular kind under the nails; the papillae being there more pointed, or in a manner conical, and

and turned obliquely toward the ends of the fingers. Thoſe which are found in the hairy ſcalp, ſcrotum, &c. are ſtill of other kinds.

In inflammations, we obſerve a reticular texture of capillary veſſels, more or leſs extended on the ſurface of the ſkin; and curious anatomiſts demonſtrate the ſame thing by fine injections, which may be looked upon as artificial inflammations. Something ſimilar to this has been injected lately by Mr Baynham of London, who thought it *rete mucoſum*; and afterwards by Mr Cruickſhank, who calls it *cuticula quarta*. (See Mr Cruickſhank's letter to Mr Clare.) But neither of theſe methods prove, that, in the natural ſtate, theſe veſſels are blood-veſſels; that is, that they contain the red portion of the blood.

It is more probable that this vaſcular texture is only a continuation or production of the very ſmall capillaries of the arteries and veins; which, in the natural ſtate, tranſmit only the ſerous part of the blood, while the red part continues its courſe through wider ramifications, which more properly retain the name of *blood-veſſels*.

This vaſcular texture is of various forms and figures in the different parts of the body. It is not the ſame in the face as it is elſewhere; neither is it alike on all the parts of the face, as may be diſcovered by the moſt ordinary microſcopes; and from hence we might perhaps be enabled to give a reaſon why one part of the body turns red more eaſily than another.

The ſkin has ſeveral conſiderable openings, ſome of which have particular names; ſuch as, the fiſſure of the palpebrae, the nares, the mouth, the external foramen of the ears, the anus, and openings of the parts of generation.

Beſides theſe, it is perforated by an infinite number of ſmall holes, called *pores*, which are of two kinds. Some

are

are more or leſs perceivable by the naked eye; ſuch as the orifices of the milky ducts of the mammae, the orifices of the excretory canals of the cutaneous glands, and the paſſage of the hairs.

The other pores are imperceptible to the naked eye, but viſible through a microſcope; and their exiſtence is likewiſe proved by the cutaneous tranſpiration, and by the effects of topical applications; and from theſe two phenomena, they have been divided into arterial and venal pores.

We ought likewiſe to obſerve the adheſions and folds of the ſkin. It is every where united to the corpus adipoſum; but it adheres to it much more cloſely in ſome parts than in others, as in the palm of the hand, ſole of the foot, elbow, and knee.

Some plicae or folds in the ſkin depend on the ſtructure of the membrana adipoſa or cellularis, as thoſe in the neck and buttocks; others do not depend on that membrane, ſuch as the rugae in the forehead, palpebrae, &c. which are formed by cutaneous muſcles, and diſpoſed more or leſs in a contrary direction to theſe muſcles. Theſe folds increaſe with age.

There is, beſides, a particular kind of folds in the ſkin of the elbow, knee, and condyles of the fingers and toes; which are owing neither to the conformation of the membrana adipoſa, nor to any muſcle.

Laſtly, there is a kind of plicae, or rather lines, which croſs the palm of the hand, ſole of the foot, and correſponding ſides of the fingers and toes, in different directions. Theſe ſerve for employment to fortune-tellers; whoſe pretended art is contrary to religion, and deſpiſed by all men of ſenſe.

Glands of the Skin.

In different parts of the body, we meet with ſmall glands or follicles of an oval form, and ſeated chiefly under the ſkin in the corpus adipoſum.

They are compoſed of convoluted veſſels; but in ſome parts of the body they appear to be formed of ſmall cylindrical tubes, or ſimple follicles, continued from the ends of the arteries, and diſcharging, by ſmall excretory ducts, a fat and oily matter, that ſerves to lubricate and moiſten the ſurface of the ſkin. When the fluid they ſecrete has acquired a certain degree of thickneſs, it approaches to the colour and conſiſtence of ſuet: and from this appearance they have derived the name of *ſebaceous glands*. They are found chiefly on the noſe, ears, axillae, likewiſe round the nipple, and about the external parts of generation, in both ſexes.

Beſides the ſebaceous glands, anatomical writers mention other ſmall ſpherical bodies placed every where over the ſurface of the body, in much greater abundance than thoſe juſt mentioned, and named *miliary*. They are ſaid to have excretory ducts that open on the ſurface of the cuticle, and diſtil the ſweat and matter of inſenſible perſpiration; but after all that has been ſaid by different authors about them, it is certain there are no diſtinct glands that can be traced by the knife.

Uses of the Skin.

It is chiefly and properly the filamentary ſubſtance, called the *body of the ſkin*, which is the univerſal integument of the body, and the baſis of all the other cutaneous parts; each of which has its particular uſes.

The ſkin is able to reſiſt external injuries to a certain degree, and ſuch impreſſions, frictions, ſtrokes, &c. to which the

the human body is often liable, as would hurt, wound, and diſorder the parts of which it is compoſed, if they were not defended by the ſkin.

The papillae are the organs of feeling, and contribute to an univerſal evacuation, called *inſenſible tranſpiration*. They likewiſe ſerve to tranſmit from without, inwards, the ſubtle particles or impreſſions of ſome things applied to the ſkin. The firſt of theſe three uſes depends on the extremities of the nerves; the ſecond, on the arterial productions; and the third, on the origin of the abſorbent ſyſtem.

The cutaneous glands ſecrete an oily humour of different conſiſtencies. But without the epidermis, both papillae and glands would be diſturbed in their functions; on which great diſorders muſt enſue.

The epidermis ſerves to keep the pencils or nervous filaments of the papillae in an even ſituation, and without confuſion; and it likewiſe moderates the impreſſions of external objects. Particular, as well as general feeling, is more or leſs perfect, in proportion to the thinneſs of the epidermis; calloſities in which weaken, and ſometimes deſtroy both.

Another uſe of the epidermis is, to regulate the cutaneous evacuations already mentioned; the moſt conſiderable of which is inſenſible tranſpiration.

Membrana Adiposa, and Fat.

The laſt univerſal integument of the human body, is the membrana adipoſa, or corpus adipoſum. This is not, however, a ſingle membrane, but a congeries of a great number of membranous laminae, joined irregularly to each other at different diſtances, ſo as to form numerous interſtices of different capacities, which communicate with each other

other. These interstices have been named *cellulae*, and the substance they compose the *cellular substance*.

The thickness of the membrana adiposa is not the same all over the body, and depends on the number of laminae of which it consists. It adheres very closely to the skin; runs in between the muscles in general, and between their several fibres in particular; and communicates with the membranes which line the inside of the thorax and abdomen.

This structure is demonstrated every day by butchers, in blowing up their meat when newly killed; in doing which, they not only swell the membrana adiposa, but the air insinuates itself likewise in the interstices of the muscles, and penetrates even to the viscera, producing a kind of artificial emphysema.

These cellular interstices are so many little bags or satchels, filled with an unctuous or oily juice, more or less liquid, which is called fat; the different consistence of which depends not only on that of the oily substance, but on the size, extent, and subdivision of the cells.

It is generally known, that the illustrious Malpighi took a great deal of pains about this substance; that in birds and frogs, the viscera and vessels of which are transparent, he thought he saw a kind of ductus adiposi; and that, by pressing these ducts, he observed oily drops to run distinctly into the small ramifications of the mesenteric veins; such ducts, however, have not been seen by later anatomists.

The manufacture of soap, the composition of some of the ointments, and the different mixtures of oils with saline and acid liquors, give us some idea, at least, of the formation of the fat in the human body; but the organ which separates it from the mass of blood, which ought to be the

ſubject of our preſent inquiry, is not yet ſufficiently known.

Fat is more fluid in living than in dead bodies. It melts with the heat of the fingers in handling it; and its fluidity is in part obſtructed by the ſacculi which contain it. The method to take it entirely out of theſe bags is, to ſet the whole over a fire in a proper veſſel; for then the bags burſt, and ſwim in cluſters in a true oily fluid.

This ſubſtance increaſes in quantity by reſt and good living: and, on the contrary, diminiſhes by hard labour and a ſpare diet. Why nouriſhment ſhould have this effect, is eaſily conceived; and it is likewiſe eaſy to ſee, that an idle ſedentary life muſt render the fat leſs fluid, and conſequently more liable to block the paſſages of inſenſible tranſpiration, through which it would otherwiſe run off.

Hard labour diſſolves it, and conſequently fits it for paſſing out of the body, with the other matter of inſenſible tranſpiration. Some authors are of opinion, that it returns into the maſs of blood by the lymphatics; and that it can, for ſome certain time, ſupply the want of nouriſhment.

By this, they think, the long abſtinence of ſome animals may be explained; but it would appear, that the mere decreaſe of cutaneous tranſpiration, occaſioned by the continual reſt and inaction of theſe animals, has a great ſhare in this effect.

The proportional differences, in the thickneſs of this membrana adipoſa, are determined, and may be obſerved, to be regular in ſome parts of the body, where either beauty or uſe require it.

Thus we find it in great quantities where the interſtices of the muſcles would otherwiſe have left diſagreeable hollow or void places; but the ſkin being filled, and as it were

ſtuffed

ſtuffed with fat, is raiſed, and an agreeable form given to the parts.

The appearance of a perſon moderately fat, of a perſon extremely lean, and of a dead carcaſe from which all the fat has been removed, proves ſufficiently what has been ſaid.

In ſome parts of the body the fat ſerves for a cuſhion, pillow, or matreſs; as on the buttocks, where the laminae and cells are very numerous. In other parts, this membrane has few or no laminae, and conſequently little or no fat; as on the forehead, elbows, &c.

In ſome places it ſeems to be braced down by a kind of natural contraction in form of a fold; as in that fold which ſeparates the baſis of the chin from the neck, and in that which diſtinguiſhes the buttocks from the reſt of the thigh. We obſerve it likewiſe to be entirely ſunk, or as it were perforated by a kind of dimple or foſſula, as in the naval of fat perſons.

Theſe depreſſions and folds are never obliterated, let the perſon be ever ſo fat; becauſe they are natural, and depend on the particular conformation of the membrana adipoſa, the laminae of which are wanting at theſe places.

The fat is likewiſe of great uſe to the muſcles in preſerving the flexibility neceſſary for their actions, and in preventing or leſſening their mutual frictions. This uſe is of the ſame kind with that of the unctuous matter found in the joints, which was explained in the deſcription of the freſh bones.

Laſtly, the fat, as a fine oily ſubſtance in its natural ſtate, may be ſome defence againſt the cold, which we find makes more impreſſion on lean than on fat perſons. It is for this reaſon, that, to guard themſelves againſt the exceſſive colds of hard winters, and to prevent chilblains, travellers rub the extremities of their bodies, and eſpecially their feet, with ſpirituous oils, ſuch as that of turpentine, &c.

This

This maſs of fat, which makes an univerſal integument of the body, is different from that which is found in the abdomen, thorax, canal of the ſpina dorſi, articulations of the bones, and in the bones themſelves.

But the difference of all theſe particular maſſes of fat conſiſts chiefly in the thickneſs or fineneſs of the pellicles, in the largeneſs or ſmallneſs of the cells, and in the conſiſtence, fluidity, and ſubtilty of the oily matter.

The Nails.

Some authors think the nails are productions of the cutaneous papillae; and others, that they are a continuation of the epidermis. This laſt opinion agrees with experiments made by maceration; by means of which the epidermis may be ſeparated entire from the hands and feet, like a glove or ſock.

In this experiment we ſee the nails part from the papillae, and go along with the epidermis, to which they remain united like a kind of appendix; and yet their ſubſtance and ſtructure appears to be very different from that of the epidermis: But like it, they are alſo inſenſible, and renewable after having been cut, or after having fallen off.

Their ſubſtance is like that of horn, and they are compoſed of ſeveral planes of longitudinal fibres ſoldered together. Theſe ſtrata begin with a ſquare root intermixed with the perioſteum a little before the laſt joints, and end at the extremity of each finger; they are nearly all of an equal thickneſs, but of different lengths.

The external plane or ſtratum is the longeſt, and the reſt decreaſe gradually, the innermoſt being the ſhorteſt; ſo that the nail increaſes in thickneſs from its union with the epidermis where it is thinneſt, to the end of the finger where it is thickeſt.

The

The graduated extremities or roots of all the fibres of which theſe planes conſiſt, are hollowed for the reception of the ſame number of very ſmall oblique papillae, which are continuations of the true ſkin: Which having reached to the root of the nail, forms a ſemilunar fold in which that root is lodged.

After this ſemilunar fold, the ſkin is continued on the whole inner ſurface of the nail, the papillae inſinuating themſelves in the manner already ſaid. The fold of the ſkin is accompanied by the epidermis, to the root of the nail exteriorly, to which it adheres very cloſely.

Three parts are generally diſtinguiſhed in the nail; the root, body, and extremity. The root is white and in form of a creſcent; and the greateſt part of it is hid under the ſemilunar fold already mentioned.

The creſcent and the fold lie in contrary directions to each other. The body of the nail is naturally arched, tranſparent, and appears of the colour of the cutaneous papillae which lie under it. The extremity of the nail does not adhere to any thing, and ſtill continues to grow as often as it is cut.

The principal uſe of the nails is to ſtrengthen the ends of the fingers and toes, and to hinder them from being inverted towards the convex ſide of the hand or foot, when we handle or preſs any thing hard. For in the hand, the ſtrongeſt and moſt frequent impreſſions are made on the ſide of the palm; and in the foot, on the ſole; and therefore the nails ſerve rather for buttreſſes than for ſhields.

The Hairs.

The hairs belong as much to the integuments as the nails. The roots or bulbs lie toward that ſide of the ſkin which is next

next the membrana adipoſa. The trunk or beginning of the ſtem perforates the ſkin, and the reſt of the ſtem advances beyond the outer ſurface of the ſkin to a certain diſtance, which is very various in the different parts of the body.

When the different hairs are examined by a microſcope, we find the roots more or leſs oval, the largeſt extremity being either turned towards, or fixed in, the corpus adipoſum. The ſmalleſt extremity is turned towards the ſkin, and in ſome places it is fixed in the ſkin.

This oval root is covered by a whitiſh ſtrong membrane, in ſome meaſure elaſtic; and it is connected either to the ſkin, to the corpus adipoſum, or to both, by a great number of very fine veſſels and nervous filaments.

Within the root, we obſerve a kind of glue, ſome very fine filaments of which advance toward the ſmall extremity, where they unite and form the ſtem, which paſſes through this ſmall extremity to the ſkin. As the ſtem paſſes through the root, the outer membrane is elongated in form of a tube, which cloſely inveſts the ſtem, and is entirely united to it.

The ſtem having reached the ſurface of the ſkin, pierces the bottom of a ſmall foſſula between the papillae, or ſometimes a particular papilla; and there it meets the epidermis, which ſeems to be inverted round it, and to unite with it entirely. A ſort of unctuous matter tranſudes through the ſides of the foſſula, which is beſtowed on the ſtem, and accompanies it more or leſs, as it runs out from the ſkin in form of an hair.

Hairs are ſcattered almoſt over the whole ſurface of the body, the palms of the hands and ſoles of the feet excepted. They differ in length, thickneſs and ſolidity, in the different parts of the body. Thoſe on the head, are called in Engliſh by the general name of *hairs*; thoſe which are diſ-

poſed

poſed archwiſe above the eyes, *ſupercilia* or the *eye-brows*; thoſe on the edges of the palpebrae, *cilia* or the *eye-laſhes*; and thoſe which ſurround the mouth, and cover the chin, the *beard*. In other parts of the body, they have no particular names; and their different lengths, thickneſſes, &c. in all theſe parts, are ſufficiently known.

Their natural figure ſeems to be rather cylindrical than angular, which is chiefly accidental. Their colour is probably the ſame with that of the glue, or medullary matter of the root, the different conſiſtence of which makes the hairs more or leſs hard, flexible, &c. Their ſtraight or crooked direction muſt depend on that of the holes through which the ſtems paſs. The hairs grow continually, and are renewed again after being cut, by a protruſion of their medullary ſubſtance from the ſkin outwards under a production of the cuticle. When they are deſtitute of this medulla in old people, they dry up, ſplit, and fall off. They ſeem to perſpire through their extremities, and poſſibly through their whole ſurface; as we may conclude from the conſtant force of protruſion in their medulla, which in the plica polonica wants a boundary to terminate it.

The uſe of the hairs, with reſpect to the human body in general, is not ſufficiently known to be determined with certainty. Their uſes with regard to ſome particular parts may be diſcovered; as we ſhall ſee in the deſcription of theſe parts.

The supposed Integuments of the Antients.

Beſides the integuments which have been already deſcribed, the antients reckoned the panniculus carnoſus, and membrana communis muſculorum.

The panniculus carnoſus is found in quadrupeds, but not in men, whoſe cutaneous muſcles are of a very ſmall extent, except that which is called *platyſma myoides* in particular; but

even

even that muſcle cannot in any tolerable ſenſe be reckoned a common integument.

There is no common membrane of the muſcles, which covers the body like an integument; it being no more than particular expanſions of the membranes of ſome muſcles, or aponeurotic expanſions from other muſcles.

The elongations from the lamina of the membrana adipoſa or cellularis, may likewiſe have given riſe to this miſtake, eſpecially in ſuch places where this membrane is cloſely united to the proper membrane of the muſcles.

Feeling.

The ſenſe of feeling is to be underſtood in a two fold manner. Every change of the nerves, produced by the heat, cold, roughneſs, ſmoothneſs, weight, moiſture, or dryneſs of external ſubſtances applied to any part of the body, is commonly called feeling. In this ſenſe, feeling is aſcribed to almoſt all parts of the human body; to ſome more, to others leſs, as in different places of the body the nerves are more numerous and bare, or covered with more tender membranes; and thus even pain, pleaſure, hunger, thirſt, anguiſh, itching, and the other ſenſations, belong to the ſenſe of feeling.

In a more peculiar ſenſe, feeling is ſaid to be the change ariſing in the mind from external bodies applied to the ſkin, more eſpecially at the ends of the fingers. For, by the fingers, we more accurately diſtinguiſh the qualities of tangible ſubſtances than by other parts of our body.

Uses of the Skin, and the Nature of Touch.

The papillae, regularly diſpoſed in the ſpiral folds at the ends of the fingers, on the inſide of the hand, may, by the attention of the mind, become erect or elevated *; and being

* This erection appears from ſhiverings, frights, and from ſimilar erections in the nipples of the breaſts.

ing in this ſtate gently preſſed or rubbed againſt a tangible ſubſtance, they receive an impreſſion from that ſubſtance into their nervous fabric, which is thence conveyed, by the trunks of the nerves, to the brain. This is what we call the *touch*, whereby we become ſenſible chiefly of the roughneſs of objects. Some perſons have this ſenſe ſo acute as to be able to diſtinguiſh colours by touching the ſurface only. By this ſenſe we perceive heat in thoſe bodies which exceed the heat of our fingers; and weight, when the body preſſes more than is uſual. Humidity we judge of by the preſence of water; ſoftneſs, by a yielding of the object; hardneſs, from a yielding of the finger; figure, from the limits, or rough circumſcribed ſurface; diſtance, from a rude calculation or eſtimate made by experience, to which the length of the arm ſerves as a meaſure. The touch ſerves to correct the miſtakes of our other ſenſes; but yet it ſometimes errs itſelf, and then the other ſenſes ſhew themſelves to be true guides to the animal without touch.

The corpus mucoſum moderates the action of the object touched, and preſerves the ſoftneſs and ſound ſtate of the papillae. The cuticle excludes the air from withering and deſtroying the ſkin; qualifies the impreſſions of bodies, ſo that they may be only ſufficient to affect the touch, without cauſing pain. When, therefore, the cuticle is become too thick by uſe, the ſenſe of feeling is either loſt or leſſened; and if it be too thin and ſoft, the touch becomes painful. The hairs ſerve to defend the cuticle from abraſion; to preſerve and increaſe the heat; to cover what ought to be concealed; to render more irritable the membranes of thoſe parts, which ought to be defended againſt the injuries of inſects; and perhaps to exhale ſome uſeleſs vapours, or oil. The nails ſerve to guard the organs of feeling, that the papillae and ends of the fingers may not be ſent back by the reſiſtance of

tangible objects: at the ſame time they increaſe our power of graſping large objects, and aſſiſt in the handling minute objects. In brute animals, they generally ſerve as weapons of offence, and might be of the ſame uſe to man, if they were not cut off.

One moſt important office of the ſkin is, to exhale from the body a large quantity of humours, and to reſorb vapours from the air. The ſkin is replete with an infinite number of ſmall arteries, either coiled up into papillae, or paſſing directly through it, which afford a paſſage to the exhaling vapour; and leſt this vapour ſhould be collected, as it is in ſome diſeaſes, between the ſkin and the cuticle, there are pores in the cuticle correſponding with the exhaling arteries of the ſkin, through which it freely paſſes. Theſe exhaling veſſels or arteries, are eaſily demonſtrated by an injection of water or iſinglaſs into the arteries, which ſweat out from all parts of the ſkin, in an infinite number of ſmall drops; and theſe ſubſtances being transfuſed under the cuticle, rendered impervious by death, raiſe it up into a bliſter.

In a living perſon, this exhalation is many ways demonſtrable. A clean looking glaſs placed againſt the warm and naked ſkin, is quickly obſcured by the moiſt vapour. In ſubterraneous caverns, where the air is more denſe, it is evidently ſeen to fly off from the whole ſurface of the body, like a thick cloud.

Whenever motion of the blood is increaſed, while at the ſame time the ſkin is hot and relaxed, the ſmall cutaneous pores, inſtead of an inviſible vapour, diſcharge *ſweat*, conſiſting of minute but viſible drops, which run together into larger drops by joining with others of the ſame kind. The hotteſt parts are moſt ſubject to ſweat, as the head, breaſt, and foldings of the ſkin. The experiment before mentioned, of injecting the arteries, the ſimplicity of nature, and the apparent cutaneous and pulmonary

monary exhalation, sufficiently persuade us, that the perspirable matter and sweat are discharged through one and the same kind of vessels, and that they differ only by the quantity and celerity of the matter. The humour of the sebaceous glands and the subcutaneous oil, which being more plentifully secreted, and diluted with the arterial juice discharged with the sweat, are of an oily and yellow consistence, and chiefly give smell and colour to the sweat. Hence we find the sweat more fetid in the arm-pits, groins, and other parts, where those glandules are most numerous or abundant. Both blood and small sand have been known to proceed from the skin along with the sweat.

The nature and quality of the perspirable matter may be investigated by experiments, and by considering its analogy to the pulmonary exhalation. What flies off from the lungs in this exhalation is chiefly water, as appears from experiments, by which the breath, being condensed in large vessels, forms into watery drops. That the perspirable matter is also chiefly water, is demonstrated by its obscuring a glass on which it is received; by obstructed perspiration producing a diuresis or diarrhoea; by the speedy passage of liquors drank warm, through the skin if the body be kept warm, or through the kidneys if the body be cold; and by several other phenomena. The water of these vapours is chiefly from what we drink, but is in part supplied from what is inhaled by the skin. The particular smell of the aliments may be sometimes plainly perceived in the perspiration.

That there are, besides water, some volatile alkaline particles, is evident, as well from the nature of our blood, as from the considerable mischiefs which follow an obstructed perspiration. This volatile alkaline matter arises from the finer particles of blood, attenuated by perpetual

tual heat and triture, and changed into an acrimonious nature. Theſe afford the ſcent which is cloſely followed by dogs, who would not know their maſters unleſs ſomething of a peculiar nature perſpired from each perſon.

The quantity of our perſpiring moiſture is very large, whether we conſider the extent of the organ by which it is ſeparated; the abundance of vapours derived from the lungs only; or barely take a review of the experiments made by Sanctorius. This indefatigable man concluded, that five pounds out of eight of the aliment was diſcharged in a healthy perſon by the inſenſible perſpiration alone, independent of the viſible ſweat, and other excretions. In colder climates, the quantity perſpired was found to be $4\frac{2}{7}$ out of 8. But the cutaneous exhalation is even much larger than this; ſince it not only throws off a quantity of the aliment, but likewiſe what is added to the blood by inhalation, which entering, often in a very conſiderable quantity, is thus again expelled. But different diſpoſitions of the air, and of the human body, cauſe great variations in theſe proportions. In warm countries, in the ſummer months, and in young perſons uſing much exerciſe, more goes off from the body by perſpiration, and leſs by the urine; while in cold climates, during the temperate or winter ſeaſons, in aged or inactive perſons, more goes off by the urine than by the inſenſible diſcharge. In temperate countries, making a computation throughout the whole year, ſomething more is perſpired than what paſſes off by urine; and joining together all the experiments made in different countries, both excretions are almoſt alike. The difference of time after feeding alſo in ſome meaſure varies the quantity perſpired; but in general it is moſt copious when the greater part of the digeſted nouriſhment is conveyed into the blood, and there attenuated ſo as to be fit for exhalation. It is naturally diminiſhed in ſleep, even in the warmer climates; but it is increaſed by the heat of bed-cloaths.

In

In general, a plentiful and uniform perſpiration, with ſtrength of body, are good ſigns of health. According to writers on this ſubject, if the perſpiration be increaſed by the weakneſs of the body, it is more hurtful than if it were altogether obſtructed. It is a ſign of health, becauſe it denotes a free pervious diſpoſition of the veſſels throughout the whole body, together with a complete digeſtion of the nouriſhment, the greater part of which is perfectly attenuated into a vapour. When it is diminiſhed, it indicates either a conſtriction of the ſkin, a weakneſs of the heart, or an imperfect digeſtion of the aliments. Perhaps in too great a perſpiration the nervous ſpirits themſelves are evaporated. This diſcharge is, by moderate exerciſe, increaſed to ſix times that of an idle perſon, even to an half or whole pound in an hour. It is likewiſe increaſed by the veſſels being ſtrong and pervious; by warm, watery, and vinous drinks; by animal food of an eaſy digeſtion; by a heavy, temperate, or moderately warm air; and, laſtly, by joy and a tranquil ſtate of mind. The contrary of theſe either leſſen or ſuppreſs the perſpiration; as a thick ſkin, a moiſt air, or a cold and dry one; reſt; more frequent bathing than uſual; a ſupervening diarrhœa; and, laſtly, a diſagreeable nervous affection of the mind. The continuance of life, however, does not depend on a ſcrupulous exactneſs in the quantity of this diſcharge, which is ſo eaſily increaſed or diminiſhed by ſlight cauſes. It is ſhut up by paints in many Indian nations; and it is inconſiderable in many animals without any ſenſible injury. When ſuppreſſed, it is extremely hurtful, producing fevers of the worſt kind. The miſchiefs ariſing from its ſuppreſſion, depend on the putreſcent quality of the particles which are retained along with it.

The ſweat is evidently of a ſaline nature; as appears from the taſte; from the minute cryſtallizations on the cloaths of people who work in glaſs-houſes; and from diſtillation, which

which ſhews the ſweat to be of an alkaline nature. Hence it is, that by this diſcharge the moſt malignant matter of many diſeaſes is thrown off from the body. But, in reality, ſweat is always a preternatural or morbid diſcharge, from which a perſon ought always to be free; unleſs by violent exerciſe, or other accidents, his conſtitution is for a ſhort time thrown into a diſeaſed ſtate. Nor is it unfrequent for ſweats to do conſiderable miſchief in acute diſeaſes, by waſting the watery parts, thickening the reſt of the blood, and at the ſame time rendering the ſalts more acrimonious. By a too violent motion of the blood, the ſweat is rendered extremely fetid; and is ſometimes even red, or mixed with blood itſelf; and being electrified, it is ſometimes lucid.

The uſes of perſpiration are, to free the blood from its redundant water, and throw out thoſe particles, which, by repeated circulations, have become alkaline or otherwiſe acrimonious; and poſſibly to exhale therewith an extremely volatile oil, prepared from the ſame blood. Perſpiration likewiſe qualifies and ſoftens the cuticle, which is a neceſſary medium extended before the tender ſenſible papillae.

Beſides the exhalent veſſels before mentioned, the ſkin is full of ſmall veſſels, which inhale or abſorb thin vapours from the air, either perpetually, or at leaſt when it is not very cold; more eſpecially when the air is damp, the body unexerciſed, the mind oppreſſed with grief, or both under conditions contrary to thoſe which increaſe perſpiration before mentioned. Theſe veins, ſays Haller, are demonſtrated by anatomical injections, which, if thin or watery, ſweat through them as through the arteries: But, according to latter phyſiologiſts, abſorption is performed ſolely by the lymphatic veſſels. That abſorption takes place, is ſufficiently proved by the manifeſt operation of medicines, pervading the air, or applied to the ſkin; ſuch as the vapours of mercury, turpentine, ſaffron, waters of baths, mercurial plaſters, tobacco, coloquintida, opium,

um, cantharides, arsenic, with the fatal effects of contagious or other poisons entering through the skin; as in the venereal infection. Another proof of absorption by the skin is, that animals live in hot moist climates without drink, and yet discharge a considerable quantity of humours both by perspiration and urine. Lastly, absorption has been proved in some diseases where a much greater quantity of urine has been discharged than the quantity of drink taken in. It is difficult to ascertain the quantity of this inhaled matter in animals; in plants it appears, from well-authenticated experiments, to be very considerable, especially during the night.

These cutaneous vessels, both exhaling and inhaling, are capable of contraction and relaxation by the power of the nerves. The truth of this appears from the effects of the passions of the mind; which, if joyful, increase the circulation, and relax the exhaling vessels, so as to yield easier to the impulse of the blood; from whence, with a shortening of the nerves, there follows a redness, moisture, and turgescence of the skin. Those passions, on the contrary, which are sorrowful, and retard the circulation, contract the exhaling vessels; as appears from the dryness and corrugation of the skin, like a *goose-skin*, after frights; and from a diarrhoea being caused by fear. And the same affections seem to open and increase the power of the inhaling vessels, whence the variolous or pestilential contagions are easily contracted by fear.

INDEX.

INDEX

TO VOLUME FIRST.

A.

ABDOMEN, muſcles of, 296
——its bulk, diviſion into regions, contents, arteries, veins, and nerves, xxvii
——muſcles within its cavity, 308
Abductor indicis manus, 345
——indicis pedis, 370
——longus pollicis manus, 343
——minimi digiti manus, ib
——medii digiti pedis, 371
——minimi digiti pedis, ib
——oculi, 272
——pollicis manus, 344
——pollicis pedis, 368
——tertii digiti pedis, 371
Acetabulum of oſſa innominata, 174
Accelerator urinae, 302
Adductor brevis femoris, 350
——ad minimum digitum, 345
——indicis pedis, 370
——longus femoris, 350
——medii digiti pedis, 370
——magnus femoris, 350
——metacarpi minimi digiti manus, 346
——oculi, 272
——pollicis manus, 345
——pollicis pedis, 369
——tertii digiti pedis, 371
Analyſis, chemical, of bones, 16
——phenomenon ſolved by it, ib
Anconaeus, 335
Angularis, 323
Ani ſphincter, 304, 307
——levator, ib ib
Anterior auris, 266
Anterior mallei, 268

Antithenar, 369
Antitragicus, 267
Anus, muſcles of the, 304
Aperiens palpebrarum, rectus, 270
Apophyſes, their different ſorts and uſes, 24. Many ſo named are epiphyſes, ib
Appendices. *Vid.* Epiphyſes.
Arm-bone, *Vid.* Humeri os.
Arthrodia, 37, 40
Articulations, 36. Symphyſis, ſynarthroſis, ib. Diarthroſis, 37. Diſpute concerning them, 38
Arytenoideus minor, 295
Arytenoideus major, ib
—— Obliquus, ib
—— tranſverſas, ib
Arytaeno epiglottideus, ib
Aſtragalus, 243. Its articulation, 245. Of children, 247
Atlas, or the firſt vertebra of the neck, 147. Of infants, 149
Attollens aurem, 265
—— oculi, 271
—— elevator oculi, ib
Auris anterior, 266
—— retrahentis, ib
—— poſterior, ib
—— ſuperior, ib
—— tranſverſus, 267
—— obliquus, 268
Axis, or third vertebra colli, 153
Axygos uvulae, 291

B.

Baſio-cerato-chondro-gloſſus, 284
Biceps brachii, 334
—— flexor cubiti, 333
—— flexor cruris, 359
—— internus, 334
—— externus, 335
Biventer maxillae inferioris, 283
Bones, what, 28. Their plates, 29. Fibres, claviculi, ib. Cancelli, 10. Arteries, 11. Veins, 12. Nerves, 13. Circulation of their liquors, nouriſhment, increaſe, decreaſe, 14. Phenomena and diſeaſes, 14. 19. 20. 31. Tranſverſe and longitudinal canals, 15. Analyſis, 16. Perioſteum internum, 17. Marrow, 18. Diſtinguiſhed into broad and round, 22. Strength increaſed by being hollow, 23. Proceſſes, 24. Cavities, 25. Epiphyſes, 27. Oſſification, 27. Articulations, 36
Brachialis, 334
—— internus, ib
—— externus, 335
Breaſt bone. *Vid.* Sternum

Bregma, 75
Broad-bones, their ſtructure, 22
Buccinator, 277
Bulbo-cavernoſus, 303

C.

Calcis os, 245. Of children, 247
Cancelli of bones, 10. Uſes, 11
Canini dentes, 131
Caninus, 275
Capitis obliquus ſuperior, 324
——obliquus inferior, ib
——par terium, 323
——rectus poſticus minor, 326
——major, 325
Carpus conſiſts of eight bones, 213. Its figure, ſubſtance, articulation, motions, 219. Uſes, ib. Of infants, ib
Cartilages, what, their plates, fibres, &c. 44. How prevented from oſſifying, 46. Their uſes, diſeaſes, ib
Cavities of bones, their different kinds, 25. Uſes, 26
Cephalo-pharyngeus, 293
Cerato-gloſſus, 284
Cervical vertebrae, their diſtinguiſhing marks, 145. Atlas, 147. Dentata, 149. Axis, 153. Seventh, ib
Cervicalis deſcendens, 322
Cheeks, their arteries, veins, and nerves, 20
Cheek-bones, *Vid.* Malarum oſſa
Circumflexus, or tenſor palati, 288
Clavicles, 192. Internal end, 193. Body, 194. External end, medullary veſſels, ſubſtance, articulation, ib. Of children, 195. Uſes, ib
Claviculi of bones, perpendicular, oblique headed, crooked, 9
Cleido-maſtoideus, 282
Coccygeus, 308
Coccygis os, 163. Its firſt, ſecond, third, and fourth bones, 164. Subſtance, motion, diſeaſes, uſes, 165
Collar-bone. *Vid.* Clavicle.
Colli latiſſimus, 282
——longus, 316
——tranſverſalis, 325
——ſemiſpinalis, ib
——interſpinalis, 328
——intertranſverſalis, ib
Complexus, 322
——minor, 323
Compreſſor naris, 278
Conſtrictor cunni, 306
——oris, 278
——iſthmi faucium, 299
——pharyngis, 293

Coraco-brachialis, 332
——hyoideus, 286
——radialis, 334
Coronal future, 57
Corpus mucoſum, 466
Corrugator ſupercilii, 265
Coſtae. *Vid.* Ribs.
Cranium, its figure, 53. Surfaces, 55. Tables, diploe, 56. Conſiſts of ſix proper, ib. and two common bones, 57. Its ſutures, ib. Muſcles of its integuments, 264.
——its contents, arteries, veins, and nerves, xxv
Cremaſter, 302
Crico-arytenoideus lateralis, 294
Crico-arytenoideus poſticus, ib.
Crico-thyroideus, 286
Crico-pharyngeus, 292
Crotaphyte muſcle, 280
Cruralis, 358
Cubit, or fore-arm. *Vid.* Fore-arm.
Cubitalis Riolani, 335
Cuboides os, 248. At the birth, ib.
Cucularis, 318
Cuneiforme os of the wriſt, 215
Cuneiformia oſſa of the foot, ; externum, 248. Medium internum, 249. At the birth, 250
Cutaneous, 281
Cuticula, 464
Cutis vera, 467. Texture, ib. Papillae, 468. Appearance in inflammation, 469. How affected by the paſſions, 486. Glands, why called ſebaceous, 471. Uſes of the ſkin, ib. See nature of Touch.

D.

Dartos, 301
Deltoides, 331
Dentata, or ſecond vertebra of the neck, 149. Its diſtinguiſhing marks, ib. Of infants, 150
Dentes. *Vid.* Teeth.
Deprimens, 272
Depreſſor anguli oris, 276
——alae naſi, 275
——labii inferioris, 276
——labii ſuperioris alaeque naſi, 275
——labii ſuperioris proprius, 276
——labiorum communis, ib.
——oculi, 272
Diaphragma, 308, 309
Diarthroſis, 37. Its three ſpecies, enarthroſis, arthrodia, ib. Ginglimus, 38
Digaſtricus, 283

Digiti. *Vid.* Fingers and Toes.
Diploe of the skull, 56
Dorsal vertebrae, their distinguishing structure, 153. Wherein they differ from each other, 155.
Dorsi interspinalis, 328
——latissimus, 318
——spinalis, 321
——longissimus, ib
——semispinalis, 324

E.

Ear, muscles of the, 265
——external parts enumerated, xxi. Arteries, veins, and nerves, xxiii
Ejaculator seminis, 302
Elasticity of muscular fibres distinguished from irritability, sensibility, and vital power, 389
Elevator oculi, 271
——labii superioris, 275
——labii inferioris, 277
——labiorum communis, 275
Enarthrosis, 37
Epicranius, 264
Epiphyses, 26. Several have apophyses, 27. Their uses, ib.
Epistaphilinus, 292
Erector clitoridis, 305
——penis, 302
Ethmoidal suture, 61
Ethmoides os, 89. Its cribriform plate, ib. Nasal plate, 90. Cellulae, ossa spongiosa, 91. Connection, uses, 92. Morbid phenomena, ib. Of a child, ib.
Extensor brevis digitorum pedis, 365
——carpi radialis brevior, 338
——carpi radialis longior, 337
——carpi ulnaris, 338
——digitorum communis, 340
——longus digitorum pedis, 364
——magnus, 361
——minor pollicis manus, 344
——ossis metacarpi pollicis manus, 343
——pollicis primus, 344
——pollicis secundus, ib.
——primi internodii pollicis manus, 343
——proprius pollicis pedis, 367
——secundi internodii pollicis manus, 344
——major pollicis manus, ib.
——pollicis secundus, ib.
——tertii internodii, ib.
——tarsi suralis, 361
——tarsi minor, ib.

Externus mallei, 268
Extremities, inferior, conſiſt each of thigh, 227; leg, 232; foot, 242; muſcles of, 349
——ſuperior, conſiſt each of ſhoulder, 192; arm, 201; fore-arm, 206; hand, 212. Uſes, 226. Muſcles of, 329
Extremities, upper parts enumerated, with their arteries, veins, nerves, and muſcles, xxx
Extremities, lower, ſame parts enumerated, xxxi
Eye, external and internal parts enumerated, xxi. Its arteries, veins, and nerves, enumerated, xxiii
Eye-ball, muſcles of, 271
Eye-lids, muſcles of, 269

F.

Face, 101. Compoſed of upper-jaw, ib. 108. Lower-jaw, 121
——Its arteries, veins, and nerves, enumerated, xxii
Fallopii capitis par tertium, 323
Faſciae latae, 355
Fat, how formed, its difference in quantity and quality, 473
Fauces, muſcles about the, 290
Faucium iſthmi conſtrictor, ib.
Female ſkeleton, diſtinguiſhing marks of it, 256
Femoris os, 227. Its upper end, ib. Body, 229. Lower end, 230. Of infants, its articulation and motions, 231
Fibres, their fabric and ſtructure, 374
Fibula, its upper end, 235. Body, 236. Inferior end, 238. Its articulation, uſes, 139. Of infants, ib.
Fingers, their bones, phalanges. firſt, ſecond, third, 223, 225
Flexor acceſſorius digitorum pedis, 366
——brevis digitorum pedis, 365
——brevis minimi digiti pedis, 369
——brevis pollicis manus, 342
——brevis pollicis pedis, 368
——longus digitorum pedis, 366
——primi internodii, 343
——tertii internodii, 342
——ſecundi internodii, 343
——carpi radialis, 337
——carpi ulnaris, ib.
——longus pollicis manus, 342
——longus pollicis pedis, 368
——oſſis metacarpi pollicis, 343
——parvus minimi digiti manus, 346
——profundus perforans, 339
——ſublimis perforatus, 338
Fontanelle, 75
Foot, compoſed of tarſus, 243; metatarſus, 251; toes, 253. Its motion, 245. Muſcles on it, 367
Fore-arm, 206. Conſiſts of ulna, ib. Radius, 219. Muſcles on it, 335

Frontal-bone, 64. Its external ſurface, ib. Proceſſes, 65. Cavities, ib. Foramina, 66. Internal ſarface, 67. Foramen, 68. Subſtance, ſinuſes, 69. Connection, 71. Uſes, 72. Of children, ib.
Frontalis, 264———

G.

Gaſtrocnemius externus, 360
———internus, ib.
Gemellus, ib.
Gemini, 353
Genae quadratus, 282
Generation, organs of; muſcles about them, 301—305
Genio-hyo-gloſſus, 284
Genio hyoideus, ib.
Ginglimus, its ſpecies, 38, 40
Gloſſo-ſtaphilinus, 290
——pharyngeus, 293
Glottis, muſcles ſituated about it, 294
Gluteus maximus, 351
——— medius, 352
——— minimus, 353
Gomphoſis, 37
Gracilis, 356
———internus, ib

H

Hairs, 477. How formed, their texture and uſe, 477, 478
Hand, 212. Conſiſts of carpus, 213. Metacarpus, 220. Fingers, 123. Its muſcles, 342
Harmonia, 37
Haunch bones. *Vid.* ilium oſſa.
Head, conſiſts of cranium, 53. Face, 101. Its motions, 151
Heel-bone. *Vid.* Calcis os.
Helicis major, 266
———minor, 267
Hip-bone. *Vid.* os iſchium.
Humeri os, 201. Its upper end, ib. Body, 203. Lower end, ib. Subſtance, 204. Articulation, 205. Motion, ib. Of children, ib. Muſcles ſituated on, 333
Hyoides os, 135. Its body, 136. Cornus, ib. Appendices, ib. Ligaments, ſubſtance, connection, uſes, of children, 137. Muſcles about it, 283. 285. 287.
Hyo-gloſſus, 284
Hyo-pharyngeus, 293
Hyo-thyroideus, 286
Hypothenar minor, 346
———Riolani, ib.

I.

Jaw, lower. *Vid.* Maxilla inferior.
——upper. ——Maxilla ſuperior.

Iliacus internus, 312
——externus, 353
Ilium ossa, 167. Spine, dorsum, ib. External surface, 168. Interior surface, 169. Medullary vessels, substance, of a child, 170
Incisivus inferior, 277
——lateralis, 275
——medius, 276
Incisores dentes, 130
Indicator, 345
Inferior prostrate, 303
Infraspinatus, 330
——costales, 315
Innominata ossa, 166. Composed of os ilium, 167. Ischium, 170. Pubis, 172. Their great foramen, 173. Acetabulum, 174. In infants, 170. Connection, phenomena, 175. Uses, 176
Intercostales externi, 314
——interni, ib.
Internus auris, 268
——mallei, ib.
Interosseus auricularis, 347
Interossei pedis externi, 370
——pedis interni, 371
Interspinales colli, 328
——dorsi et lumborum, ib.
Intertransversales colli, ib.
——dorsi, ib.
——lumborum, 329
Irritability, 389
Ischium os, 170. Process, depressions, tuber, 171. Substance, ib. Of infants, 172
Ischio cavernosus, 302
Isthmi faucium constrictor, 290

L.

Lambdoid suture, 57. Its additaments, 58
Latissimus colli, 282
——dorsi, 318
——Lattice-work of bones. *Vid.* Cancelli.
Laxator tympani, 267
Leg, composed of tibia, 232. Fibula, 235. Rotula, 239. Its articulation and motions, 241
Levator palati, 289
——palati mollis, ib.
——parvus mollis, 303
——menti, 277
——palpebrae superioris, 270
——oculi, 271
——anguli oris, 274
——labii superioris alaeque nasi, 275

Levator labii inferioris, 277
—— ani, 304. 307
—— scapulæ, 323
Ligamenta lateralia, 430
—— cervicale seu nuchæ, ib.
—— transversum vertebræ primæ colli, 434
—— anticum commune vertebrarum, ib.
—— intervertebralia, 435
—— interspinosa, ib.
—— intertransversalia, ib.
—— capsularia, ib.
Ligaments, 42. Their fibres, ib. Vessels, nerves, diseases, 43. Uses, phenomena, 44. Enumeration of them, 428
—— of the elbow joint, 442
—— joining the heads of the radius and ulna, 444
—— of the fore-arm, 445. Carpus, 446. Carpus and metacarpus, ib. Of the fingers, 447
—— of the pelvis, 448
—— of the hip-joint, 453
—— of the knee, 456
—— of the fibula, 458
—— at the bottom of the fibula and tibia, 459
—— of the toes, 460
Ligamentum capsulare, 429
—— latum anterius, 430
—— latum posterius, ib
—— processus odontoides seu perpendiculare, ib.
—— posticum commune vertebrarum, 435
—— cartilaginis ensiformis, 437
—— costarum verarum, ib.
—— costarum ipsarum proprium, 438
—— interclaviculare, 440
—— rhomboideum, ib.
—— trapezoidæum scapulæ, 441
—— conoidæum, ib.
—— proprium anticum, ib.
—— proprium posticum, ib.
—— capsulare humeri, ib.
Lingualis, 285
Longus colli, 316
Longissimus dorsi, 321
Lumbar vertebræ, their particular structure, 156. Differences, 157
Lumbaris internus, 311
—— externus, ib.
Lumbricales, 339
—— pedis, 362
Lunare os, 214

M.

Magnum os of the wriſt, 216
Malarum oſſa, their proceſſes, 106. Holes, ſubſtance, connection, 107. Of infants, 108
Marrow, what, 18. Its chemical analyſis, its arteries, veins, nerves, ib. Tranſverſe and longitudinal canals for conveying it, 14. 20. Uſes, ib. Phenomena and diſeaſes, 21
Maxilla inferior, 121. The chin, ſides, baſe, ib. Phenomena, angles, proceſſes, 122. Foramina, ſubſtance, 123. Articulation, 124. Motions, ib. Of infants, uſes, 125.
—— ſuperior, conſiſts of 13 bones, 101. Their connection, 102. 113.
Maxillaria oſſa, their proceſſes, 108. Cavities, 109. Foramina, 110. Sinuſes, 111. Morbid phenomena, 112. Subſtance, connection, 113. Uſes, ib. Of children, 114.
Marſupialis, 308
Maſtoideus, 282
—— lateralis, 323
Maſſeter, 280
Membrana adipoſa, 472. Texture, its different affections, 473
Metacarpus of the hand, compoſed of four bones, 220. Subſtance at the Birth, articulation, ib. Figure, uſes, digiti indicis, ib. Medii, 221. Annularis, minimi, 222
Metacarpus muſcle, 346
Metatarſeus, 369
Metatarſus, compoſed of five bones, 151
Molares, dentes, 131
Mouth, external and internal parts enumerated, xxi
—— its arteries, veins, and nerves, xxiv
Mucilaginous glands, 49. Liquor ſecreted by them, 50. Their cellular ſubſtance, veſſels, nerves, diſeaſes, 51. Uſes, 50. Phenomena, ib.
Multifidus ſpinæ, 324
Muſcular motion, 373
Muſcles, tables of them, 397. 425
Muſculus cutaneus, 281
—— ſupercilii, 265
—— frontales, ib.
—— ſtapedii, 268
—— tubæ novus, 288
—— patientiæ, 323
Mylo pharyngeus, 293
—— hyoideus, 283

N.

Nails, 476. How formed, their texture and uſe, ib.
Naſalis, 274
Naſi oſſa, their ſides, 104. Connection, ib. Uſes, of an infant, 105
Naviculare os, of the tarſus, 247. Of children, 248

Neck. *Vide* Cervical vertebra.
Neck, its muſcles, arteries, veins, and nerves, enumerated, xxv.
Noſe, external and internal parts enumerated, xxi. Arteries and nerves, xxiii.

O.

Obliquus inferior, 273. 326
—— auris, 268
—— major, 273. 326
—— minor, 326. 273
—— externus abdominis, 298
—— deſcendens, ib.
—— internus abdominis, 299
—— aſcendens, 298
—— ſuperior, 326. 372
—— deſcendens externus, 296
—— internus, 299
—— capitis ſuperior, 326
—— capitis inferior, ib.
Obturator internus, 307
—— externus, 351
Occipito-frontalis, 264
Occipitalis, ib.
Occipitis os, 82. Its external ſurface, 93. Proceſſes, ib. Internal ſurface, 85. Holes, 86. Subſtance, 87. Connection, 88. Uſes, in infants, ib.
Omo-hyoideus, 285
Orbicularis palpebrarum, 269
—— oris, 278
Oſſification of bones, 27. Accounted for from particular diſpoſition of veſſels, and from preſſure, 29. Phenomena, 27. 31. Its hiſtory, 32
Oſteogeny, its hiſtory, 34. Of what uſe, 35

P.

Palati os, 114. Its Square palate plate, ib. Pterygoid proceſs, naſal lamella, 115. Orbitar proceſs, 116. Subſtance, connection, uſes, ib. Of children, 117. Its diſeaſes, ib.
Palato-ſalpingeus, 288
—— ſtaphilinus, 292
—— pharyngeus, 290
Palmaris longus, 336
—— brevis, ib.
—— cutaneus, ib.
Parathenar major, 369.
—— minor, 370
Parietal bones, their external ſurface, 72. Internal ſurface, 73. Subſtance, Connection, 74. Uſes, 75. Of children, ib. Practical obſervations concerning bregma, ib.
Patella. *Vide* Rotula.

Pelvis,

Pelvis, 166. 177. Confiſts of os ſacrum, 154. Coccygis, 163. Oſſa innominata, 166. Its cartilages, ligaments, &c. 448.
Perichondrium, its veſſels, 45
Perioſteum externum, 5. Its fibres, ib. Arteries, veins, nerves, 7. Uſes, 8. Diſeaſes, ib.
Perioſteum internum, its ſtructure, 17. Uſes, ib.
Perſpiration, what, 482. Proofs of, 483. Mixed with alkaline particles, 484. Quantity, ib. Uſes, 485
Petro-ſalpingo-ſtaphilinus, 289
Pectoralis, 313
—— major, 312
—— minor, 313
Pectinalis, 349
Pectineus, 350
Perforatus, 338. 365
Perforans, 339
Peroneus maximus, 363
—— poſterior, ib
—— primus, ib.
—— medius, 364
—— anticus, ib.
—— ſecundus, ib.
—— longus, 363
—— brevis, ib.
Phalanges of the fingers, 225
—— toes, 253
Piſiforme os of the wriſt, 215
Plates of bones, their fibres, 9
Platiſma myoides, 281
Plantaris, 361
Pollicis manus oſſa. *Vide* Thumb.
—— pedis oſſa. *Vide* Toes.
Poplitaeus, 359
Pores, tranſverſe and longitudinal of bones, 15. Their uſes, ib.
Poſterior auris, 266
—— indicis, 347
—— annularis, 348
—— medii, ib.
Proceſſes of bones. *Vide* Apophyſes.
Profundus, 339
Pronator radii teres, 341
—— quadratus, ib.
Prior indicis, 347
—— annularis, ib.
—— medii, 48
Pſoas, 311
—— parvus, ib.
—— magnus, ib. 349
Pterygoideus major, 280

Ptery-

Pterygoideus minor, 281
—— internus, 280
—— externus, 280
Pterygo-ſtaphilinus externus vulgo, 280
—— pharyngius, 293
Pubis os, 172. Subſtance, of children, 173
Pyramidalis, 275. 301
Pyriformis, 353

Q.

Quadratus, 276
—— genæ, 282
Quadratus, lumborum, 311
—— femoris, 354

R.

Radius, its upper end, 209. Body, ib. Lower end, 210. Of children, articulation, 211
Radialis internus, 337
—— externus longior, 338
—— externus primus, ib.
—— externus brevior, ib.
—— ſecundus, 335
Rectus abdominus, 300
—— major, 326
—— anterior longus, 317
—— anterior brevis, ib.
—— internus, 356
Rectus, ib.
—— capitis internus major, 316
—— capitis internus minor, 317
—— capitis lateralis, ib.
—— capitis poſticus major, 325
—— capitis poſticus minor, 326
Retrahentes auris, 266
Retractor anguli oris, 277
Rhomboideus, 319
—— major, ib.
—— minor, ib.
Ribs, 178. Their middle, ib. Poſterior end, 179. Anterior end, ſubſtance, cartilages, 180. Articulation and motion, 181. Differ from each other, 182. Diſtinguiſhed into true and falſe, ib. Diſtinguiſhing marks of the 1ſt, 2d, 3d, 4th, 5th, 6th, 7th, 11th, 12th, ib. Of infants, 185. Motion, ib. Their cartilages, ligaments, &c. 431
Rinaeus vel naſalis, 274
Rotula deſcribed, 239. Its ſubſtance, 240. Articulation, ib. Of children, 241
Round bones, their ſtructure, 22. Strength, how increaſed, ib.

S.

Sacro lumbalis, 322

Sacrum os, its body, 159. Proceſſes oblique tranſverſe, ib. Spinal, 161. Foramina, 162. Subſtance, ib. Articulation, 163. Uſes, ib. Of children, ib.
Sagittal ſuture, 59
Salpingo-ſtaphilinus, 289
—— ſtaphilinus internus, ib.
—— pharyngeus, 291
Sartorius, 356
Scalenus prior, 327
—— ſecond, ib.
—— third, ib.
—— anticus, ib.
—— medius, ib.
—— poſticus, ib.
Scaphoid bone of the wriſt, 214
Scapula, 195. Its baſe, 196. Inferior coſta, 197. Superior coſta, dorſum, ib Anterior ſurface, 198. Proceſſes, ſpine, acromion, caracoid proceſs, ib. Third proceſs, medullary veſſels, 199. Subſtance, ib. Articulation, 200. Uſes, 201. Of children, ib.
Schindyleſis, 37
Semiſpinalis, 324 325
—— internus, 324
—— externus, ib.
Semi-interoſſeus, 345
Semi-membranoſus, 358
Semi-nervoſus, ib.
Semi-orbicularis, 278
Semi-tendinoſus, 358
Serratus anticus, 313
Senſibility, 389
—— magnus, ib.
—— poſticus inferior, 319
—— poſticus ſuperior, 320
Seſamoid bones, how formed, 254
Where to be found, ib.
Share-bone. *Vide* oſſa pubis.
Shoulder, conſiſts of clavicle, 192, and ſcapula, 195.
Shoulder blade. *Vide* Scapula.
Sinus frontal, 69. Sphenoidal, 93. Maxillary, 112.
Skeleton, natural, artificial, 52. Situation of the bones, 62. Divided into head, 53, trunk, 137. Extremities, ſuperior, 192. Inferior, 226. Table of its bones, 397. 412
Skeleton of a female, its differences from that of the male, 256.
Skull. *Vide* Cranium.
Soleus, 360
Sphenoid bone, 93. Its external ſurfaces, proceſſes, ib. Cavities, 95. Holes proper, 96 common, 98. Sinuſes, 99. Subſtance, connection, ib. Uſes, 100. Of children, ib.
Sphenoidal ſuture, 61

Spheno-ſalpingo-ſtaphilinus, 288
———ſtaphilinus, 289
Sphincter ani, 304. 307
———vaginae, 306
———labiorum, 278
———externus, 304
———cutaneus, ib
Spinalis, 325
———cervices, ib
——— dorſi, 321
Spine, its ſhape, 138. Conſiſts of true vertebrae, ib. and falſe, 159
Splenius, 320
Spongioſa inferiora oſſa, 117. Their ſubſtance, connection, uſes; of children, 118
———ſuperiora oſſa, 91
Squamous ſutures, 59. How formed, 60
Staphilinus, 292
———externus, 288
Stapedius, 269
Sterno-maſtoideus, 282
——cleido-maſtoideus, ib
——hyoideus, 285
——thyroideus, 286
——coſtalis, 315
Sternum, 186. Its ſubſtance, 187. Firſt bone, ib. Second bone, 188. Xiphoid cartilage, ib. Connection, 190. Of children, ib. Uſes, ib. Motion, ib. Its cartilages, ligaments, &c. 437
Stylo-gloſſus, 289
——hyoideus, ib
——pharyngeus, 288
Subclavius, 313
Subſcapularis, 332
Sublimis, 339. 366
Superior auris, 266
Supinator longus, 341
——radii longus, 340
——brevis, 341
Supra coſtales, 315
Supra ſpinatus, 329
Suture, 36. How formed, 37
Sutures of the cranium, 57. Coronal, lambdoid, ib. Sagittal, 58. Squamous, 59. Ethmoidal, ſphenoidal, tranſverſe, 61. Zygomatic, 62. Uſes, 63
Sutures of the face diſtinguiſhed by names, 102. Uſes, 103
Symphyſis, divided into ſynchondroſis, ſynneuroſis, ſyſſarcoſis, 36

Synarthrosis, divided into future, harmonia, 37. Gomphosis, schindylesis, ib
Synchondrosis, 36
Syndesmo pharyngaeus, 293
Syneurosis, 36
Synovia of joints, what composed of, 48. Its uses, diseases, 49
Syssarcosis, 36

T.

Tarsus consists of seven bones, 243. Its articulation, uses, 250.
Teeth, 126. Their base, ib. Roots, ib. Periosteum, ib. Substance, 127. Canals, ib. Vessels, 128. Formation, 129. Shedding, ib. Connection, 130. Uses, ib. Distinguished into incisores, ib. Canini, molares, 131. Phenomena, 133
Temporal bones, 76. Their external surface, processes, ib. Cavities, 78. Holes, ib. Internal surface, 80. Substance, 81. Connection, 82. Uses, of infants, ib
Temporalis, 279
Tensor tympani, 268
——palati, 288
——vaginae femoris, 355
Teres minor, 330
——major, 331
Thenar, 369.—345
Thigh-bone. *Vide* Femoris os.
Thorax, 177. Consists of vertebræ, 138. ribs, 178. sternum, 186. Its motion, 196
——its division, muscles, contents, arteries, veins, and nerves xxvi
Thumb, consists of three bones; the first, its articulation, 223. At the time of birth, 224. Second bone, its articulation, ib. Third bone, ib
Thyro-staphilinus, 291
—— pharyngo-staphilinus, ib.
——pharyngeus, 292
Thyreo-arytænoideus, 294
——epiglottideus, 295
——hyoidæus, 286
Tibia, its upper end, cartilages, 232. Body, 233. Lower end, 234. Articulation; of children, 235
Tibialis gracilis, 361
——anticus, 362
——posticus, ib
Toes, 253. Differ from the fingers; use, of children, ib
Touch, what, 480. How performed by the cutaneous papillæ, 481
Transverse future, 61
Trachelo-mastoideus, 323
Tragicus, 267

Transverſalis, 299
——colli, 325
——pedis, 372
——urethrae, 303
——lumborum, vulgo ſacer, 325
——dorſi, ib
——colli, ib
Transverſo ſpinalis dorſi, 324
——lumborum, veterib. ſacer, ib
——ſpinalis colli pars interna, 325
——ſpinalis colli pars externa, ib
Transverſus abdominis, 300
——auris, 267
——perinei, 303. 306
Trapezium os, 215
Trapezius, 318
Trapezoides os, 216
Triangularis, 276, 315
Triceps minor, 350
——ſecundus, ib
——tertius, 351
——extenſor cubiti, 334
——adductor femoris, 350
——Triquetra oſſa, 58
Trochlearis, 272
Trochoides ginglimus, 41
Trunk of the ſkeleton, conſiſts of the ſpine, pelvis, thorax, 177
Turbinata oſſa. *Vid.* Spongioſa.

U.

Vaſtus externus, 357
——internus, ib
Vertebrae cervical. *Vid.* Cervical.
——dorſal. *Vid.* Dorſal.
——falſe, compoſed of os ſacrum, 155. Coccigis, 163
——lumbar. *Vid.* Lumbar.
——true, 138. Their bodies, cartilages, ib. Articulations, phenomena, 141. Of infants, 143. Diſeaſes, 144. Divided into cervical, 145. Dorſal, 153. Lumbar, 166. Uſes, ſecured from luxations, 158
Veſſels of perioſteum, 7. Of the bones, 11. Of the marrow, 19. Of ligaments, 42. Cartilages, 45. Mucilaginous glands, 49.
Vires animales, divided into elaſticity, ſenſibility, and irritability, 389
Vis nervea, 381. Inſita, ib. How produced and influenced, 382. Dr Monro's objections to the vis inſita, 392
Vital power deſcribed, ib
Vomer, 118. Its figure, ſituation, 119. Subſtance, connection, uſes, 120

Ulna, its upper end, 206. Body, 208. Lower end, ib. Articulation, 209. Of children, ib
Ulnaris gracilis, 336
——internus, 357
——externus, 338

W

Women, the difference of their bones from thoſe of men, 256
Wormiana oſſa. *Vid.* Triquetra.
Wriſt. *Vid.* Carpus.

X.

Xiphoid cartilage, 188. Its diſeaſes, ib

Z.

Zygoma, 77. 107
Zygomatic ſuture, 62
Zygomaticus major, 277
——minor, 278

DIRECTIONS

DIRECTIONS to the BINDER for placing the PLATES.

Vol. I. Table I.	to face page	397
II.	———	401
III.	———	404
IV.	———	407
V.	———	412
VI.	———	414
VII.	———	416
VIII.	———	420
IX.	———	425
X.	———	427
Vol. II. Table XI.	———	458
XII.	———	459
XIII.	———	460
XIV.	———	462
Vol. III. Table XV.	———	189
XVI.	———	191
XVII.	———	202
XVIII.	———	218
XIX.	———	311
XX.	———	312

END OF THE FIRST VOLUME.

I
Edinburgh
1801
v.1